

**EXISTENCE AND UNIQUENESS FOR ONE-PHASE STEFAN  
PROBLEMS OF NON-CLASSICAL HEAT EQUATIONS WITH  
TEMPERATURE BOUNDARY CONDITION AT A FIXED FACE**

ADRIANA C. BRIOZZO, DOMINGO A. TARZIA

ABSTRACT. We prove the existence and uniqueness, local in time, of a solution for a one-phase Stefan problem of a non-classical heat equation for a semi-infinite material with temperature boundary condition at the fixed face. We use the Friedman-Rubinstein integral representation method and the Banach contraction theorem in order to solve an equivalent system of two Volterra integral equations.

1. INTRODUCTION

The one-phase Stefan problem for a semi-infinite material for the classical heat equation requires the determination of the temperature distribution  $u$  of the liquid phase (melting problem) or of the solid phase (solidification problem), and the evolution of the free boundary  $x = s(t)$ . Phase-change problems appear frequently in industrial processes and other problems of technological interest [2, 3, 6, 8, 9, 10, 11, 12, 18, 29]. A large bibliography on the subject was given in [25].

Non-classical heat conduction problem for a semi-infinite material was studied in [4, 7, 17, 27, 28], e.g. problems of the type

$$\begin{aligned}u_t - u_{xx} &= -F(u_x(0, t)), & x > 0, t > 0, \\u(0, t) &= 0, & t > 0 \\u(x, 0) &= h(x), & x > 0\end{aligned}\tag{1.1}$$

where  $h(x), x > 0$ , and  $F(V), V \in \mathbb{R}$ , are continuous functions. The function  $F$ , henceforth referred as control function, is assumed to satisfy the condition

$$(H1) \quad F(0) = 0.$$

As observed in [27, 28], the heat flux  $w(x, t) = u_x(x, t)$  for problem (1.1) satisfies a classical heat conduction problem with a nonlinear convective condition at  $x = 0$ ,

---

2000 *Mathematics Subject Classification*. 35R35, 80A22, 35C05, 35K20, 35K55, 45G15, 35C15.

*Key words and phrases*. Stefan problem; non-classical heat equation; free boundary problem; similarity solution; nonlinear heat sources; Volterra integral equations.

©2006 Texas State University - San Marcos.

Submitted November 10, 2005. Published February 9, 2006.

which can be written in the form

$$\begin{aligned} w_t - w_{xx} &= 0, & x > 0, t > 0, \\ w_x(0, t) &= F(w(0, t)), & t > 0, \\ w(x, 0) &= h'(x) \geq 0, & x > 0. \end{aligned} \tag{1.2}$$

The literature concerning problem (1.2) has increased rapidly since the publication of the papers [19, 21, 22]. Related problems have been also studied; see for example [1, 14, 16]. In [26], a one-phase Stefan problem for a non-classical heat equation for a semi-infinite material was presented. There the free boundary problem consists in determining the temperature  $u = u(x, t)$  and the free boundary  $x = s(t)$  with a control function  $F$  which depends on the evolution of the heat flux at the extremum  $x = 0$  is given by the conditions

$$\begin{aligned} u_t - u_{xx} &= -F(u_x(0, t)), & 0 < x < s(t), 0 < t < T, \\ u(0, t) &= f(t) \geq 0, & 0 < t < T, \\ u(s(t), t) &= 0, & u_x(s(t), t) = -\dot{s}(t), & 0 < t < T, \\ u(x, 0) &= h(x) \geq 0, & 0 \leq x \leq b = s(0) \quad (b > 0). \end{aligned} \tag{1.3}$$

The goal in this paper is to prove the existence and uniqueness, local in time, of a solution to the one-phase Stefan problem (1.3) for a non-classical heat equation with temperature boundary condition at the fixed face  $x = 0$ . First, we prove that problem (1.3) is equivalent to a system of two Volterra integral equations (2.4)-(2.5) following the Friedman-Rubinstein's method given in [13, 23]. Then, we prove that the problem (2.4)-(2.5) has a unique local solution by using the Banach contraction theorem.

## 2. EXISTENCE AND UNIQUENESS OF SOLUTIONS

We have the following equivalence for the existence of solutions to the non-classical free boundary problem (1.3).

**Theorem 2.1.** *The solution of the free-boundary problem (1.3) is*

$$\begin{aligned} u(x, t) &= \int_0^b G(x, t; \xi, 0)h(\xi)d\xi + \int_0^t G_\xi(x, t; 0, \tau)f(\tau) d\tau \\ &+ \int_0^t G(x, t; s(\tau), \tau)v(\tau) d\tau - \iint_{D(t)} G(x, t; \xi, \tau)F(V(\tau))d\xi d\tau, \end{aligned} \tag{2.1}$$

$$s(t) = b - \int_0^t v(\tau) d\tau, \tag{2.2}$$

where  $D(t) = \{(x, \tau) : 0 < x < s(\tau), 0 < \tau < t\}$ , with  $f \in C^1[0, T]$ ,  $h \in C^1[0, b]$ ,  $h(b) = 0$ ,  $h(0) = f(0)$ ,  $F$  is a Lipschitz function over  $C^0[0, T]$ , and the functions  $v \in C^0[0, T]$ ,  $V \in C^0[0, T]$  defined by

$$v(t) = u_x(s(t), t), \quad V(t) = u_x(0, t) \tag{2.3}$$

must satisfy the following system of Volterra integral equations

$$\begin{aligned} v(t) &= 2 \int_0^b N(s(t), t; \xi, 0) h'(\xi) d\xi - 2 \int_0^t N(s(t), t; 0, \tau) \dot{f}(\tau) d\tau \\ &\quad + 2 \int_0^t G_x(s(t), t; s(\tau), \tau) v(\tau) d\tau \\ &\quad + 2 \int_0^t [N(s(t), t; s(\tau), \tau) - N(s(t), t; 0, \tau)] F(V(\tau)) d\tau. \end{aligned} \quad (2.4)$$

$$\begin{aligned} V(t) &= \int_0^b N(0, t; \xi, 0) h'(\xi) d\xi \\ &\quad - \int_0^t N(0, t; 0, \tau) \dot{f}(\tau) d\tau + \int_0^t G_x(0, t; s(\tau), \tau) v(\tau) d\tau \\ &\quad + \int_0^t [N(0, t; s(\tau), \tau) - N(0, t; 0, \tau)] F(V(\tau)) d\tau, \end{aligned} \quad (2.5)$$

where  $G$ ,  $N$  are the Green and Neumann functions and  $K$  is the fundamental solution of the heat equation, defined respectively by

$$\begin{aligned} G(x, t, \xi, \tau) &= K(x, t, \xi, \tau) - K(-x, t, \xi, \tau), \\ N(x, t, \xi, \tau) &= K(x, t, \xi, \tau) + K(-x, t, \xi, \tau), \\ K(x, t, \xi, \tau) &= \begin{cases} \frac{1}{2\sqrt{\pi(t-\tau)}} \exp\left(-\frac{(x-\xi)^2}{4(t-\tau)}\right) & t > \tau \\ 0 & t \leq \tau, \end{cases} \end{aligned}$$

where  $s(t)$  is given by (2.2),

*Proof.* Let  $u(x, t)$  be the solution to (1.3). We integrate, on the domain  $D_{t,\varepsilon} = \{(\xi, \tau) : 0 < \xi < s(\tau), \varepsilon < \tau < t - \varepsilon\}$ , the Green identity

$$(Gu_\xi - uG_\xi)_\xi - (Gu)_\tau = GF(u_\xi(0, \tau)). \quad (2.6)$$

Now we let  $\varepsilon \rightarrow 0$ , to obtain the following integral representation for  $u(x, t)$ ,

$$\begin{aligned} u(x, t) &= \int_0^b G(x, t; \xi, 0) h(\xi) d\xi + \int_0^t G_\xi(x, t; 0, \tau) f(\tau) d\tau \\ &\quad + \int_0^t G(x, t; s(\tau), \tau) u_\xi(s(\tau), \tau) d\tau - \iint_{D(t)} G(x, t; \xi, \tau) F(u_\xi(0, \tau)) d\xi d\tau. \end{aligned}$$

From the definition of  $v(t)$  and  $V(t)$  by (2.3), we obtain (2.1) and (2.2). If we differentiate  $u(x, t)$  in variable  $x$  and we let  $x \rightarrow 0^+$  and  $x \rightarrow s(t)$ , by using the jump relations, we obtain the integral equations for  $v$  and  $V$  given by (2.4) and (2.5).

Conversely, the function  $u(x, t)$  defined by (2.1) where  $v$  and  $V$  are the solutions of (2.4) and (2.5), satisfy the conditions (1.3) (i),(ii),(iv) and (v). In order to prove condition (1.3) (iii) we define  $\psi(t) = u(s(t), t)$ . Taking into account that  $u$  satisfy the conditions (1.3) (i),(ii),(iv) and (v), if we integrate the Green identity (2.6) over

the domain  $D_{t,\varepsilon}$ , ( $\varepsilon > 0$ ) and we let  $\varepsilon \rightarrow 0$  we obtain that

$$\begin{aligned} u(x, t) &= \int_0^b G(x, t; \xi, 0)h(\xi)d\xi + \int_0^t G(x, t; s(\tau), \tau)v(\tau) d\tau \\ &\quad + \int_0^t \psi(\tau)[G_x(x, t; s(\tau), \tau) - G(x, t; s(\tau), \tau)v(\tau)] d\tau \\ &\quad + \int_0^t G_\xi(x, t; 0, \tau)f(\tau) d\tau - \iint_{D(t)} G(x, t; \xi, \tau)F(V(\tau))d\xi d\tau. \end{aligned}$$

Then, if we compare this last expression with (2.1), we deduce that

$$M(x, t) = \int_0^t \psi(\tau)[G_x(x, t; s(\tau), \tau) - G(x, t; s(\tau), \tau)v(\tau)] d\tau \equiv 0 \quad (2.7)$$

for  $0 < x < s(t)$ ,  $0 < t < \sigma$ . We let  $x \rightarrow s(t)$  in (2.7) and by using the jump relations we have that  $\psi$  satisfy the integral equation

$$\frac{1}{2}\psi(t) + \int_0^t \psi(\tau)[G_x(s(t), t; s(\tau), \tau) - G(s(t), t; s(\tau), \tau)v(\tau)] d\tau = 0.$$

Then we deduce that

$$\begin{aligned} |\psi(t)| &\leq C \int_0^t \frac{|\psi(\tau)|}{\sqrt{t-\tau}} d\tau \\ &\leq C^2 \int_0^t \frac{d\tau}{\sqrt{t-\tau}} \int_0^\tau \frac{|\psi(\eta)|}{\sqrt{\tau-\eta}} d\eta \\ &= C^2 \int_0^t |\psi(\eta)| d\eta \int_\eta^t \frac{d\tau}{[(t-\tau)(\tau-\eta)]^{1/2}} \\ &= \pi C^2 \int_0^t |\psi(\eta)| d\eta \end{aligned}$$

where  $C = C(t)$ ; therefore by using the Gronwall inequality we have that  $\psi(t) = 0$  over  $[0, \sigma]$ .  $\square$

Next, we use the Banach fixed point theorem in order to prove the local existence and uniqueness of solution  $v, V \in C^0[0, \sigma]$  to the system of two Volterra integral equations (2.4)-(2.5) where  $\sigma$  is a positive small number. Consider the Banach space

$$C_{M,\sigma} = \left\{ \vec{w} = \begin{pmatrix} v \\ V \end{pmatrix} : v, V : [0, \sigma] \rightarrow \mathbb{R}, \text{ continuous, with } \|\vec{w}\|_\sigma \leq M \right\}$$

with

$$\|\vec{w}\|_\sigma := \|v\|_\sigma + \|V\|_\sigma := \max_{t \in [0, \sigma]} |v(t)| + \max_{t \in [0, \sigma]} |V(t)|$$

We define  $A : C_{M,\sigma} \rightarrow C_{M,\sigma}$ , such that

$$\vec{\tilde{w}}(t) = A(\vec{w}(t)) = \begin{pmatrix} A_1(v(t), V(t)) \\ A_2(v(t), V(t)) \end{pmatrix}$$

where

$$A_1(v(t), V(t)) = F_0(v(t)) + 2 \int_0^t [N(s(t), t, s(\tau), \tau) - N(s(t), t, 0, \tau)]F(V(\tau)) d\tau \quad (2.8)$$

with

$$F_0(v(t)) = 2 \int_0^b N(s(t), t, \xi, 0) h'(\xi) d\xi - 2 \int_0^t N(s(t), t, 0, \tau) \dot{f}(\tau) d\tau \\ + 2 \int_0^t G_x(s(t), t, s(\tau), \tau) v(\tau) d\tau$$

and

$$A_2(v(t), V(t)) = \int_0^b N(0, t, \xi, 0) h'(\xi) d\xi - \int_0^t N(0, t, 0, \tau) \dot{f}(\tau) d\tau \\ + \int_0^t G_x(0, t, s(\tau), \tau) v(\tau) d\tau \quad (2.9) \\ + \int_0^t [N(0, t, s(\tau), \tau) - N(0, t, 0, \tau)] F(V(\tau)) d\tau.$$

**Lemma 2.2.** *If  $v \in C^0[0, \sigma]$ ,  $\max_{t \in [0, \sigma]} |v(t)| \leq M$  and  $2M\sigma \leq b$  then  $s(t)$  defined by (2.2) satisfies*

$$|s(t) - s(\tau)| \leq M|t - \tau| \quad |s(t) - b| \leq \frac{b}{2}, \quad \forall t, \tau \in [0, \sigma].$$

To prove the following Lemmas we need the inequality

$$\exp\left(\frac{-x^2}{\alpha(t-\tau)}\right)/(t-\tau)^{n/2} \leq \left(\frac{n\alpha}{2ex^2}\right)^{n/2}, \quad \alpha, x > 0, t > \tau, n \in \mathbb{N}. \quad (2.10)$$

**Lemma 2.3.** *Let  $\sigma \leq 1$ ,  $M \geq 1$ ,  $f \in C^1[0, T]$ ,  $h \in C^1[0, b]$ ,  $F$  a Lipschitz function over  $C^0[0, T]$ . Under the hypothesis of Lemma 2.2, we have the following properties:*

$$\int_0^t |N(s(t), t, 0, \tau)| |\dot{f}(\tau)| d\tau \leq \|\dot{f}\|_t C_1(b)t \quad (2.11)$$

$$\int_0^t |G_x(s(t), t, s(\tau), \tau)| |v(\tau)| d\tau \leq M^2 C_2(b) \sqrt{t} \quad (2.12)$$

$$\int_0^b |N(s(t), t, \xi, 0)| |h'(\xi)| d\xi \leq \|h'\| \quad (2.13)$$

$$\int_0^t |N(s(t), t, s(\tau), \tau) - N(s(t), t, 0, \tau)| |F(V(\tau))| d\tau \leq C_4(L)M\sqrt{t} \quad (2.14)$$

$$\int_0^b |N(0, t, \xi, 0)| |h'(\xi)| d\xi \leq \|h'\| \quad (2.15)$$

$$\int_0^t |N(0, t, 0, \tau)| |\dot{f}(\tau)| d\tau \leq \frac{2\|\dot{f}\|_\sigma}{\sqrt{\pi}} \sqrt{t} \quad (2.16)$$

$$\int_0^t |G_x(0, t, s(\tau), \tau)| |v(\tau)| d\tau \leq C_3(b)Mt \quad (2.17)$$

$$\int_0^t |N(0, t, s(\tau), \tau) - N(0, t, 0, \tau)| |F(V(\tau))| d\tau \leq C_4(L)M\sqrt{t} \quad (2.18)$$

where  $L$  is the Lipschitz constant of  $F$  and

$$\begin{aligned} C_1(b) &= \left(\frac{8}{eb^2}\right)^{1/2} \frac{1}{\sqrt{\pi}}, & C_2(b) &= \frac{1}{2\sqrt{\pi}} + \frac{3b}{4\sqrt{\pi}} \left(\frac{2}{3eb^2}\right)^{3/2} \\ C_3(b) &= \frac{3b}{8\sqrt{\pi}} \left(\frac{24}{eb^2}\right)^{3/2}, & C_4(L) &= \frac{4L}{\sqrt{\pi}}. \end{aligned} \quad (2.19)$$

*Proof.* To prove (2.11), we have

$$\begin{aligned} |N(s(t), t, 0, \tau)| &= |K(s(t), t, 0, \tau) + K(-s(t), t, 0, \tau)| = 2K(s(t), t, 0, \tau) \\ &= \exp\left(\frac{-s^2(t)}{4(t-\tau)}\right) \frac{(t-\tau)^{-1/2}}{\sqrt{\pi}} \\ &\leq \exp\left(\frac{-b^2}{16(t-\tau)}\right) \frac{(t-\tau)^{-1/2}}{\sqrt{\pi}} \\ &\leq \left(\frac{8}{eb^2}\right)^{1/2} \frac{1}{\sqrt{\pi}} = C_1(b) \end{aligned}$$

then (2.11) holds. To prove (2.12), we have

$$\begin{aligned} |G_x(s(t), t, s(\tau), \tau)| &= |K_x(s(t), t, s(\tau), \tau) + K_x(-s(t), t, s(\tau), \tau)| \\ &= \frac{(t-\tau)^{-3/2}}{4\sqrt{\pi}} \left| (s(t) - s(\tau)) \exp\left(\frac{-(s(t) - s(\tau))^2}{4(t-\tau)}\right) \right. \\ &\quad \left. - (s(t) + s(\tau)) \exp\left(\frac{-(s(t) + s(\tau))^2}{4(t-\tau)}\right) \right| \\ &\leq \frac{(t-\tau)^{-3/2}}{4\sqrt{\pi}} \left( M(t-\tau) + 3b \exp\left(\frac{-9b^2}{4(t-\tau)}\right) \right) \\ &\leq \frac{1}{4\sqrt{\pi}} \left( M(t-\tau)^{-1/2} + 3b \left(\frac{2}{3eb^2}\right)^{3/2} \right). \end{aligned}$$

Then

$$\begin{aligned} \int_0^t |G_x(s(t), t, s(\tau), \tau)| |v(\tau)| d\tau &\leq \frac{M}{4\sqrt{\pi}} \left( 2M\sqrt{t} + 3b \left(\frac{2}{3eb^2}\right)^{3/2} t \right) \\ &\leq M^2 \sqrt{t} \left( \frac{1}{2\sqrt{\pi}} + \frac{3b}{M4\sqrt{\pi}} \left(\frac{2}{3eb^2}\right)^{3/2} \right) \\ &\leq M^2 C_2(b) \sqrt{t}, \end{aligned}$$

which implies (2.12). To prove (2.13), we have

$$\int_0^b |N(s(t), t, \xi, 0)| |h'(\xi)| d\xi \leq \|h'\| \int_0^\infty |N(s(t), t, \xi, 0)| d\xi \leq \|h'\|$$

because

$$\int_0^\infty |N(s(t), t, \xi, 0)| d\xi \leq 1.$$

To prove (2.14), by taking into account that

$$|N(s(t), t, s(\tau), \tau) - N(s(t), t, 0, \tau)| \leq \frac{2}{\sqrt{\pi(t-\tau)}}$$

we obtain

$$\begin{aligned} \int_0^t |N(s(t), t, s(\tau), \tau) - N(s(t), t, 0, \tau)|F(V(\tau))| d\tau &\leq LM \int_0^t \frac{2}{\sqrt{\pi(t-\tau)}} d\tau \\ &= C_4(L)M\sqrt{t}. \end{aligned}$$

The inequality (2.15) is prove in the same way as (2.13). To prove (2.16), we have

$$\begin{aligned} \int_0^t |N(0, t, 0, \tau)|\dot{f}(\tau)| d\tau &\leq \|\dot{f}\|_\sigma \int_0^t |N(0, t, 0, \tau)| d\tau \\ &= \|\dot{f}\|_\sigma \int_0^t \frac{1}{\sqrt{\pi(t-\tau)}} d\tau \\ &= \frac{\|\dot{f}\|_\sigma}{\sqrt{\pi}} 2\sqrt{t}. \end{aligned}$$

To prove (2.17), we have

$$\begin{aligned} |G_x(0, t, s(\tau), \tau)| &= \frac{(t-\tau)^{-3/2}}{4\sqrt{\pi}} s(\tau) \exp\left(\frac{-(s(\tau))^2}{4(t-\tau)}\right) \\ &\leq \frac{3b}{8\sqrt{\pi}} (t-\tau)^{-3/2} \exp\left(\frac{-b^2}{16(t-\tau)}\right) \\ &\leq \frac{3b}{8\sqrt{\pi}} \left(\frac{24}{eb^2}\right)^{3/2}. \end{aligned}$$

To prove (2.18), as in (2.14), we prove that

$$|N(0, t, s(\tau), \tau) - N(0, t, 0, \tau)| \leq \frac{2}{\sqrt{\pi(t-\tau)}}$$

and therefore (2.18) holds.  $\square$

**Lemma 2.4.** *Let  $s_1, s_2$  be the functions corresponding to  $v_1, v_2$  in  $C^0[0, \sigma]$ , respectively, with  $\max_{t \in [0, \sigma]} |v_i(t)| \leq M$ ,  $i = 1, 2$ , Then we have*

$$\begin{aligned} |s_2(t) - s_1(t)| &\leq t\|v_2 - v_1\|_t, \\ |s_i(t) - s_i(\tau)| &\leq M|t - \tau|, \quad i = 1, 2, \\ \frac{b}{2} &\leq s_i(t) \leq \frac{3b}{2}, \quad \forall t \in [0, \sigma], \quad i = 1, 2. \end{aligned} \tag{2.20}$$

**Lemma 2.5.** *Let  $f \in C^1[0, T]$ ,  $h \in C^1[0, b]$ ,  $F$  a Lipschitz function in  $C^0[0, T]$ . We have*

$$|F_0(v_2(t)) - F_0(v_1(t))| \leq E(b, h, f)\sqrt{t}\|v_2 - v_1\|_t; \quad (2.21)$$

$$\begin{aligned} & \int_0^t |N(s_2(t), t, s_2(\tau), \tau) - N(s_2(t), t, 0, \tau)|F(V_2(\tau)) - F(V_1(\tau))| d\tau \\ & \leq C_4(L)\sqrt{t}\|V_2 - V_1\|_t; \end{aligned} \quad (2.22)$$

$$\begin{aligned} & \int_0^t |N(s_2(t), t, 0, \tau) - N(s_1(t), t, 0, \tau)|F(V_1(\tau))| d\tau \\ & \leq C_5(b, L, M)t\|v_2 - v_1\|_t; \end{aligned} \quad (2.23)$$

$$\begin{aligned} & \int_0^t |N(s_2(t), t, s_2(\tau), \tau) - N(s_1(t), t, s_1(\tau), \tau)|F(V_1(\tau))| d\tau \\ & \leq [C_6(L, M)\sqrt{t} + C_7(b, L, M)t]\|v_2 - v_1\|_t; \end{aligned} \quad (2.24)$$

$$\int_0^t |G_x(0, t, s_2(\tau), \tau)|v_2(\tau) - v_1(\tau)| d\tau \leq C_8(b)t\|v_2 - v_1\|_t; \quad (2.25)$$

$$\begin{aligned} & \int_0^t |G_x(0, t, s_2(\tau), \tau)v_2(\tau) - G_x(0, t, s_1(\tau), \tau)v_1(\tau)| d\tau \\ & \leq (C_8(b)t + C_9(b, M)t^2)\|v_2 - v_1\|_t; \end{aligned} \quad (2.26)$$

$$\begin{aligned} & \int_0^t \left| [N(0, t, s_2(\tau), \tau) - N(0, t, 0, \tau)]F(V_2(\tau)) \right. \\ & \quad \left. - [N(0, t, s_1(\tau), \tau) - N(0, t, 0, \tau)]F(V_1(\tau)) \right| d\tau \\ & \leq C_4(L)\sqrt{t}\|V_2 - V_1\|_t + C_5(b, L, M)t^2\|v_2 - v_1\|_t, \end{aligned} \quad (2.27)$$

where the constants are defined by

$$\begin{aligned} C_4(L) &= \frac{4L}{\sqrt{\pi}}, \quad C_5(b, L, M) = LM \frac{3b}{8\sqrt{\pi}} \left(\frac{24}{eb^2}\right)^{3/2}, \\ C_6(L, M) &= \frac{LM^3}{\sqrt{\pi}}, \quad C_7(b, L, M) = \left(\frac{6}{eb^2}\right)^{3/2} \frac{3bLM^2}{2\sqrt{\pi}}, \\ C_8(b) &= \frac{3}{4\sqrt{\pi}} \left(\frac{24}{eb^2}\right)^{3/2}, \quad C_9(b, M) = \left[\left(\frac{40}{eb^2}\right)^{5/2} \frac{9b^2}{16\sqrt{\pi}} + \frac{1}{2\sqrt{\pi}} \left(\frac{24}{eb^2}\right)^{3/2}\right] \frac{M}{2}. \end{aligned} \quad (2.28)$$

*Proof.* The proof of (2.21) can be found in [13]. To prove (2.22), we have

$$|N(s_2(t), t, s_2(\tau), \tau) - N(s_2(t), t, 0, \tau)| \leq \frac{2}{\sqrt{\pi(t-\tau)}}.$$

Then

$$\begin{aligned} & \int_0^t |N(s_2(t), t, s_2(\tau), \tau) - N(s_2(t), t, 0, \tau)|F(V_2(\tau)) - F(V_1(\tau))| d\tau \\ & \leq \frac{4L}{\sqrt{\pi}}\sqrt{t}\|V_2 - V_1\|_t \end{aligned}$$



To prove (2.23), we use the mean value theorem: There exists  $c = c(t, \tau)$  between  $s_1(t)$  and  $s_2(t)$  such that

$$\begin{aligned} & |N(s_2(t), t, 0, \tau) - N(s_1(t), t, 0, \tau)| |F(V_1(\tau))| \\ &= |N_x(c, t, 0, \tau)| |s_2(\tau) - s_1(\tau)| |F(V_1(\tau))| \\ &\leq |c| \exp\left(-\frac{c^2}{4(t-\tau)}\right) \frac{(t-\tau)^{-3/2}}{2\sqrt{\pi}} LM\tau |v_2(\tau) - v_1(\tau)| \\ &\leq \frac{3b}{4\sqrt{\pi}} \exp\left(-\frac{b^2}{16(t-\tau)}\right) (t-\tau)^{-3/2} LM\tau |v_2(\tau) - v_1(\tau)| \\ &\leq \frac{3b}{4\sqrt{\pi}} \left(\frac{24}{eb^2}\right)^{3/2} LM\tau |v_2(\tau) - v_1(\tau)|. \end{aligned}$$

Then

$$\begin{aligned} & \int_0^t |N(s_2(t), t, 0, \tau) - N(s_1(t), t, 0, \tau)| |F(V_1(\tau))| d\tau \\ &\leq \frac{3b}{8\sqrt{\pi}} \left(\frac{24}{eb^2}\right)^{3/2} LMt \|v_2 - v_1\|_t = C_5(b, L, M)t \|v_2 - v_1\|_t. \end{aligned}$$

To prove (2.24), we have

$$\begin{aligned} & N(s_2(t), t, s_2(\tau), \tau) - N(s_1(t), t, s_1(\tau), \tau) \\ &= K(s_2(t), t, s_2(\tau), \tau) - K(s_1(t), t, s_1(\tau), \tau) \\ &\quad + K(-s_2(t), t, s_2(\tau), \tau) - K(-s_1(t), t, s_1(\tau), \tau). \end{aligned}$$

As in [24], for each  $(t, \tau)$ ,  $0 < \tau < t$ , we define

$$f_{t,\tau}(x) = \exp\left(\frac{-x^2}{4(t-\tau)}\right).$$

Then we have

$$\begin{aligned} & K(s_2(t), t, s_2(\tau), \tau) - K(s_1(t), t, s_1(\tau), \tau) \\ &= \frac{(t-\tau)^{-1/2}}{2\sqrt{\pi}} \left[ \exp\left(-\frac{(s_2(t) - s_2(\tau))^2}{4(t-\tau)}\right) - \exp\left(-\frac{(s_1(t) - s_1(\tau))^2}{4(t-\tau)}\right) \right] \\ &= \frac{(t-\tau)^{-1/2}}{2\sqrt{\pi}} [f_{t,\tau}(s_2(t) - s_2(\tau)) - f_{t,\tau}(s_1(t) - s_1(\tau))] \end{aligned}$$

and

$$\begin{aligned} & K(-s_2(t), t, s_2(\tau), \tau) - K(-s_1(t), t, s_1(\tau), \tau) \\ &= \frac{(t-\tau)^{-1/2}}{2\sqrt{\pi}} \left[ \exp\left(-\frac{(s_2(t) + s_2(\tau))^2}{4(t-\tau)}\right) - \exp\left(-\frac{(s_1(t) + s_1(\tau))^2}{4(t-\tau)}\right) \right] \\ &= \frac{(t-\tau)^{-1/2}}{2\sqrt{\pi}} [f_{t,\tau}(s_2(t) + s_2(\tau)) - f_{t,\tau}(s_1(t) + s_1(\tau))] \end{aligned}$$

By the mean value theorem there exists  $c = c(t, \tau)$  between  $s_2(t) - s_2(\tau)$  and  $s_1(t) - s_1(\tau)$  such that

$$\begin{aligned} & f_{t,\tau}(s_2(t) - s_2(\tau)) - f_{t,\tau}(s_1(t) - s_1(\tau)) \\ &= f'_{t,\tau}(c)(s_2(t) - s_2(\tau) - s_1(t) + s_1(\tau)) \\ &= \frac{-c}{2(t-\tau)} \exp\left(-\frac{c^2}{4(t-\tau)}\right)(s_2(t) - s_2(\tau) - s_1(t) + s_1(\tau)) \end{aligned}$$

Taking into account that

$$|c| \leq \max\{|s_i(t) - s_i(\tau)|, i = 1, 2\} \leq M(t - \tau)$$

it results

$$\begin{aligned} |f_{t,\tau}(s_2(t) - s_2(\tau)) - f_{t,\tau}(s_1(t) - s_1(\tau))| &\leq \frac{M}{2} [|s_2(t) - s_1(t)| + |s_2(\tau) - s_1(\tau)|] \\ &\leq M^2 \|v_2 - v_1\|_t. \end{aligned}$$

Then we have

$$|K(s_2(t), t, s_2(\tau), \tau) - K(s_1(t), t, s_1(\tau), \tau)| \leq \frac{M^2}{2\sqrt{\pi}(t-\tau)} \|v_2 - v_1\|_t.$$

In the same way we have

$$\begin{aligned} & f_{t,\tau}(s_2(t) + s_2(\tau)) - f_{t,\tau}(s_1(t) + s_1(\tau)) \\ &= f'_{t,\tau}(c^*)(s_2(t) + s_2(\tau) - s_1(t) - s_1(\tau)) \\ &= \frac{-c^*}{2(t-\tau)} \exp\left(-\frac{c^{*2}}{4(t-\tau)}\right)(s_2(t) + s_2(\tau) - s_1(t) - s_1(\tau)) \end{aligned}$$

where  $c^* = c^*(t, \tau)$  is between  $s_2(t) + s_2(\tau)$  and  $s_1(t) + s_1(\tau)$ . Since  $s_1(t) + s_1(\tau) \leq c^* \leq s_2(t) + s_2(\tau)$ , (or viceversa), we deduce that  $b \leq c^* \leq 3b$ , that is  $\exp(-c^{*2}/4(t-\tau)) \leq \exp(-b^2/4(t-\tau))$ . Then we obtain

$$\begin{aligned} & |K(-s_2(t), t, s_2(\tau), \tau) - K(-s_1(t), t, s_1(\tau), \tau)| \\ &= \frac{(t-\tau)^{-1/2}}{2\sqrt{\pi}} |f_{t,\tau}(s_2(t) + s_2(\tau)) - f_{t,\tau}(s_1(t) + s_1(\tau))| \\ &\leq \frac{3b}{4\sqrt{\pi}(t-\tau)^{3/2}} \exp\left(-\frac{b^2}{4(t-\tau)}\right) 2M \|v_2 - v_1\|_t \\ &\leq \left(\frac{6}{eb^2}\right)^{3/2} \frac{3bM}{2\sqrt{\pi}} \|v_2 - v_1\|_t \end{aligned}$$

and

$$\begin{aligned} & |N(s_2(t), t, s_2(\tau), \tau) - N(s_1(t), t, s_1(\tau), \tau)| \\ &\leq \left(\frac{M^2}{2\sqrt{\pi}(t-\tau)} + \left(\frac{6}{eb^2}\right)^{3/2} \frac{3bM}{2\sqrt{\pi}}\right) \|v_2 - v_1\|_t. \end{aligned}$$

Therefore,

$$\begin{aligned} & \int_0^t |N(s_2(t), t, s_2(\tau), \tau) - N(s_1(t), t, s_1(\tau), \tau)|F(V_1(\tau))| d\tau \\ & \leq \int_0^t \left( \frac{M^2}{2\sqrt{\pi}(t-\tau)} + \left(\frac{6}{eb^2}\right)^{3/2} \frac{3bM}{2\sqrt{\pi}} \right) \|v_2 - v_1\|_t |F(V_1(\tau))| d\tau \\ & \leq LM \left( \frac{M^2\sqrt{t}}{\sqrt{\pi}} + \left(\frac{6}{eb^2}\right)^{3/2} \frac{3bM}{2\sqrt{\pi}} t \right) \|v_2 - v_1\|_t \\ & = (C_6(L, M)\sqrt{t} + C_7(L, M, b)t) \|v_2 - v_1\|_t. \end{aligned}$$

To prove (2.25), we take into account (2.10):

$$\begin{aligned} G_x(0, t, s_2(\tau), \tau) &= K(0, t, s_2(\tau), \tau) \frac{s_2(\tau)}{t-\tau} \\ &= \exp\left(-\frac{s_2^2(\tau)}{4(t-\tau)}\right) \frac{(t-\tau)^{-3/2}}{2\sqrt{\pi}} s_2(\tau) \\ &\leq \frac{1}{2\sqrt{\pi}} \left(\frac{24}{eb^2}\right)^{3/2} s_2(\tau) \leq \frac{3b}{4\sqrt{\pi}} \left(\frac{24}{eb^2}\right)^{\frac{3}{2}} = C_8(b). \end{aligned}$$

To prove (2.26), we have

$$\begin{aligned} & |G_x(0, t, s_2(\tau), \tau)v_2(\tau) - G_x(0, t, s_1(\tau), \tau)v_1(\tau)| \\ & \leq |G_x(0, t, s_2(\tau), \tau)| \|v_2(\tau) - v_1(\tau)\| \\ & \quad + |G_x(0, t, s_2(\tau), \tau) - G_x(0, t, s_1(\tau), \tau)| \|v_1(\tau)\|. \end{aligned}$$

Using the mean value theorem there exists  $c = c(\tau)$  between  $s_2(\tau)$  and  $s_1(\tau)$  such that  $G_x(0, t, s_2(\tau), \tau) - G_x(0, t, s_1(\tau), \tau) = G_{x\xi}(0, t, c, \tau)(s_2(\tau) - s_1(\tau))$ . Taking into account the following properties

$$\begin{aligned} G_{x\xi}(0, t, c, \tau) &= \frac{K(0, t, c, \tau)}{t-\tau} \left( \frac{c^2}{2(t-\tau)} + 1 \right), \\ \frac{K(0, t, c, \tau)}{t-\tau} &= \frac{1}{2\sqrt{\pi}} \exp\left(-\frac{c^2}{4(t-\tau)}\right) (t-\tau)^{-3/2} \leq \frac{1}{2\sqrt{\pi}} \left(\frac{24}{eb^2}\right)^{3/2}, \\ K(0, t, c, \tau) \frac{c^2}{2(t-\tau)^2} &= \frac{1}{4\sqrt{\pi}} \exp\left(-\frac{c^2}{4(t-\tau)}\right) (t-\tau)^{-\frac{5}{2}} c^2 \leq \frac{9b^2}{16\sqrt{\pi}} \left(\frac{40}{eb^2}\right)^{5/2} \end{aligned}$$

we have

$$\begin{aligned} & |G_x(0, t, s_2(\tau), \tau) - G_x(0, t, s_1(\tau), \tau)| \|v_1(\tau)\| \\ & \leq \left( \frac{1}{2\sqrt{\pi}} \left(\frac{24}{eb^2}\right)^{\frac{3}{2}} + \frac{9b^2}{16\sqrt{\pi}} \left(\frac{40}{eb^2}\right)^{\frac{5}{2}} \right) |s_2(\tau) - s_1(\tau)| \|v_1(\tau)\| \\ & \leq M \left( \frac{1}{2\sqrt{\pi}} \left(\frac{24}{eb^2}\right)^{\frac{3}{2}} + \frac{9b^2}{16\sqrt{\pi}} \left(\frac{40}{eb^2}\right)^{\frac{5}{2}} \right) \tau \|v_2(\tau) - v_1(\tau)\|. \end{aligned}$$

Then

$$\begin{aligned} & \int_0^t |G_x(0, t, s_2(\tau), \tau) - G_x(0, t, s_1(\tau), \tau)| \|v_1(\tau)\| d\tau \\ & \leq \left( \frac{1}{2\sqrt{\pi}} \left(\frac{24}{eb^2}\right)^{\frac{3}{2}} + \frac{9b^2}{16\sqrt{\pi}} \left(\frac{40}{eb^2}\right)^{\frac{5}{2}} \right) \frac{Mt^2}{2} \|v_2 - v_1\|_t. \end{aligned}$$

Then (2.26) holds by using (2.25). To prove (2.27), we have

$$\begin{aligned} & [N(0, t, s_2(\tau), \tau) - N(0, t, 0, \tau)]F(V_2(\tau)) \\ & - [N(0, t, s_1(\tau), \tau) - N(0, t, 0, \tau)]F(V_1(\tau)) \\ & = [N(0, t, s_2(\tau), \tau) - N(0, t, 0, \tau)][F(V_2(\tau)) - F(V_1(\tau))] \\ & \quad + [N(0, t, s_2(\tau), \tau) - N(0, t, s_1(\tau), \tau)]F(V_1(\tau)) \end{aligned} \quad (2.29)$$

Using  $|N(0, t, s_2(\tau), \tau) - N(0, t, 0, \tau)| \leq \frac{2}{\sqrt{\pi(t-\tau)}}$  we get

$$|N(0, t, s_2(\tau), \tau) - N(0, t, 0, \tau)||F(V_2(\tau)) - F(V_1(\tau))| \leq \frac{2}{\sqrt{\pi(t-\tau)}}L|V_2(\tau) - V_1(\tau)|,$$

and

$$\begin{aligned} & \int_0^t |N(0, t, s_2(\tau), \tau) - N(0, t, 0, \tau)||F(V_2(\tau)) - F(V_1(\tau))| d\tau \\ & \leq \frac{4\sqrt{t}}{\sqrt{\pi}}L\|V_2 - V_1\|_t = C_4(L)\sqrt{t}\|V_2 - V_1\|_t. \end{aligned} \quad (2.30)$$

Furthermore,

$$|N(0, t, s_2(\tau), \tau) - N(0, t, s_1(\tau), \tau)| = |N_\xi(0, t, c, \tau)||s_2(\tau) - s_1(\tau)|$$

where  $c = c(\tau)$  is between  $s_2(\tau)$  and  $s_1(\tau)$  and

$$\begin{aligned} |N_\xi(0, t, c, \tau)||s_2(\tau) - s_1(\tau)| & = |-G_x(0, t, c, \tau)||s_2(\tau) - s_1(\tau)| \\ & \leq \frac{|c|}{2\sqrt{\pi}}\left(\frac{24}{eb^2}\right)^{3/2}\tau|v_2(\tau) - v_1(\tau)| \\ & \leq \frac{3b}{4\sqrt{\pi}}\left(\frac{24}{eb^2}\right)^{3/2}\tau|v_2(\tau) - v_1(\tau)|. \end{aligned}$$

Then

$$\begin{aligned} & \int_0^t |N(0, t, s_2(\tau), \tau) - N(0, t, s_1(\tau), \tau)||F(V_1(\tau))| d\tau \\ & \leq LM\frac{3b}{4\sqrt{\pi}}\left(\frac{24}{eb^2}\right)^{3/2}\frac{t^2}{2}\|v_2 - v_1\|_t = C_5(L, M, b)t^2\|v_2 - v_1\|_t \end{aligned} \quad (2.31)$$

Therefore, by (2.29), (2.30), and (2.31), the inequality (2.27) holds.  $\square$

**Theorem 2.6.** *The map  $A : C_{M,\sigma} \rightarrow C_{M,\sigma}$  is well defined and is a contraction map if  $\sigma$  satisfies the following inequalities:*

$$\sigma \leq 1, 2M\sigma \leq b \quad (2.32)$$

$$(2\|\dot{f}\|_\sigma C_1(b) + MC_3(b))\sigma + (2M^2C_2(b) + \frac{2\|\dot{f}\|_\sigma}{\sqrt{\pi}} + 3MC_4(L))\sqrt{\sigma} \leq 1 \quad (2.33)$$

$$D(b, f, h, L, M)\sqrt{\sigma} < 1, \quad (2.34)$$

where

$$M = 1 + 3\|h'\| \quad (2.35)$$

and

$$\begin{aligned} D_1(b, f, h, L, M) &= E(b, f, h) + 2C_6(L, M) + 3C_4(L) \\ D_2(b, L, M) &= 2[C_5(b, L, M) + 2C_7(b, L, M) + C_8(b)] \\ D_3(b, L, M) &= C_9(b, M) + C_5(b, L, M) \\ D(b, f, h, L, M) &= D_1(b, f, h, L, M) + D_2(b, L, M) + D_3(b, L, M). \end{aligned}$$

Then there exists a unique solution on  $C_{M, \sigma}$  to the system of integral equations (2.4), (2.5).

*Proof.* Firstly we demonstrate that  $A$  maps  $C_{\sigma, M}$  into itself, that is

$$\|A(\vec{w})\|_{\sigma} = \max_{t \in [0, \sigma]} |A_1(v(t), V(t))| + \max_{t \in [0, \sigma]} |A_2(v(t), V(t))| \leq M \quad (2.36)$$

Using the Lemmas 2.3, 2.4 and the definitions (2.8)-(2.9), we have

$$\begin{aligned} |A_1(v(t), V(t))| &\leq 2\|\dot{f}\|_{\sigma} C_1(b)t + 2M^2 C_2(b)\sqrt{t} + 2\|h'\| + 2C_4(L)M\sqrt{t}, \\ |A_2(v(t), V(t))| &\leq \|h'\| + \left(\frac{2\|\dot{f}\|_{\sigma}}{\sqrt{\pi}} + C_4(L)M\right)\sqrt{t} + C_3(b)Mt. \end{aligned}$$

Then

$$\begin{aligned} \|A(\vec{w})\|_{\sigma} &= \max_{t \in [0, \sigma]} |A_1(v(t), V(t))| + \max_{t \in [0, \sigma]} |A_2(v(t), V(t))| \\ &\leq 3\|h'\| + (2\|\dot{f}\|_{\sigma} C_1(b) + C_3(b)M)\sigma \\ &\quad + \left(2M^2 C_2(b) + \frac{2\|\dot{f}\|_{\sigma}}{\sqrt{\pi}} + 3MC_4(L)\right)\sqrt{\sigma}. \end{aligned}$$

Selecting  $M$  by (2.35) and  $\sigma$  such that (2.32) and (2.33) hold, we obtain (2.36).

Now, we prove that

$$\|A(\vec{w}_2) - A(\vec{w}_1)\|_{\sigma} \leq D(b, h, f, L, M)\sqrt{\sigma}\|\vec{w}_2 - \vec{w}_1\|_{\sigma}$$

where  $\vec{w}_1 = \begin{pmatrix} v_1 \\ V_1 \end{pmatrix}$ ,  $\vec{w}_2 = \begin{pmatrix} v_2 \\ V_2 \end{pmatrix}$ . By selecting  $\sigma$  such that (2.34) holds,  $A$  becomes a contraction mapping on  $C_{\sigma, M}$  and therefore it has a unique fixed point. To prove this assertion we consider

$$A(\vec{w}_1)(t) - A(\vec{w}_2)(t) = \begin{pmatrix} A_1(v_2(t), V_2(t)) - A_1(v_1(t), V_1(t)) \\ A_2(v_2(t), V_2(t)) - A_2(v_1(t), V_1(t)) \end{pmatrix}$$

where

$$\begin{aligned} &A_1(v_2(t), V_2(t)) - A_1(v_1(t), V_1(t)) \\ &= F_0(v_2(t)) - F_0(v_1(t)) + 2 \int_0^t [N(s_2(t), t, s_2(\tau), \tau) - N(s_2(t), t, 0, \tau)] F(V_2(\tau)) d\tau \\ &\quad - 2 \int_0^t [N(s_1(t), t, s_1(\tau), \tau) - N(s_1(t), t, 0, \tau)] F(V_1(\tau)) d\tau \end{aligned}$$

and

$$\begin{aligned} & A_2(v_2(t), V_2(t)) - A_2(v_1(t), V_1(t)) \\ &= \int_0^t [G_x(0, t, v_2(\tau), \tau)v_2(\tau) - G_x(0, t, v_1(\tau), \tau)v_1(\tau)] d\tau \\ & \quad + \int_0^t \{ [N(0, t, s_2(\tau), \tau) - N(0, t, 0, \tau)]F(V_2(\tau)) \\ & \quad - [N(0, t, s_1(\tau), \tau) - N(0, t, 0, \tau)]F(V_1(\tau)) \} d\tau. \end{aligned}$$

Taking into account the Lemmas 2.4 and 2.5 it results

$$\begin{aligned} & |A_1(v_2(t), V_2(t)) - A_1(v_1(t), V_1(t))| \\ & \leq E(b, h, f)\sqrt{t}\|v_2 - v_1\|_t + 2C_4(L)\sqrt{t}\|V_2 - V_1\|_t \\ & \quad + 2C_5(b, L, M)t\|v_2 - v_1\|_t + 2[C_6(L, M)\sqrt{t} + C_7(b, L, M)t]\|v_2 - v_1\|_t, \end{aligned}$$

and

$$\begin{aligned} & |A_2(v_2(t), V_2(t)) - A_2(v_1(t), V_1(t))| \\ & \leq (C_8(b)t + C_9(b, M)t^2)\|v_2 - v_1\|_t \\ & \quad + C_4(L)\sqrt{t}\|V_2 - V_1\|_t + C_5(b, L, M)t^2\|v_2 - v_1\|_t. \end{aligned}$$

Therefore,

$$\begin{aligned} & \|A(\vec{w}_2) - A(\vec{w}_1)\|_\sigma \\ & \leq \max_{t \in [0, \sigma]} |A_1(v_2(t), V_2(t)) - A_1(v_1(t), V_1(t))| \\ & \quad + \max_{t \in [0, \sigma]} |A_2(v_2(t), V_2(t)) - A_2(v_1(t), V_1(t))| \\ & \leq \{D_1(b, f, h, L, M)\sqrt{\sigma} + D_2(b, L, M)\sigma + D_3(b, L, M)\sigma^2\}\|\vec{w}_2 - \vec{w}_1\|_\sigma \\ & \leq D(b, f, h, L, M)\sqrt{\sigma}\|\vec{w}_2 - \vec{w}_1\|_\sigma. \end{aligned}$$

By hypothesis (2.34) we have that  $A$  is a contraction.  $\square$

**Remark.** If  $F$  satisfies the conditions

$$(H2) \quad F(V) > 0, \text{ for all } V \neq 0 \text{ and } F(0) = 0,$$

then by the maximum principle [5],  $u$  is a sub-solution for the same problem with  $F \equiv 0$ , that is

$$u(x, t) \leq u_0(x, t), \quad s(t) \leq s_0(t)$$

where  $u_0(x, t)$  and  $s_0(t)$  solve the classical Stefan problem

$$\begin{aligned} & u_{0t} - u_{0xx} = 0, \quad 0 < x < s_0(t), \quad 0 < t < T, \\ & u_0(0, t) = f(t) \geq 0, \quad 0 < t < T, \\ & u_0(s_0(t), t) = 0, \quad u_{0x}(s_0(t), t) = -\dot{s}_0(t), \quad 0 < t < T, \\ & u_0(x, 0) = h(x) \quad 0 \leq x \leq b = s_0(0). \end{aligned}$$

**Acknowledgments.** The authors have been partially sponsored by Project PIP No. 5379 from CONICET - UA (Rosario, Argentina), by Project #53900/4 from Fundaci3n Antorchas (Argentina), and and by ANPCYT PICT # 03-11165 from Agencia (Argentina).

## REFERENCES

- [1] E. K. Afenya and C. P. Calderon, *A remark on a nonlinear integral equation*, Rev. Un. Mat. Argentina, 39 (1995), 223-227.
- [2] V. Alexiades and A. D. Solomon, *Mathematical modeling of melting and freezing processes*, Hemisphere - Taylor & Francis, Washington (1983).
- [3] I. Athanassopoulos, G. Makrakis, and J. F. Rodrigues (Eds.), *Free Boundary Problems: Theory and Applications*, CRC Press, Boca Raton (1999).
- [4] L. R. Berrone, D. A. Tarzia, and L. T. Villa, *Asymptotic behavior of a non-classical heat conduction problem for a semi-infinite material*, Math. Meth. Appl. Sci., 23 (2000), 1161-1177.
- [5] M. Bertsch, *A class of degenerate diffusion equations with a singular nonlinear term*, Nonlinear Anal. Th. Meth. Appl., 7(1983), 117-127.
- [6] J. R. Cannon, *The one-dimensional heat equation*, Addison-Wesley, Menlo Park (1984).
- [7] J. R. Cannon and H. M. Yin, *A class of non-linear non-classical parabolic equations*, J. Diff. Eq., 79 (1989), 266-288.
- [8] H. S. Carslaw and J. C. Jaeger, *Conduction of heat in solids*, Oxford University Press, London (1959).
- [9] J. M. Chadam and H. Rasmussen H (Eds.), *Free boundary problems involving solids*, Pitman Research Notes in Mathematics Series #281, Longman Essex (1993).
- [10] J. Crank, *Free and moving boundary problems*, Clarendon Press Oxford (1984).
- [11] J. I. Diaz, M. A. Herrero, A. Liñan and J. L. Vazquez (Eds.), *Free boundary problems: theory and applications*, Pitman Research Notes in Mathematics Series #323, Longman, Essex (1995).
- [12] A. Fasano and M. Primicerio (Eds.), *Nonlinear diffusion problems*, Lecture Notes in Math N. 1224, Springer Verlag, Berlin (1986).
- [13] A. Friedman, *Free boundary problems for parabolic equations I. Melting of solids*, J. Math. Mech., 8 (1959), 499-517.
- [14] K. Glashoff and J. Sprekels, *The regulation of temperature by thermostats and set-valued integral equations*, J. Integral Eq., 4 (1982), 95-112.
- [15] G. Gripenberg, S.O. Londen, and O. Staffans, *Volterra integral and functional equations*, Cambridge Univ. Press, Cambridge (1990).
- [16] R. A. Handelsman and W. E. Olmstead, *Asymptotic solution to a class of nonlinear Volterra integral equations*, SIAM J. Appl. Math., 22 (1972), 373-384.
- [17] N. Kenmochi and M. Primicerio, *One-dimensional heat conduction with a class of automatic heat source controls*, IMA J. Appl. Math. 40 (1988), 205-216.
- [18] V. J. Lunardini, *Heat transfer with freezing and thawing*, Elsevier, Amsterdam (1991).
- [19] W. R. Mann and F. Wolf, *Heat transfer between solid and gases under nonlinear boundary conditions*, Quart. Appl. Math., 9 (1951), 163-184.
- [20] R. K. Miller, *Nonlinear Volterra integral equations*, W. A. Benjamin, Menlo Park (1971).
- [21] W. E. Olmstead and R. A. Handelsman, *Diffusion in a semi-infinite region with nonlinear surface dissipation*, SIAM Rev., 18 (1976), 275-291.
- [22] J. H. Roberts and W. R. Mann, *A certain nonlinear integral equation of the Volterra type*, Pacific J. Math., 1 (1951), 431-445.
- [23] L. I. Rubinstein, *The Stefan problem*, Trans. Math. Monographs # 27, Amer. Math. Soc., Providence (1971).
- [24] B. Sherman, *A free boundary problem for the heat equation with prescribed flux at both fixed face and melting interface*, Quart. Appl. Math., 25 (1967), 53-63,
- [25] D. A. Tarzia, *A bibliography on moving-free boundary problems for the heat-diffusion equation. The Stefan and related problems*, MAT-Serie A, Rosario, # 2 (2000), (with 5869 titles on the subject, 300 pages). See [www.austral.edu.ar/MAT-SerieA/2\(2000\)/](http://www.austral.edu.ar/MAT-SerieA/2(2000)/)
- [26] D. A. Tarzia, *A Stefan problem for a non-classical heat equation*, MAT-Serie A, Rosario, # 3 (2001), 21-26.
- [27] D. A. Tarzia and L. T. Villa, *Some nonlinear heat conduction problems for a semi-infinite strip with a non-uniform heat source*, Rev. Un. Mat. Argentina, 41 (2000), 99-114.
- [28] L. T. Villa, *Problemas de control para una ecuación unidimensional del calor*, Rev. Un. Mat. Argentina, 32 (1986), 163-169.

- [29] L. C. Wrobel and C. A. Brebbia (Eds.), *Computational methods for free and moving boundary problems in heat and fluid flow*, Computational Mechanics Publications, Southampton (1993).

ADRIANA C. BRIOZZO

DEPARTAMENTO DE MATEMÁTICA, FCE, UNIVERSIDAD AUSTRAL, PARAGUAY 1950, S2000FZF  
ROSARIO, ARGENTINA

*E-mail address:* `Adriana.Briozzo@fce.austral.edu.ar`

DOMINGO ALBERTO TARZIA

DEPARTAMENTO DE MATEMÁTICA - CONISET, FCE, UNIVERSIDAD AUSTRAL, PARAGUAY 1950,  
S2000FZF ROSARIO, ARGENTINA

*E-mail address:* `Domingo.Tarzia@fce.austral.edu.ar`