

Overexpression of SPARC obliterates the *in vivo* tumorigenicity of human hepatocellular carcinoma cells

Catalina Atorrasagasti^{1†}, Mariana Malvicini^{1†}, Jorge B. Aquino^{1,2}, Laura Alaniz^{1,2}, Mariana Garcia^{1,2}, Marcela Bolontrade^{2,3}, Manglio Rizzo¹, Osvaldo L. Podhajcer^{2,3} and Guillermo Mazzolini^{1,2}

¹Gene Therapy Laboratory, Liver Unit School of Medicine, Austral University, Av. Presidente Perón 1500, (B1629ODT) Derqui-Pilar, Buenos Aires, Argentina

²CONICET (Consejo Nacional de Investigaciones Científicas y Técnicas)

³Laboratory of Molecular and Cellular Therapy, Fundación Instituto Leloir, Patricias Argentinas 435, (C1405BWE) Buenos Aires, Argentina

Hepatocellular carcinoma (HCC) is the sixth most common cancer and the third leading cause of cancer-related death worldwide. Current treatments are extremely disappointing. SPARC (Secreted protein, acidic and rich in cysteine) is a matricellular glycoprotein with differential expression in several tumors, including HCC, which significance remains unclear. We infected HCC cells (HepG2, Hep3B and Huh7) with an adenovirus expressing SPARC (AdsSPARC) to examine the role of SPARC expression on HCC cells and its effect on tumor aggressiveness. The *in vitro* HCC cells substrate-dependent proliferation and cell cycle profile were unaffected; however, SPARC overexpression reduced HCC proliferation when cells were grown in spheroids. A mild induction of cellular apoptosis was observed upon SPARC overexpression. SPARC overexpression resulted in spheroid growth inhibition *in vitro* while no effects were found when recombinant SPARC was exogenously applied. Moreover, the clonogenic and migratory capabilities were largely decreased in SPARC-overexpressing HCC cells, altogether suggesting a less aggressive HCC cell phenotype. Consistently, AdsSPARC-transduced cells showed increased E-cadherin expression and a concomitant decrease in N-cadherin expression. Furthermore, SPARC overexpression was found to reduce HCC cell viability in response to 5-FU-based chemotherapy *in vitro*, partially through induction of apoptosis. *In vivo* experiments revealed that SPARC overexpression in HCC cells inhibited their tumorigenic capacity and increased animal survival through a mechanism that partially involves host macrophages. Our data suggest that SPARC overexpression in HCC cells results in a reduced tumorigenicity partially through the induction of mesenchymal-to-epithelial transition (MET). These evidences point to SPARC as a novel target for HCC treatment.

HCC is the sixth most common cancer and the third leading cause of cancer-related death in the world.¹ Unfortunately, the incidence and mortality associated with HCC is increas-

Key words: Secreted protein, acidic and rich in cysteine, gene therapy, hepatocellular carcinoma, cadherins, mesenchymal-to-epithelial transition, 5-FU-based chemotherapy

Abbreviations: HCC: hepatocellular carcinoma; SPARC: secreted protein, acidic and rich in cysteine; TAM: tumor associated macrophages; MET: mesenchymal-to-epithelial transition; FBS: fetal bovine serum; TGF- β 1: transforming growth factor beta 1; rSPARC: recombinant human SPARC

[†]These author's equally contributed to this work.

Grant sponsor: Universidad Austral, Agencia Nacional de Promoción Científica y Tecnológica; **Grant numbers:** PICT-2006-1882, PICT-2005-34788, PICTO-CRUP-2005-31179, CTE-06, AECI 2007, AECI 2008

DOI: 10.1002/ijc.24966

History: Received 17 Jun 2009; Accepted 6 Oct 2009; Online 14 Oct 2009

Correspondence to: Guillermo Mazzolini, Liver Unit, School of Medicine, Austral University, Av. Presidente Perón 1500, (B1629ODT) Derqui-Pilar, Buenos Aires, Argentina, Tel: +54-2322-482618, Fax: +54-2322-482204, E-mail: gmazzoli@cas.austral.edu.ar

ing steadily in the United States as in Europe.² Current curative options can be applied to a paucity of patients and, in general, the prognosis of HCC is dismal due to underlying cirrhosis as well as to poor tumor response to chemotherapeutic regime.^{3,4} Therefore, novel therapies are urgently needed for advanced HCC.⁵

There is a complex cross-talk between cancer cells and different tumor microenvironmental components, such as fibroblasts, endothelial cells, tumor-associated macrophages (TAM) and matricellular proteins.⁶ Accumulating evidence indicates that this dynamic cross-talk can modulate tumor cell capacity to invade and disseminate.⁶ It is therefore highly relevant to assess whether and how those environmental factors are able to regulate tumor cellular processes in cancer disease. With this regard and especially in the case of HCC, in which cirrhosis is the underlying disease of most patients,⁴ the focus is placed on analyzing cellular events triggered by cell-matrix interactions.

SPARC, also named BM40 or osteonectin, is a secreted multifunctional matricellular glycoprotein involved in a wide number of biological processes during development, tissue repair and remodeling.⁷⁻¹⁰ Among them, it was shown to inhibit cell cycle in fibroblasts and endothelial cells, by arresting cells at G1.¹¹ In certain cell types, SPARC has also been shown to reduce cell adhesion to integrin-ligand coated

surfaces due to disruption of focal adhesion complexes, causing inhibition of cell spreading.¹² SPARC is known to interact with several extracellular matrix components and to bind to, and modulate the expression and activity of several growth factors and matrix metalloproteinases.^{13–15} For instance, SPARC has been shown to induce antiangiogenesis likely through blockade of the VEGF and FGF-2 induced endothelial cell proliferation, or to block tumor stromal cell growth through inhibition of PDGF activity.^{16–18}

Changes in SPARC expression levels were evidenced in several malignant tumors of epithelial and nonepithelial origin.^{19–22} Nevertheless, the role of SPARC in cancer is controversial and it seems to depend on the type of tumor.²³ For example, overexpression of SPARC in neuroblastoma,²⁴ pancreatic adenocarcinoma cells,²⁵ ovarian cancer,²⁶ colorectal carcinoma²⁷ and acute myeloid leukemia²⁸ was associated with good prognosis. Induction of cancer cell apoptosis and enhanced sensitivity to chemotherapeutic drugs are mechanisms likely involved in the beneficial antitumoral effects observed upon SPARC enhanced expression in colon cancer cells.²⁹

Despite beneficial effects of SPARC overexpression in certain tumor types, the induction of SPARC has been positively associated with increased aggressiveness in melanoma,^{30,31} glioblastoma,³² prostate³³ and breast cancer³⁴. It seems that SPARC effects on different cancer types might depend on whether SPARC acts directly on tumor cells or indirectly, by influencing adjacent stromal cells.²³

The expression of SPARC was shown to be induced in myofibroblasts of cirrhotic livers in HCV chronically infected patients³⁵ and in experimental liver cirrhosis.³⁶ Similarly, SPARC overexpression was observed in HCC, along capillaries present in the tumor capsule.³⁷ Lau *et al.* showed HCC antitumoral effects of SPARC forced expression in human tumor xenografts which correlated with a decreased neo-angiogenesis,²² but no mechanism was proposed. Thus, even though some authors showed SPARC overexpression in HCC^{22,35,37,38} and others reported some antitumor effects²² the significance of SPARC expression changes in HCC still remains unclear and the mechanisms involved, unknown.

Cadherins are a superfamily of transmembrane glycoproteins that mediate intercellular adhesion as well as other signaling events.³⁹ Disruptions in E-cadherin expression have been associated with enhanced aggressiveness of certain types of tumors including HCC.^{40–42} Moreover, enhanced expression of mesenchymal-associated cadherins, such as N-cadherin, in cancer cells increased tumor cell aggressiveness and facilitated tumor dissemination.⁴³

In an effort to assess the role and action mechanisms of SPARC in HCC, we transiently overexpressed SPARC in HepG2 HCC cells by transduction with an adenoviral vector (AdsSPARC). Our results hereby show for the first time that a transient increase in SPARC expression levels on HCC cells reduced their spheroid growth, clonogenic, migratory and adherent capabilities. The *in vivo* growth capacity of SPARC-overexpressing HCC cells was strongly inhibited in nude

mice, a feature now associated with an increased number of host macrophage cells. Interestingly, up-regulation of SPARC expression decreases HCC cell viability in response to 5-FU-based chemotherapy. More importantly, overexpression of SPARC was associated with an up-regulation of E-cadherin and a consistent down-regulation of N-cadherin, a tandem that likely results in a less aggressive HCC cell phenotype.

Material and Methods

Generation of recombinant adenoviral vectors

AdsSPARC, a first-generation replication-defective adenovirus was constructed and produced as previously described.³⁶ Briefly, a 1.7 kb *Sa*I fragment containing the coding sequence of human SPARC or a 527 bp *Sa*II fragment containing the bacterial β -galactosidase gene were cloned into a pADPSY-LTRSVpolyA vector to generate adenoviral vectors carrying SPARC cDNA in sense orientation (AdsSPARC) or Ad- β gal, respectively. AdsSPARC and Ad- β gal were expanded in HEK-293 cells, purified by cesium chloride density gradient, desalted using a PD-10 Sephadex G-25 column (Amersham Biosciences, Piscataway, NJ), and stored at -80°C . The concentration of recombinant vector was expressed as 50% tissue culture infectious doses (TCID₅₀) per milliliter.⁴⁴ For SPARC downregulation experiments, adenoviral vectors carrying SPARC cDNA in antisense orientation (AdasSPARC) were generated and experimentally applied.³⁶

Cells and cell culture

HepG2, Hep3B and HuH7 human hepatocellular carcinoma cells (kindly provided by Prof. Prieto, University of Navarra) were grown in Dulbecco's Modified Eagles Medium (DMEM) supplemented with 10% v/v fetal bovine serum (FBS) and antibiotics. Cell cultures were maintained at 37°C in a 5% CO₂ humidified incubator. HCC cells, transduced with AdsSPARC, AdasSPARC or Ad β gal at MOI of 100 or nontransduced cells were used in different experiments.

Immunofluorescence assays

For immunofluorescence studies of SPARC expression HepG2 cells were fixed in 4% paraformaldehyde and blocked with 10% goat serum in phosphate buffered saline (PBS)-0.2%Tween for 60 min at room temperature, followed by overnight incubation at 4°C in a humid chamber with a polyclonal mouse anti-SPARC antibody (32 $\mu\text{g}/\text{ml}$; Hybridoma). After 3 washes with PBS, bound antibodies were detected with FITC-conjugated goat anti-mouse IgG (Jackson Immuno Research, West Grove, PA). Secondary antibody was diluted 1:40 in PBS-0.2% Tween and incubated for 2 hr at 37°C . Nuclear morphology was examined by staining with DAPI. Images were captured from a Nikon E800 microscope coupled to a CCD camera. Control experiments without primary antibody showed only a faint background staining (not shown).

Three-dimensional spheroids

Ninety-six-well tissue culture plates were coated with 75 μ l of 1% agarose in PBS. Nearly confluent non-transduced HCC, HCC/AdsSPARC, HCC/AdasSPARC or HCC/Ad- β gal (HepG2 and Hep3B) were washed twice with PBS, trypsinized, and seeded at 5×10^3 cells/well in 150 μ l of 2% FBS DMEM to obtain a single homotypic spheroid per well. Seventy-five microliters of supernatant were carefully removed from each well every 3 days and replaced with fresh medium. Spheroid size was measured at days 2 and 6 using an inverted microscope and photographed. Length and width were measured using the ImageJ program (NIH). Spheroid volume was expressed as arbitrary units. Noninfected HepG2 cells were seeded at 5×10^3 cells/well in 150 μ l of 2% FBS DMEM with or without 0.5 μ g/ml of recombinant SPARC. Spheroids volume was measured as described above.

Proliferation assays and *in vitro* apoptosis assessment

Cell proliferation was measured using the colorimetric MTT assay (Invitrogen). Briefly, 3×10^3 HCC cells (HepG2, Hep3B and Huh7)/well were seeded onto 96-well plates in a final volume of 100 μ l per well. At each time point, culture medium was replaced with 100 μ l of 5 mg/ml 3-(4,5-dimethylthiazol-3-yl)-2,5-diphenyl tetrazolium bromide. Four hours later, the formazan dye was solubilized and read at 490 nm optical density as described (Denizot and Lang, 1986). Each assay was performed 3 times in triplicate. HepG2 spheroids were trypsinized and proliferation was similarly analyzed. For *in vitro* chemotherapy assays, HepG2 cells were transduced with AdsSPARC, Ad- β gal or left nontransduced for 2 days, washed and further incubated for 24 hr with no 5-FU or with different doses of 5-FU ranging from 0.1 to 10 μ g/ml. Cell viability was assessed by MTT assay as described above.

Morphological changes associated with apoptosis were assessed by acridine orange-ethidium bromide mixture staining (Sigma). Single cell suspensions were stained with 10 μ g/ml of the mixture and cells were visualized under a fluorescence microscope. Apoptotic cells were defined as those stained in yellow and showing cytoplasmic and nuclear shrinkage and chromatin condensation or fragmentation. At least 100 cells were counted from 4 independent experiments and the percentage of apoptotic and necrotic cells was determined. Apoptosis analysis was performed on HepG2 cells infected with AdsSPARC, Ad- β gal or left uninfected by flow cytometry at 48 hr using an apoptosis detection kit based on annexin-V staining (eBioscienceTM, San Diego, CA). Briefly, HepG2 cells (1×10^6) were collected with EDTA, washed twice in PBS, centrifuged and incubated with 5 μ l of annexin-V in binding buffer at room temperature for 15 min. Cells were vortexed and centrifuged, and the resultant pellets were washed and stained with 1 μ l propidium iodide in binding buffer. The percentage of apoptotic cells were immediately analyzed in a FACSCalibur (Becton Dickinson).

Results are representative of 4 different experiments. For quantifications, mean values from each independent experiment were normalized to their respective control values, prior to statistical comparisons.

Cell cycle analysis

For cell cycle analysis, 2×10^6 cells were collected, washed in PBS and fixed in a mixture of ice-cold 70% (v/v) ethanol, FBS and distill water. Fixed cells were centrifuged and stained with propidium iodide (PI) solution (50 mg/ml PI, 180 U/ml RNase). DNA content was determined using a FACScan flow cytometer (Becton Dickinson Immunocytometry Systems, San Jose, CA).

Colony formation and cell migration assay

For colony formation assay, 1×10^3 HCC cells (HepG2, Hep3B and Huh7) left untreated (control) or transduced with AdsSPARC, AdasSPARC or Ad- β gal were plated onto 60-mm dishes and incubated for 2 weeks before staining with crystal violet. Colonies, composed by 20–25 cells, were quantified under phase-contrast light microscopy. Three independent experiments were performed, in triplicates. For transwell cell migration assays, non-transduced or AdsSPARC or Ad- β gal transduced HepG2 or Huh7 cells (5×10^4 in 100 μ l of 0.1% FBS DMEM) were seeded on the upper chambers of 48-well chamber plates (Neuroprobe). In the lower chamber, 5 ng/ml TGF- β 1 was added as the chemoattractant. After 16 hr incubation at 37°C in 5% CO₂, the cells that remained on the upper surface of the membrane were removed by wiping with a cotton bud. Migrated cells attached to the lower surface of the membrane were fixed with 2% formaldehyde/PBS and stained with 10% May Gr \ddot{u} ndwald-Giemsa. The number of migrated cells on each membrane was counted under a microscope ($\times 100$), for ten random microscopic fields per membrane, and averaged. To analyze the differential adhesive capacity of treated cells onto polycarbonate membranes, HepG2 cells were seeded on the upper chambers and 4 hr later, the cells attached to the upper surface of the membrane were fixed and stained.

Western blot analysis

Human SPARC was detected in supernatants by monoclonal anti-SPARC antibody (diluted 1:500). Supernatants were collected and centrifuged twice at 4°C for 20 min at 10,000g. After centrifugation, cleared supernatants were stored at –80°C until analysis. E-cadherin was detected in HepG2 extracts by mouse monoclonal anti-E-cadherin antibody (diluted 1:2,000; kindly provided by Dr. Berasain, University of Navarra). Briefly, cells were collected and incubated in lysis buffer with protease inhibitors (50 mM Tris-HCl buffer, pH7.4, containing 0.1% Tween-20, 150mM NaCl, 10 μ g/ml aprotinin, 5 μ g/ml leupeptin, 1 mM PMSF) 30 min on ice. Measurement of total protein concentration was performed using Bradford assay.⁴⁵ For immunoblotting, 100 μ g of total protein was loaded and separated on 10% SDS-PAGE gels

and transferred to polyvinylidene difluoride membranes as described previously.³¹ Blots were then developed with HRP-conjugated goat anti-mouse IgG (Jackson ImmunoResearch Labs, West Grove, PA) diluted 1:5,000 in blocking buffer.³¹ Bands were detected using the ECL detection system. Protein loading and transfer for E-cadherin was monitored using an anti-actin antibody (diluted 1/700, Santa Cruz Biotechnology, Santa Cruz, CA) and incubated with horseradish peroxidase-conjugated goat anti-mouse antibody (diluted 1/5,000, Santa Cruz Biotechnology, Santa Cruz, CA). For SPARC protein loading membranes were stained with Ponceau Red. Bands intensities were measured by densitometer analysis using the Scion Image software (Scion Corporation, USA).

N-cadherin flow cytometry

For flow cytometry analysis, HepG2 cells grown for 48 and 72 hr in medium supplemented with 2% FBS were washed and detached with 1.25 mM EDTA in PBS. After washing with ice-cold medium, cells were incubated with 2 mg/mL of a monoclonal anti-N-cadherin antibody (anti-A-CAM, clone CG-4, Sigma, St. Louis, MO) for 30 min, washed twice with PBS 0.1% BSA. After washings, samples were incubated for 30 min with a secondary goat FITC-conjugated anti-mouse IgG antibody (Jackson Immuno Research, West Grove, PA) and fixed in 4% w/v paraformaldehyde in PBS for 10 min at room temperature. Cells were washed, resuspended in PBS and subjected to flow cytometry using a FACSCalibur (Becton Dickinson) flow cytometer.

Animal studies

For assessment of the *in vivo* tumor growth, HepG2 or HuH7 HCCs were left untransduced or were *ex vivo*-transduced with AdsSPARC, AdasSPARC or Ad- β gal, at a MOI of 100. Twenty hours later, cells were trypsinized, counted, and resuspended in 100 μ l of saline. Six to eight-week-old male athymic N:NIH(S)-nu mice were subcutaneously injected into the right flank with 1.5×10^6 HepG2 cells or 5×10^6 Huh7 cells. Perpendicular diameters were used to determine tumor volume ($V = (dl \times ds^2 \times 0.52)$, where ds is the smaller diameter and dl is the larger one). Ten animals were allocated to each treatment group. For histological analysis, tumor tissue was obtained at day 5 post HCC cells inoculation. All procedures were performed according to the "Guide for the Care and Use of Laboratory Animals" published by the U.S. National Research Council (National Academy Press, Washington, D.C. 1996) and approved by the School of Biomedical Sciences of the Austral University.

Histology and immunocyto/histochemistry

Tumor sections from individual mice were fixed in 10% formalin, embedded in paraffin, sectioned, and stained with hematoxylin-eosin for morphological evaluation. Five-micrometer-thick tumor sections were de-paraffinized in xylene and rehydrated in graded ethanol. Endogenous peroxidase activity was blocked by incubating with 3% hydrogen peroxide in

90% ethanol for 20 min. Nonspecific primary antibody binding was blocked by incubating sections in normal goat serum (10% in PBS). Endogenous biotin and avidin was blocked with blocking agent complex (Vectastain ABC Elite; Vector Labs., Burlingame, CA). For detection of macrophages, tissue sections were incubated with a rat anti-F4/80 monoclonal antibody (1:100; Serotec) overnight at 4°C in PBS containing 0.1% triton 0.1% bovine serum albumin. Slides were incubated with peroxidase-linked biotinylated goat anti-mouse secondary antibodies for 60 min, washed and further incubated with the ABC kit (Vector Laboratories). Sections were washed and incubated in a mixture of 3.3-diaminobenzidine (DAB) and nickel-ammonium salt for enhancement of the signal.⁴⁶ Controls for immunostaining specificity in which the primary antibody was replaced by nonimmune mouse serum or omitted, were negative. Quantitative analysis of immunohistochemical staining for F4/80 was performed using the ImageJ software, from 400 \times magnification images.

For N-cadherin and smooth muscle actin (SMA) immunocytochemistries, 1,000 HepG2 cells/well were plated onto poly-L-lysine/fibronectin coated coverslips placed on 24-well plates for 8 hr and then fixed in 4% paraformaldehyde. Cells were incubated overnight with a mouse anti-N-cadherin antibody (1:50; Zymed) or a mouse anti- α -SMA (1:200, Sigma), and specific antibody binding was evidenced by incubation with a donkey anti-mouse secondary antibody (1:100; Vector Labs.) followed by chromogenic reaction (ABC kit, Vector Laboratories) with nickel salt enhancement of the signal.⁴⁶ At least 2 independent experiments were performed, with similar results. For quantifications, only spread cells were considered. The N-cadherin and α -SMA immunostaining intensities were measured using the ImageJ software by analyzing 20 randomly sampled cells per condition. For this purpose, a mean intensity value per cell was obtained from 5 random values taken from similar areas of the cytoplasm. Cell morphology was determined by establishing similar cultures of HepG2 cells after 16 hr of incubation.

In vivo apoptosis TUNEL assay

For apoptosis assay, 5-micrometer-thick tumor sections were fixed with 10% formalin for 20 m and the terminal deoxynucleotidyl transferase-mediated DUPT nick-end labeling (TUNEL) assay was done following the manufacturer's protocol (Fluorescein-FragELTM DNA Fragmentation Detection Kit, Calbiochem, Darmstadt, Germany). TUNEL-positive cells were analyzed by using a standard fluorescein filter (465–495) and were viewed with a microscope (Nikon).

Statistical analysis

Data are expressed as mean \pm SEM. Statistical analysis was performed using Student's *t*, Mann-Whitney tests or Kruskal-Wallis, when appropriate. Differences at $p < 0.05$ were considered to be significant. Survival rates were calculated with the Kaplan-Meier method and their differences were evaluated by the log-rank test. Data analysis was performed with

the Prism GraphPad (GraphPad Software, Inc., San Diego, CA).

Results

Endogenous SPARC overexpression inhibits HCC cell spheroid growth

As shown by western blot (Fig. 1a) and by immunofluorescence (Fig. 1b), SPARC expression was identified in HepG2 cells. AdsSPARC transduction (MOI 100) resulted in a significant SPARC overexpression at 3 days after gene transfer (Fig. 1a).

To establish the role of SPARC overexpression in HCC cells we initially performed *in vitro* studies growing HCC cells in tri-dimensional spheroids. A profound inhibitory effect on spheroid growth of HepG2 cells was observed following SPARC overexpression (Fig. 2a, upper panel; 2b). On the contrary, transduction of HCC cells with Ad- β gal or AdasSPARC had no effects on spheroid growth capacity comparable to untransduced cells. Interestingly, when HepG2 cells were exogenously treated with recombinant SPARC (Fig. 2c) spheroid growth rate was not affected, indicating that only endogenous increased levels of this protein was able to induce the inhibitory effects on HCC growth. A similar inhibition in spheroid growth formation was found in SPARC-overexpressing Hep3B cells (Fig. 2a, lower panel).

Increased endogenous SPARC expression levels do not affect proliferation of HCC cells

To examine whether SPARC overexpression on HCC cells (HepG2, Hep3B and Huh7) could influence tumor cell substrate-dependent proliferation capacity, cells were assessed for their *in vitro* growth capacity in plastic. No significant differences were found in between groups up to 72 hr (Fig. 3a). Similar results were observed at 120 hr (not shown). We next aimed to address whether AdsSPARC transduction on HCC cells could induce apoptosis. For this purpose, HepG2 cells were stained with acridine orange-ethidium bromide mixture solution and they were subsequently visualized under immunofluorescence microscope. SPARC overexpression resulted in a mild and nonsignificant increased induction of cellular apoptosis (Fig. 3b). However, this effect was shown to be significant by flow cytometry when the annexin V staining was used instead (Fig. 3c). Thus, to confirm the lack of significant effects of SPARC overexpression on HCC cell proliferation and to address the extent of apoptosis events, the pattern of cell cycle progression was analyzed. No differences were found among treatments (Fig. 3d). Similar results were obtained when cell cycle analysis was performed on Hep3B cells (G1 phase = 55.5%, 61% and 66%; S phase = 35.6%, 31% and 24.5%; G2 phase = 8.9%, 7.2% and 9.1%; for untransduced, Ad- β gal and AdsSPARC transduced cells, respectively). Since both the proliferation rate and the cell cycle pattern remained unaffected, we conclude that SPARC overexpression-mediated increase in HCC cellular apoptosis




Figure 1. Overexpression of SPARC by AdsSPARC. Western blot analysis of SPARC on HepG2 hepatocellular carcinoma cells. AdsSPARC (at MOI 100) induces high levels of secreted SPARC. (a) Western blot of conditioned media obtained from HepG2 cells following transduction with the different adenoviral vectors. Samples were collected at day 3 after cell transduction. (b) Immunofluorescent staining of HepG2 naïve cells. Fluorescence signals specific to SPARC antibody are visualized in green and nuclei, in blue (DAPI staining).

was not remarkable in terms of substrate-dependent cell growth.

We next wonder whether substrate-independent growth of HCC cells might be influenced by SPARC overexpression. To address this issue, HepG2 cells were grown in spheroids. Interestingly, and contrarily to what we have observed in monolayer-cultured HCC cells, a significant reduction in proliferation activity was observed in spheroids composed by AdsSPARC-treated cells (Fig. 3e). To exclude the partial involvement of indirect mechanisms mediated by cellular apoptosis, HepG2 spheroids were dissociated and stained with acridine orange-ethidium bromide mixture solution. No significant differences were found (not shown), which overall suggests that an increase in endogenous levels of SPARC in HCC cells reduces their substrate-independent proliferation capacity, therefore resulting in a partial inhibition of their tri-dimensional growth capacity.

Clonogenic and migratory capabilities of HCC cells are inhibited by SPARC over-expression. Possible implications on HCC cell aggressiveness

To assess whether SPARC overexpression in HCC cells might affect their colony formation capacity, a feature of cell aggressiveness, cells were plated at a density of 10^3 cells per 60-mm dish. After 15 days of incubation, the number of colonies was significantly decreased (16-fold) in cells with forced SPARC overexpression when compared with Ad- β gal- or AdasSPARC-treated cells (Fig. 4a).




Figure 2. Positive modulation of SPARC levels strongly inhibits HCC multicellular spheroid growth. (a) Spheroids prepared from HepG2 (upper panel) and Hep3B (lower panel) cells were transduced with AdsSPARC, AdasSPARC or Ad-βgal. Comparisons of spheroid mean volumes at days 2 and 6. Data expressed as the mean \pm SEM correspond to 3 experiments performed in 5 replicas. (*) vs. Ad-βgal. (*) <0.05 using Mann-Whitney test. (b) Photographs of the spheroids from HepG2 cells at day 6, taken under phase-contrast light microscopy. (c) Exogenous administration of SPARC as a recombinant protein did not affect spheroid growth on HepG2 cells. Kruskal-Wallis showed no significant differences among groups. [Color figure can be viewed in the online issue, which is available at www.interscience.wiley.com.]

Considering the ability of SPARC to modulate adhesiveness of certain tumor cells, we next wonder whether the increased SPARC expression in HCC cells could affect cell adhesiveness. For this purpose, cells were placed on fibronectin for 4 hr. AdsSPARC overexpressing cells showed a reduced ability to adhere to fibronectin (data not shown).

Migration is considered to be a critical mechanism in tumor cell dissemination. SPARC was previously shown to act as a chemotactic factor for prostate and breast carcinoma cell migration towards bone extracts^{47,48}; nevertheless, endogenous SPARC expression was reported to have opposite effects

in glioma cells⁴⁹ or to render no effects on MDA-MB-231 breast cancer cells.⁵⁰ Therefore, we decided to explore whether overexpression of SPARC might affect HCC cells chemotactic capacity towards TGF-β1. As shown in Figure 4b, AdsSPARC-HepG2 cells exhibited significantly less chemotactic migration capacity than cells transduced with Ad-βgal or untransduced cells. Similar results were obtained when Hep3B cells were used instead (not shown). Importantly, no differences in transwell membrane adherence of HepG2 cells were found among conditions (not shown). Therefore, the reduced migration capacity of SPARC-




Figure 3. SPARC upregulation does not profoundly affect HCC cell growth *in vitro*. (a) HCC cells (HepG2, Hep3B and Huh7) were incubated with no addition of viral vectors or with AdsSPARC, AdasSPARC or Ad-βgal for 3 days, at a MOI of 100. Cell viability was determined by MTT assay (Invitrogen) at 490 nm, in 3 independent studies; ns: non-significant. (*) $p < 0.05$ using Mann-Whitney test. (b) SPARC overexpression results in a non-significant increase in HepG2 cellular apoptosis when analyzed by the acridine-orange/ethidium bromide method. (*) $p < 0.05$ using Mann-Whitney test. (c) Flow cytometric analysis of HepG2 cells stained with annexin-V obtained after 72 hr in culture showed a slight significant induction in cellular apoptosis in SPARC overexpressing cells. Results are representative of 4 experiments similarly performed. (*) $p < 0.05$ using Mann-Whitney test. (d) Cell cycle analysis from DNA content. Cell cycle profiles correspond to one representative experiment of the 3 that were performed. Values are expressed as mean \pm SEM. (e) Proliferation activity of cells grown in spheroids, analyzed by MTT. (*) $p < 0.05$ using Student's *t* test.

overexpressing cells seems not to be principally related to differential adherence of HCC cells (Fig. 4b) upper. Thus, overall these data, including colony formation capacity, cell adhesiveness and cell migration capacity results, suggest that intracellular SPARC overexpression likely diminishes the aggressive-like behavior of HCC cells.

E-cadherins are involved in maintaining the epithelial structure of a number of tumor types and their down-regulation was associated with an increased invasiveness of human hepatocellular carcinomas.^{51,52} In addition, N-cad-

herin expression is frequently induced in highly aggressive tumors.⁴³ To further investigate the potential involvement of cadherins in SPARC-induced effects, changes in both E- and N-cadherin expression levels were investigated. Western blot analysis of AdsSPARC-HCC cells showed increased E-cadherin expression levels at day 3 (Fig. 5a). On the other hand, a 50% reduction in N-cadherin cell surface expression levels was observed in AdsSPARC-transduced HepG2 cells when compared to Ad-βgal-transduced or untransduced cells (Figs. 5b and 5c). A similar degree of N-cadherin inhibition was




Figure 4. SPARC profoundly influences colony formation and migratory capability in HCC cells. (a) AdsSPARC reduced HCC cell clonogenic capacity. For colony formation assay 10^3 untreated (Control) or AdsSPARC, AdasSPARC or Ad-βgal transduced (MOI 100) HCC cells (HepG2, Hep3B and Huh7) were cultured in a 6-well plate for 2 weeks before crystal violet staining, and the number of colonies (20–25 cells) was quantified under phase-contrast light microscopy (*) $p < 0.05$ vs. Control; Mann-Whitney test. (b) AdsSPARC inhibits migration of HepG2 cells towards TGF-β1. HepG2 cells were transduced with AdsSPARC or Ad-βgal or left untransduced for 3 days at a MOI of 100. Migration assays were performed in the absence of FBS as described in M&M. Representative pictures of transwell membrane adherence of HCC cells are shown (×200). For quantification, the average number of migrated cells per field was assessed. (***) $p < 0.001$ vs. Control and Ad-βgal; Student's *t* test. All assays were done in triplicates.

found in SPARC-overexpressing HCC cultured cells by immunocytochemistry (Figs. 5d–5h). To confirm the previous result, we have performed new experiments and immunostained cultures against α-SMA, another mesenchymal marker. Consistently, α-SMA expression was also inhibited in AdsSPARC-treated cells (Fig. 5i). From previous results, we conclude that SPARC overexpression in HepG2 cells results in the upregulation of E-cadherin levels and in the downregulation of N-cadherin as well as of α-SMA levels, which further suggest they have acquired a less aggressive phenotype. In addition, the finding of cell rounding in HCC overexpressing SPARC could be the result of its counteradhesive properties.

Upregulation of SPARC expression decreases HCC cells viability and increases their apoptosis in response to 5-FU-based chemotherapy

HCC cells are naturally resistant to any chemotherapeutic drug. One of the proposed mechanisms responsible for the observed antitumoral effects of SPARC is its ability to induce apoptosis of cancer cells.²⁹ To assess whether SPARC over-

pression might lead HCC cells more susceptible to chemotherapeutic agents, we tested the effects of 5-FU and adriamycin on HepG2 cells proliferation capacity and cell apoptosis. SPARC overexpression resulted in an increased sensitivity to 5-FU chemotherapy since both a 50% decreased cellular viability (Fig. 6a) and an increased apoptosis (~70% at 10 μg/ml 5-FU) (Fig. 6b) were observed. No changes were found when adriamycin was used as a chemotherapeutic agent instead (not shown).

The *In vivo* growth capacity of SPARC-overexpressing HCC cells is strongly inhibited in nude mice

We aimed to explore whether transient SPARC overexpression in 2 different HCC cell lines might affect their *in vivo* growth. Nude mice injected with control HepG2 cells showed an average tumor volume of 500–600 to 900 mm³ at day 40. Contrarily, transient overexpression of SPARC in HepG2 cells significantly inhibited tumor growth (reaching an average maximum tumor volume value of 30 mm³) (Fig. 7a) and significantly increased animal survival (log rank test $p < 0.05$) (Fig. 7b). No *in vivo* effects were observed on tumor




Figure 5. Gene transfer of SPARC induces mesenchymal-to-epithelial transition in hepatocellular carcinoma cells. (a) HepG2 cells were infected (at a MOI of 100) with AdSPARC, AdasSPARC or Ad-βgal for 3 days and whole-cell lysates were generated. E-cadherin expression was assessed by western blot. A representative of 3 independent experiments is shown. A goat antibody against actin was used as control of protein loading. (b) SPARC overexpression induces a decrease of N-cadherin expression in HepG2 cells. HepG2 cells were infected (at a MOI of 100) with AdSPARC or Ad-βgal for 3 days, fixed and incubated with antibodies to N-cadherin. By using fluorescence-activated flow cytometry analysis, N-cadherin expression is displayed by a shift in mean fluorescent intensity when compared with incubation without addition of any primary antibody. The percentages of N-cadherin positive cells are indicated. (c) Quantitative analysis from comparisons of results obtained from independent studies similar to that shown in b. (d–g) Representative pictures from N-cadherin immunostained HepG2 cells, which were previously transduced with Ad-Bgal (D,E) or AdSPARC (F,G) and plated onto poli-L-lysine/fibronectin coated coverslips. (h, i) Quantification of the N-cadherin (H) and α-SMA (I) immunostaining intensities in HepG2 cells is represented. The data are expressed as the Mean ± SEM. (*) vs. Control; (τ) vs. Ad-βgal. (*) $p < 0.05$; (ττ) $p < 0.01$; (***,σσσ) $p < 0.001$. Student's *t* test.

growth upon transient SPARC downregulation. Similar results although less potent resulted when Huh7 HCCs were used instead, with a consistent increased in animal survival (Figs. 7c and 7d). H&E staining of HepG2 tumors overexpressing SPARC showed a decreased rim of viable tumor cells in comparison with tumors of untransduced or Ad-βgal or AdasSPARC treated cells (Fig. 8a and not shown). In addition, we have observed a reduced number of cells with fibroblast morphology in the AdSPARC-HCC treated group in comparison with controls.

Over-expression of SPARC in HCC cells results in increased macrophage infiltration

Considering the importance of macrophages in the progression of tumors, we assessed their possible involvement in the *in vivo* observed SPARC effects. Interestingly, the number

of infiltrating macrophages (F4/80⁺ cells) was significantly higher in AdSPARC-HepG2 HCC tumors in comparison with control animal groups (F4/80 labeled area: 56463 ± 7209 vs. 17072 ± 2767 vs. 20785 ± 2634 pixels²; AdSPARC vs. Ad-βgal vs. Control; $p < 0.0001$; Fig. 8b). Significantly higher numbers of F4/80-positive cells were found in the core region of tumors from AdSPARC-treated animals than in those from mice injected with HepG2 or Ad-βgal HepG2 cells (Fig. 8d). This feature is likely a consequence of enhanced tumor cell apoptosis, since an increase in apoptosis induction was observed by TUNEL in SPARC overexpressing tumors (Fig. 8c).

Discussion

In this article, we provide for the first time, clear evidence that SPARC has a role on the invasive potential of HCC.




Figure 6. Up-regulation of SPARC expression decreases HCC cell viability in response to 5-FU-based chemotherapy and increases apoptosis. (a) HepG2 cells were infected with AdSPARC, Ad-βgal or left uninfected for 2 days, washed and further incubated with 5-FU for 24 hr. Cell viability was assessed by MTT assay (Invitrogen) at 490 nm. The data is expressed as the mean absorbance \pm SEM. (b) HepG2 cells were infected with AdSPARC, Adβgal or left uninfected for 2 days, washed and further incubated with 5-FU for 24 hr and stained with an AO/EB bromide mixture. Percentage of apoptotic cells is shown. The data are expressed as the Mean \pm SEM. (*) vs. Control; (σ) vs. Ad-βgal. (*,σ) $p < 0.05$; (**,σσ) $p < 0.01$; (***,σσσ) $p < 0.001$. Student's *t* test.

Transient, overexpression of SPARC in HCC cells transduction resulted in a phenotype indicative of decreased tumor cell aggressiveness. Moreover, the increase of endogenous SPARC expression levels in HepG2 cells resulted in: (i) reduced capacity to form 3-dimensional spheroids and decreased substrate-independent proliferation; (ii) decreased ability to migrate and to generate cellular colonies; (iii) increased expression of E-cadherin and a concomitant decrease in N-cadherin expression; (iv), increased susceptibility to 5-FU-based chemotherapy and (v) a potent inhibition of *in vivo* tumorigenicity leading to increased animal survival. The results from key experiments were also confirmed in other 2 HCC cell lines (Hep3B and Huh7) demonstrating that these findings do not correspond to a cell line specific phenomena.

HCC is a complex disease that often progresses as a consequence of genetic anomalies in cancer cells affecting many cell-growth regulatory pathways and also as a result of tumor cell interactions with microenvironmental factors.⁵³ In this context, we have addressed the role of SPARC, a matricellular glycoprotein with a complex regulatory function associated with increased aggressiveness in a number of human cancers.⁵⁴ There is contrasting evidence in the literature regarding SPARC influence on cancer cell behavior.⁵⁴ Although SPARC has been shown to be expressed by cancer cells, in certain tumors it is produced as well and at much higher levels by fibroblasts and endothelial cells.⁵⁵ SPARC is almost undetectable in normal hepatic tissue by immunohistochemistry, whereas substantial induction of its expression is observed in the HCC stroma.³⁷ Our data indicate that only endogenous increased levels of SPARC could exert a beneficial effect against HCC growth. Even though there is abundant information on the role of SPARC in other cancer types,²³ little is known about its involvement in hepatocarcinogenesis.

To analyze the effects of increasing SPARC expression levels on HCC cells we chose to take advantage of available gene therapy vectors.⁵⁶ Among them, replication deficient recombinant adenoviruses have been widely used for gene transfer to the liver and especially for HCC experimental treatment.^{57–59} In particular, type 5 adenoviruses very efficiently infect hepatic cells including HCC cells.^{59,60}

SPARC has been previously shown to induce the inhibition of cancer cells proliferation,⁶¹ arresting cells at G₀. The mechanisms by which SPARC inhibits proliferation were previously associated with alteration of growth factor signaling events through diverse mechanisms including interaction with PDGF receptors⁶² with the result of Mitogen-Activated Protein Kinase (MAPK) inhibition, cyclin E-Cyclin Dependent Kinase 2 (CDK2) inactivation, cyclin A down-regulation, and maintenance of RB activation.¹⁸ Antiproliferative effects of SPARC were also associated with indirect effects on IGF signals.⁶³ Interestingly, we show here that SPARC overexpression in HCC cells did not affect substrate-dependent cell proliferation. In addition, SPARC overexpression has been shown to slightly induce cellular apoptosis in HepG2 cells. It is noteworthy that the AdSPARC-mediated induction of SPARC inhibited spheroid formation of HCC cells, whereas its knock-down using SPARC antisense mRNA had no effects on spheroid growth. Moreover, when cultured as monolayer and even though cellular growth was unaffected by SPARC overexpression (not shown), a clear inhibitory effect on their clonogenic capacity was noted. Furthermore, no effects on spheroid growth were observed when recombinant SPARC was exogenously applied, indicating that SPARC effects on HCC tumor cells largely rely on the modulation of its cellular levels.

Migration is a key step for the invasive and disseminating capabilities of cancer cells. In our study, forced upregulation of SPARC expression inhibited migration of HCC cells




Figure 7. The *in vivo* growth capacity of SPARC overexpressing HCC cells is strongly inhibited in nude mice. AdsSPARC significantly increased animal survival. Nude mice were s.c. inoculated with 1.5×10^6 HepG2 (a) or 5×10^6 Huh7 (c) cells transduced with AdsSPARC or AdasSPARC or Ad-βgal or left untransduced and tumor size was measured with caliper twice a week over a period of 42 days. Effects of AdsSPARC treatment on survival of the tumor-bearing nude mice (b, HepG2 cells; d, Huh7 cells) (Kaplan-Meier survival curve). The data are expressed as the Mean \pm SEM. (*) vs. Control; (σ) vs. Ad-βgal. (*,σ) $p < 0.05$; (**,σσ) $p < 0.01$; (σσσ) $p < 0.001$. Mann-Whitney test.

towards TGF-β1, used as a chemoattractant. Although the mechanisms have been unknown, some evidences in glioma cell lines suggested that cell migration capability may depend on the type of extracellular matrix proteins present in their microenvironment.⁴¹

One of the key processes providing cancer cells with the capacity to migrate, invade and metastasize is their ability to undergo an epithelial to mesenchymal transition (EMT).⁶⁴ EMT is characterized by loss of intercellular adhesion (E-cadherin to N-cadherin switch),⁶⁴ downregulation of epithelial markers (cytokeratins), upregulation of MMP-2 and MMP-9 and the acquisition of a fibroblast-like motile phenotype. In fact, the induction of an EMT has been implicated in the malignant progression of HCC.^{41,42} We

observed that SPARC overexpression resulted in E-cadherin upregulation and N-cadherin downregulation as well as in a reduced expression of α-SMA protein altogether suggesting the involvement of mesenchymal to epithelial transition (MET) events. However, no significant changes were observed on MMP-2 and MMP-9 activities which might be due to low frequency of EMT events in HepG2 cells as suggested by the low levels of N-cadherin expression found in this specific cell line (not shown). These data are in contrast with our and others previous observations in melanoma cells, in which high levels of SPARC are associated with N-cadherin upregulation and E-cadherin downregulation^{65,66}; however, they are consistent with the increase of tumor aggressiveness observed in other melanoma models when




Figure 8. Host macrophage cells are involved in the rejection of SPARC overexpressing HCC cells. Nude mice were s.c. inoculated with 1.5×10^6 HepG2 cells transduced with AdsSPARC, AdasSPARC, Ad β gal or left untransduced. (a) H&E staining showing a reduced tumoral rim in mice from AdsSPARC group. Arrows: fibroblastic-like cells; dotted arrows: tumor cells. (b) Representative photographs of immunohistochemical staining of macrophages (F4/80) in HepG2 tumors at 5 days after inoculation, corresponding to HepG2 (upper, $\times 100$; lower, $\times 400$) and AdsSPARC-HepG2 (upper, $\times 100$; lower, $\times 400$) treated mice. (c) Quantitative graph showing comparisons of the F4/80 immunostained area in different experimental conditions. The data are expressed as the Mean \pm SEM. (*) vs. Control; (σ) vs. Ad β gal. (*, σ) $p < 0.05$; (**, $\sigma\sigma$) $p < 0.01$; (***, $\sigma\sigma\sigma$) $p < 0.001$. Mann-Whitney test. (d) SPARC increases *in vivo* HCC apoptosis. The effect of SPARC overexpression on apoptosis was assessed by TUNEL assay. Representative photographs of TUNEL assay in HepG2 tumors at day 5 after treatment, corresponding to Ad β gal, AdsSPARC, AdasSPARC or untransduced HepG2 cells.

SPARC is overexpressed^{30,31} Thus, SPARC overexpression in different kinds of cancer cells seems to be differently involved in modulating EMT or MET, likely affecting cells

migratory properties. Together, our data suggest that SPARC overexpression in HCC cells induces a MET in HCC cells, a mechanism which is likely involved in differences in cell behavior hereby shown, which are indicative of a less aggressive-like phenotype.

Many chemotherapeutic agents have been applied in the treatment of HCC.⁶⁷ High doses of chemotherapeutic drugs are necessary to obtain a low rate of response. One of the chemotherapeutic agents most extensively studied is 5-FU, a pyrimidine antimetabolite which has an overall response rate lower than 10%.⁶⁷ Chemotherapy in cirrhotic patients is poorly tolerated and it is no longer applied to patients with advanced or metastatic HCC.⁶⁷ It is clear that there is a need for new nontoxic agents or new strategies that can re-sensitize HCC cells to standard chemotherapy. On the basis of the present data SPARC appears to function as a tumor suppressor in some cancer cells and one of the mechanisms is the induction of cellular apoptosis. For instance, SPARC has been shown to reverse chemotherapy resistance in colorectal carcinoma cells through induction of caspase-8 activation.²⁹ Consistent with those results, we found that SPARC overexpression in HCC cells resulted in a decreased proliferation and increased apoptosis when cells were subjected to lower concentrations of 5-FU. This evidence provides a proof-of-principle that the reversal of resistance to 5-FU-based chemotherapy can potentially be exploited therapeutically for HCC. Thus, increasing SPARC expression on HCC cells through gene therapy could be an attractive strategy to render cells more susceptible to standard chemotherapy or to combined treatments.

Confirming previous results,²² nude animals inoculated with SPARC-overexpressing HCC cells showed a strong inhibition of tumor and significant increase in long-term survival when compared to controls. In agreement with *in vitro* effects, knock-down of SPARC had no effects on *in vivo* tumor growth. It still remains unclear to a certain extent why SPARC is able to inhibit tumor growth in certain types of tumor while in others it has the opposite effect.²³ The timing of SPARC expression might partially explain those contrasting results; thus, in our work, high but transient SPARC overexpression seems to be a sufficient stimulus to decrease the aggressive-like behavior of HCC cells. The induction in E-cadherin and downregulation of N-cadherin by SPARC overexpression might be partially involved in the decreased growth ability of tumor cells likely through interfering with other microenvironmental factors in the cell-cell context. We observed that the density of host tumor invading macrophages is increased in AdsSPARC-HCC cells-treated animals. This feature might be the consequence of *in vivo* apoptosis induction in SPARC overexpressing HCC cells and/or it could be an active event mediated by the enhancement in SPARC expression. Another particular observation emerging from histological analysis of HCC tumors is the reduced number of cells with fibroblast morphology in the AdsSPARC-HCC treated group, suggesting that SPARC might have a role in extracellular matrix

organization and/or fibroblast invasiveness to tumor mass periphery. On the other hand, only mild changes in tumor vasculature were observed after CD31 or von-Willebrand immunostainings, at 5 days after tumor cell injection, although it is well known that SPARC has antiangiogenic properties.

In summary, our study implicates SPARC as a HCC tumor modulator protein able to inhibit volumetric tumor growth both *in vitro* and *in vivo* and to influence several other morphological and behavioral cellular features intimately linked to cancer cell aggressiveness. In addition, we show that SPARC overexpression in HCC cells likely increase

tumor cells sensitivity to 5-FU-based chemotherapy. For previous reasons, the enhancement of SPARC expression in HCC cells is proposed as a potential strategy for the treatment of HCC, a disease in which other therapeutic approaches are nowadays mostly neglected.

Acknowledgements

We acknowledge the continuous support from Ms. Inés Bemberg. We thank Ms. Soledad Arregui and Mr. Guillermo Gastón for their expert technical assistance; Dr. Fernando Pitossi (Fundación Instituto Leloir) for gently sharing some reagents and Mr. Leonardo Sganga for flow cytometry analysis.

References

- Parkin DM, Bray F, Ferlay J, Pisani P. Global cancer statistics, 2002. *CA Cancer J Clin* 2005;55:74–108.
- El-Serag HB, Mason AC. Rising incidence of hepatocellular carcinoma in the United States. *N Engl J Med* 1999;340:745–50.
- El-Serag HB, Marrero JA, Rudolph L, Reddy KR. Diagnosis and treatment of hepatocellular carcinoma. *Gastroenterology* 2008;134:1752–63.
- Llovet JM, Burroughs A, Bruix J. Hepatocellular carcinoma. *Lancet* 2003;362:1907–17.
- Llovet JM, Di Bisceglie AM, Bruix J, Kramer BS, Lencioni R, Zhu AX, Sherman M, Schwartz M, Lotze M, Talwalkar J, Gores GJ, for the Panel of Experts in HCCDCT. Design and endpoints of clinical trials in hepatocellular carcinoma. *J Natl Cancer Inst* 2008;100:698–711.
- Liotta LA, Kohn EC. The microenvironment of the tumour-host interface. *Nature* 2001;411:375–9.
- Termine JD, Kleinman HK, Whitson SW, Conn KM, McGarvey ML, Martin GR. Osteonectin, a bone-specific protein linking mineral to collagen. *Cell* 1981;26:99–105.
- Sage EH, Bornstein P. Extracellular proteins that modulate cell-matrix interactions. SPARC, tenascin, and thrombospondin. *J Biol Chem* 1991;266:14831–4.
- Bradshaw AD, Sage EH. SPARC, a matricellular protein that functions in cellular differentiation and tissue response to injury. *J Clin Invest* 2001;107:1049–54.
- Mann K, Deutzmann R, Paulsson M, Timpl R. Solubilization of protein BM-40 from a basement membrane tumor with chelating agents and evidence for its identity with osteonectin and SPARC. *FEBS Lett* 1987;218:167–72.
- Funk SE, Sage EH. Differential effects of SPARC and cationic SPARC peptides on DNA synthesis by endothelial cells and fibroblasts. *J Cell Physiol* 1993;154:53–63.
- Barker TH, Baneyx G, Cardo-Vila M, Workman GA, Weaver M, Menon PM, Dedhar S, Rempel SA, Arap W, Pasqualini R, Vogel V, Sage EH. SPARC regulates extracellular matrix organization through its modulation of integrin-linked kinase activity. *J Biol Chem* 2005;280:36483–93.
- Barker TH, Baneyx G, Cardo-Vila M, Workman GA, Weaver M, Menon PM, Dedhar S, Rempel SA, Arap W, Pasqualini R, Vogel V, Sage EH. SPARC regulates extracellular matrix organization through its modulation of integrin-linked kinase activity. *J Biol Chem* 2005;280:36483–93. Epub 2005 Aug 22.
- Brekken RA, Sage EH. SPARC, a matricellular protein: at the crossroads of cell-matrix communication. *Matrix Biol* 2001;19:816–27.
- Sage EH, Bornstein P. Extracellular proteins that modulate cell-matrix interactions. SPARC, tenascin, and thrombospondin. *J Biol Chem* 1991;266:14831–4.
- Yunker CK, Golembieski W, Lemke N, Schultz CR, Cazacu S, Brodie C, Rempel SA. SPARC-induced increase in glioma matrix and decrease in vascularity are associated with reduced VEGF expression and secretion. *Int J Cancer* 2008;122:2735–43.
- Hasselaar P, Sage EH. SPARC antagonizes the effect of basic fibroblast growth factor on the migration of bovine aortic endothelial cells. *J Cell Biochem* 1992;49:272–83.
- Motamed K, Funk SE, Koyama H, Ross R, Raines EW, Sage EH. Inhibition of PDGF-stimulated and matrix-mediated proliferation of human vascular smooth muscle cells by SPARC is independent of changes in cell shape or cyclin-dependent kinase inhibitors. *J Cell Biochem* 2002;84:759–71.
- Porte H, Chastre E, Prevot S, Nordlinger B, Empereur S, Basset P, Chambon P, Gspach C. Neoplastic progression of human colorectal cancer is associated with overexpression of the stromelysin-3 and BM-40/SPARC genes. *Int J Cancer* 1995;64:70–5.
- Porte H, Triboulet JP, Kotelevets L, Carrat F, Prevot S, Nordlinger B, DiGioia Y, Wurtz A, Comoglio P, Gspach C, Chastre E. Overexpression of stromelysin-3, BM-40/SPARC, and MET genes in human esophageal carcinoma: implications for prognosis. *Clin Cancer Res* 1998;4:1375–82.
- Ledda MF, Adris S, Bravo AI, Kairiyama C, Bover L, Chernajovsky Y, Mordoh J, Podhajcer OL. Suppression of SPARC expression by antisense RNA abrogates the tumorigenicity of human melanoma cells. *Nat Med* 1997;3:171–6.
- Lau CP, Poon RT, Cheung ST, Yu WC, Fan ST. SPARC and Hevin expression correlate with tumour angiogenesis in hepatocellular carcinoma. *J Pathol* 2006;210:459–68.
- Podhajcer OL, Benedetti LG, Girotti MR, Prada F, Salvatierra E, Llera AS. The role of the matricellular protein SPARC in the dynamic interaction between the tumor and the host. *Cancer Metastasis Rev* 2008;27:691–705.
- Chlenski A, Liu S, Crawford SE, Volpert OV, DeVries GH, Evangelista A, Yang Q, Salwen HR, Farrer R, Bray J, Cohn SL. SPARC is a key Schwannian-derived inhibitor controlling neuroblastoma tumor angiogenesis. *Cancer Res* 2002;62:7357–63.
- Sato N, Fukushima N, Maehara N, Matsubayashi H, Koopmann J, Su GH, Hruban RH, Goggins M. SPARC/osteonectin is a frequent target for aberrant methylation in pancreatic adenocarcinoma and a mediator of tumor-stromal interactions. *Oncogene* 2003;22:5021–30.
- Said N, Motamed K. Absence of host-secreted protein acidic and rich in cysteine (SPARC) augments peritoneal ovarian carcinomatosis. *Am J Pathol* 2005;167:1739–52.
- Cheetham S, Tang MJ, Mesak F, Kennecke H, Owen D, Tai IT. SPARC promoter hypermethylation in colorectal cancers can be reversed by 5-Aza-2'-deoxycytidine to increase SPARC expression and improve

- therapy response. *Br J Cancer* 2008;98:1810–19.
28. DiMartino JF, Lacayo NJ, Varadi M, Li L, Saraiya C, Ravindranath Y, Yu R, Sikic BI, Raimondi SC, Dahl GV. Low or absent SPARC expression in acute myeloid leukemia with MLL rearrangements is associated with sensitivity to growth inhibition by exogenous SPARC protein. *Leukemia* 2006;20:426–32.
 29. Tang MJ, Tai IT. A novel interaction between procaspase 8 and SPARC enhances apoptosis and potentiates chemotherapy sensitivity in colorectal cancers. *J Biol Chem* 2007;282:34457–67.
 30. Massi D, Franchi A, Borgognoni L, Reali UM, Santucci M. Osteonectin expression correlates with clinical outcome in thin cutaneous malignant melanomas. *Hum Pathol* 1999;30:339–44.
 31. Ledda F, Bravo AI, Adris S, Bover L, Mordoh J, Podhajcer OL. The expression of the secreted protein acidic and rich in cysteine (SPARC) is associated with the neoplastic progression of human melanoma. *J Invest Dermatol* 1997;108:210–14.
 32. Golembieski WA, Ge S, Nelson K, Mikkelsen T, Rempel SA. Increased SPARC expression promotes U87 glioblastoma invasion in vitro. *Int J Dev Neurosci* 1999;17:463–72.
 33. Thomas R, True LD, Bassuk JA, Lange PH, Vessella RL. Differential expression of osteonectin/SPARC during human prostate cancer progression. *Clin Cancer Res* 2000;6:1140–9.
 34. Minn AJ, Gupta GP, Siegel PM, Bos PD, Shu W, Giri DD, Viale A, Olshen AB, Gerald WL, Massague J. Genes that mediate breast cancer metastasis to lung. *Nature* 2005;436:518–24.
 35. Nakatani K, Seki S, Kawada N, Kitada T, Yamada T, Sakaguchi H, Kadoya H, Ikeda K, Kaneda K. Expression of SPARC by activated hepatic stellate cells and its correlation with the stages of fibrogenesis in human chronic hepatitis. *Virchows Arch* 2002;441:466–74. Epub 2002 Apr 26.
 36. Camino AM, Atorrasagasti C, Maccio D, Prada F, Salvatierra E, Rizzo M, Alaniz L, Aquino JB, Podhajcer OL, Silva M, Mazzolini G. Adenovirus-mediated inhibition of SPARC attenuates liver fibrosis in rats. *J Gene Med* 2008;10:993–1004.
 37. Le Bail B, Faouzi S, Boussarie L, Guirouilh J, Blanc JF, Carles J, Bioulac-Sage P, Balabaud C, Rosenbaum J. Osteonectin/SPARC is overexpressed in human hepatocellular carcinoma. *J Pathol* 1999;189:46–52.
 38. Goldenberg D, Ayes S, Schneider T, Pappo O, Jurim O, Eid A, Fellig Y, Dadon T, Ariel I, de Groot N, Hochberg A, Galun E. Analysis of differentially expressed genes in hepatocellular carcinoma using cDNA arrays. *Mol Carcinog* 2002;33:113–24.
 39. Wheelock MJ, Johnson KR. Cadherins as modulators of cellular phenotype. *Annu Rev Cell Dev Biol* 2003;19:207–35.
 40. Takeichi M. Cadherins in cancer: implications for invasion and metastasis. *Curr Opin Cell Biol* 1993;5:806–11.
 41. Kozyraki R, Scoazec JY, Flejou JF, D'Errico A, Bedossa P, Terris B, Fiorentino M, Bringuier AF, Grigioni WF, Feldmann G. Expression of cadherins and alpha-catenin in primary epithelial tumors of the liver. *Gastroenterology* 1996;110:1137–49.
 42. Wei Y, Van Nhieu JT, Prigent S, Srivatanakul P, Tiollais P, Buendia MA. Altered expression of E-cadherin in hepatocellular carcinoma: correlations with genetic alterations, beta-catenin expression, and clinical features. *Hepatology* 2002;36:692–701.
 43. Hazan RB, Qiao R, Keren R, Badano I, Suyama K. Cadherin switch in tumor progression. *Ann NY Acad Sci* 2004;1014:155–63.
 44. Reed LJ MH. A simple methods of estimating fifty per cent points. *Am J Hyg* 1938;27:493–7.
 45. Bradford MM. A rapid and sensitive method for the quantitation of microgram quantities of protein utilizing the principle of protein-dye binding. *Anal Biochem* 1976;72:248–54.
 46. Aquino JB, Musolino PL, Coronel MF, Villar MJ, Setton-Avruj CP. Nerve degeneration is prevented by a single intraneural apotransferrin injection into colchicine-injured sciatic nerves in the rat. *Brain Res* 2006;1117:80–91.
 47. Jacob K, Webber M, Benayahu D, Kleinman HK. Osteonectin promotes prostate cancer cell migration and invasion: a possible mechanism for metastasis to bone. *Cancer Res* 1999;59:4453–7.
 48. Campo McKnight DA, Sosnoski DM, Koblinski JE, Gay CV. Roles of osteonectin in the migration of breast cancer cells into bone. *J Cell Biochem* 2006;97:288–302.
 49. Seno T, Harada H, Kohno S, Teraoka M, Inoue A, Ohnishi T. Downregulation of SPARC expression inhibits cell migration and invasion in malignant gliomas. *Int J Oncol* 2009;34:707–15.
 50. Koblinski JE, Kaplan-Singer BR, VanOsdol SJ, Wu M, Engbring JA, Wang S, Goldsmith CM, Piper JT, Vostal JG, Harms JF, Welch DR, Kleinman HK. Endogenous osteonectin/SPARC/BM-40 expression inhibits MDA-MB-231 breast cancer cell metastasis. *Cancer Res* 2005;65:7370–7.
 51. Liu J, Lian Z, Han S, Wayne MM, Wang H, Wu MC, Wu K, Ding J, Arbuthnot P, Kew M, Fan D, Feitelson MA. Downregulation of E-cadherin by hepatitis B virus X antigen in hepatocellular carcinoma. *Oncogene* 2006;25:1008–17.
 52. Endo K, Ueda T, Ueyama J, Ohta T, Terada T. Immunoreactive E-cadherin, alpha-catenin, beta-catenin, and gamma-catenin proteins in hepatocellular carcinoma: relationships with tumor grade, clinicopathologic parameters, and patients' survival. *Hum Pathol* 2000;31:558–65.
 53. Avila MA, Berasain C, Sangro B, Prieto J. New therapies for hepatocellular carcinoma. *Oncogene* 2006;25:3866–84.
 54. Podhajcer OL, Benedetti L, Girotti MR, Prada F, Salvatierra E, Llera AS. The role of the matricellular protein SPARC in the dynamic interaction between the tumor and the host. *Cancer Metastasis Rev* 2008;27:523–37.
 55. Sage H, Johnson C, Bornstein P. Characterization of a novel serum albumin-binding glycoprotein secreted by endothelial cells in culture. *J Biol Chem* 1984;259:3993–4007.
 56. Prieto J, Qian C, Hernandez-Alcoceba R, Gonzalez-Aseguinolaza G, Mazzolini G, Sangro B, Kramer MG. Gene therapy of liver diseases. *Expert Opin Biol Ther* 2004;4:1073–91.
 57. Wilson JM. Adenovirus-mediated gene transfer to liver. *Adv Drug Deliv Rev* 2001;46:205–9.
 58. Ilan Y, Saito H, Thummala NR, Chowdhury NR. Adenovirus-mediated gene therapy of liver diseases. *Semin Liver Dis* 1999;19:49–59.
 59. Yu Q, Que LG, Rockey DC. Adenovirus-mediated gene transfer to nonparenchymal cells in normal and injured liver. *Am J Physiol Gastrointest Liver Physiol* 2002;282:G565–G72.
 60. Garcia-Banuelos J, Siller-Lopez F, Miranda A, Aguilar LK, Aguilar-Cordova E, Armendariz-Borunda J. Cirrhotic rat livers with extensive fibrosis can be safely transduced with clinical-grade adenoviral vectors. Evidence of cirrhosis reversion. *Gene Ther* 2002;9:127–34.
 61. Framson PE, Sage EH. SPARC and tumor growth: where the seed meets the soil? *J Cell Biochem* 2004;92:679–90.
 62. Raines EW, Lane TF, Iruela-Arispe ML, Ross R, Sage EH. The extracellular glycoprotein SPARC interacts with platelet-derived growth factor (PDGF)-AB and -BB and inhibits the binding of PDGF to its receptors. *Proc Natl Acad Sci USA* 1992;89:1281–5.
 63. Francki A, Motamed K, McClure TD, Kaya M, Murri C, Blake DJ, Carbon JG, Sage EH. SPARC regulates cell cycle progression in mesangial cells via its inhibition of IGF-dependent signaling. *J Cell Biochem* 2003;88:802–11.

64. Thiery JP. Epithelial-mesenchymal transitions in tumour progression. *Nat Rev Cancer* 2002;2:442–54.
65. Sosa MS, Girotti MR, Salvatierra E, Prada F, de Olmo JA, Gallango SJ, Albar JP, Podhajcer OL, Llera AS. Proteomic analysis identified N-cadherin, clusterin, and HSP27 as mediators of SPARC (secreted protein, acidic and rich in cysteines) activity in melanoma cells. *Proteomics* 2007;7:4123–34.
66. Robert G, Gaggioli C, Bailet O, Chavey C, Abbe P, Aberdam E, Sabatie E, Cano A, Garcia de Herreros A, Ballotti R, Tartare-Deckert S. SPARC represses E-cadherin and induces mesenchymal transition during melanoma development. *Cancer Res* 2006;66:7516–23.
67. Bruix J, Sherman M, Llovet JM, Beaugrand M, Lencioni R, Burroughs AK, Christensen E, Pagliaro L, Colombo M, Rodes J. Clinical management of hepatocellular carcinoma. Conclusions of the Barcelona-2000 EASL conference. European Association for the Study of the Liver. *J Hepatol* 2001;35:421–30.