BEHAVIOR OF THE SOLUTION OF THE TWO-PHASE STEFAN PROBLEM WITH REGARD TO THE CHANGING OF THERMAL COEFFICIENTS OF THE SUBSTANCE

M. C. $OLGUIN^{\dagger}$, M. C. $SANZIEL^{\ddagger}$ and D. A. $TARZIA^{\ast}$

†Dto. de Matemática, Fac. de Cs. Exactas, Ing. y Agrimensura, Univ. Nac. de Rosario,

(S2000BTP) Rosario, Argentina. mcolguin@fceia.unr.edu.ar

‡Consejo de Investigaciones, Dto. de Matemática, Fac. de Cs. Exactas, Ing. y Agrimensura,

Univ. Nac. de Rosario, (S2000BTP) Rosario, Argentina. sanziel@fceia.unr.edu.ar

* Dto. de Matemática FCE, Univ. Austral-CONICET, (S2000FZF) Rosario, Argentina

DTarzia@austral.edu.ar

Abstract— We consider one-dimensional two-phase Stefan problems for a finite substance with different boundary conditions at the fixed faces. The goal of this paper is to determine the behavior of the free boundary and the temperature when the thermal coefficients of the material change.

We obtain properties of monotony with respect to the latent heat, the common mass density, the specific heat of each phase and the thermal conductivity of the liquid phase.

We show that the solution is not monotone with respect to the thermal conductivity of solid phase, in some cases, by computing a numerical solution through a finite difference scheme.

The results obtained are important in technological applications as the climate of buildings, the storage of energy in satellites and clothes and the transport of biological substances and telecommunications.

Keywords— Phase change material, Two phase Stefan problem, Finite difference method

I. INTRODUCTION

Several technological applications can be modeled through heat transfer problems with phase-change. The Phase Change Materials (PCMs) are substances whose phase-change temperature make them available to moderate oscillations of temperature and to store energy of another substance in contact with them (Jiji and Gaye, 2006). For this reason PCMs are used for the climate of buildings, the storage of energy in satellites and clothes and the transport of biological substances (Asako *et al.*, 2002; Lamberg and Sirn, 2003), among a variety of applications.

Usually, the technological way to select a PCM is through its phase-change temperature, but when there

are several products in the correct range of temperature, it is necessary to study another properties of the substance, for example the thermal coefficients, in order to choose the most convenient PCM.

When we consider a packaging of a PCM that recovers an organic substance to be transported, it is essential to find the thickness of this pack to insure the optimal temperature of conservation in the organic substance during total time of transport (Bouciguez *et al.*, 2001; Farid *et al.*, 2004; Medina *et al.*, 2004; Zalba *et al.*, 2003; Zivkovic and Fujii, 2001). Because the sizes of the pack (wide, length and height) are sufficiently greater than its thickness, we can assume that the heat transfer occurs in only one direction. Moreover, if we consider a material with one portion at solid state and the other at the liquid state, and take into account the environment conditions, we have an one dimensional two-phase Stefan problem (Lamberg, 2004; Lamberg *et al.*, 2004).

In Olguín *et al.* (2007), we considered a onedimensional one-phase Stefan problem for fusion of a semi-infinite material. We showed that, both, the temperature and the free boundary present a monotonous behavior with respect to the latent heat, the mass density and the specific heat. We also showed that the solution is not monotone with respect to the thermal conductivity.

At the present work we continue this line of research. We consider several two-phase one-dimensional Stefan problems for a finite material, with different boundary conditions at the two fixed faces, in order to determine the behavior of the solutions when the thermal coefficients change.

In Section 2 we study a phase-change problem with temperature specification on both fixed faces of the finite material and we obtain results of monotony for latent heat, mass density, specific heat of the solid and the liquid phase, and thermal conductivity of the liquid phase, by using the maximum principle (Protter and Weinberger, 1967). Analogous results are obtained in Section 3, by considering heat flux specification at both fixed faces of the finite material and, in Section 4, with temperature specification at the left fixed face and a heat flux specification at the right fixed face.

When we consider the thermal conductivity of the solid phase, it was not possible to establish analytical results of monotony. In order to obtain some conclusion for this case, in Section 5 we consider a numerical solution. Since the free boundary change with time, the domain of the problems is variable, then it is necessary to develop a particular scheme with a time variable mesh. We can show that, in some cases, the solutions are not monotone.

II. PROBLEM WITH TEMPERATURE BOUNDARY SPECIFICATION AT BOTH FACES

A. Mathematical problem and preliminary results

We consider a finite material represented by the interval [0, 1]. At the initial time, we suppose that one portion of the material is at solid phase and the other is at liquid phase. If we consider that s(t) is the position of the free boundary at each time and

$$w(x,t) = \begin{cases} u(x,t) & 0 \le x \le s(t), \ 0 \le t \le T, \\ v(x,t) & s(t) \le x \le 1, \ 0 \le t \le T, \end{cases}$$

is the temperature of the material, the problem (P1)
is to find functions $w(x,t) = s(t)$ and a time $T > 0$, so

is to find functions w(x,t), s(t) and a time T > 0, so that they satisfy the following conditions:

$$\alpha_l u_{xx} = u_t, \quad 0 < x < s(t), 0 < t < T,$$
 (1)

$$\alpha_s v_{xx} = v_t, \quad s(t) < x < 1, 0 < t < T,$$
 (2)

$$u(0,t) = f(t) > 0, \quad 0 < t < T,$$
 (3)

$$v(1,t) = g(t) < 0, \quad 0 < t < T,$$
 (4)

$$u(s(t), t) = v(s(t), t) = 0, \quad 0 < t < T,$$
 (5)

$$k_s v_x(s(t), t) - k_l u_x(s(t), t) = \rho \ell s'(t), \ 0 < t < T, \ (6)$$

$$\iota(x,0) = \varphi(x) \ge 0, \quad 0 \le x \le b, \tag{7}$$

$$\psi(x,0) = \psi(x) \le 0, \quad b \le x \le 1,$$
 (8)

$$s(0) = b, \quad 0 < b < 1,$$
 (9)

where $\alpha_i = \frac{k_i}{\rho c_i}$ is the thermal diffusivity of the phase $i \ (i = s, l)$ and $\varphi(b) = 0 = \psi(b)$ (see Fig.1).

In Cannon and Primicerio (1971), Tarzia (1987) and Cannon (1984), under suitable hypothesis for data, it was proved that:

*) there is a unique solution for problem (P1) for all T > 0;

**) the Stefan condition (Eq.6) is equivalent to the integral equation:

$$s(t) = b + \int_0^1 \Phi(x) dx - \frac{c_l}{\ell} \int_0^{s(t)} u(x, t) dx$$



Figure 1: Scheme of the problem (P1)

$$-\frac{c_s}{\ell} \int_{s(t)}^1 v(x,t) dx + \frac{k_s}{\rho \ell} \int_0^t v_x(1,\tau) d\tau$$
$$-\frac{k_l}{\rho \ell} \int_0^t u_x(0,\tau) d\tau t \ge 0$$
(10)

where

$$\Phi(x) = \begin{cases} \frac{c_l}{\ell} \varphi(x), & 0 \le x \le b, \\ \frac{c_s}{\ell} \psi(x), & b < x \le 1. \end{cases}$$

B. Analytical results of monotony

In this Section we use the maximum principle, Hopf's lemma (Protter and Weinberger, 1967; Cannon, 1984) and the result of the following Lemma, when it is required, in order to establish some properties of monotony for the solution of problem (P1).

Lemma 1 Let w be the solution of (P1). Then: i) $u_x(s(t),t) < 0$ and $v_x(s(t),t) < 0$ for t > 0;

ii) If the function f(t) is continuously differentiable for t > 0 and $\varphi(x)$ is twice continuously differentiable at 0 < x < b, $f'(t) \leq 0$ and $\varphi''(x) \geq 0$ then $u_{xx}(x,t) \geq 0$ for 0 < x < s(t), 0 < t < T.

iii) If g(t) is continuously differentiable for t > 0and $\psi(x)$ is twice continuously differentiable at b < x < 1, $g'(t) \ge 0$ and $\psi''(x) \ge 0$ then $v_{xx}(x,t) \ge 0$ for s(t) < x < 1, 0 < t < T.

Proof.

i) The results are obtained when the maximum principle and Hopf's lemma are applied to functions u and v respectively.

ii) We define the auxiliary function $z = u_{xx}$ which verify the associated problem

$$\begin{aligned} z_t &= \alpha_l \, z_{xx}, & 0 < x < s(t), \, t > 0, \\ z(0,t) &= \frac{1}{\alpha_l} \, f'(t) & t > 0, \\ z(x,0) &= \varphi''(x) & 0 < x < b, \\ z(s(t),t) &\ge 0, & t > 0, \\ s(0) &= b. \end{aligned}$$

By using the maximum principle, we obtain $z \ge 0$. *iii*) The proof follows (*ii*) by defining now the aux*iliary* function $z = v_{xx}$. **Proposition 2** The solution $\{w(x,t), s(t)\}$ of problem (P1) depends monotonically on the latent heat ℓ , the mass density ρ , the specific heat c_i (i = s, l) and the thermal conductivity of the liquid phase k_l .

Proof. The five results are similar but the proofs are different because we must change the coefficients only in the Stefan condition for the latent heat ℓ , only in the heat equation for the specific heats $c_i(i = s, l)$ and in both equations in the remaining cases.

a) We consider $\ell_1 < \ell_2$. We call $\{w_1, s_1\}, \{w_2, s_2\}$ the solutions of problem (P1) with $\ell = \ell_i$ and $s_i(0) = b_i$ for i = 1, 2. We consider two cases: $s_1(0) > s_2(0)$ and $s_1(0) \ge s_2(0)$.

Case I: Let $b_1 = s_1(0) > s_2(0) = b_2$. We suppose t_0 is the first time such that $s_1(t_0) = s_2(t_0)$ and $s_2(t) < s_1(t)$ for all $0 < t < t_0$. Because of that, it must occur (Fig.2):

$$\dot{s}_{1}(t_{0}) \leq \dot{s}_{2}(t_{0})$$
 (11)

If we define the functions

 $\int U(x,t) = u_1(x,t) - u_2(x,t), \ 0 \le x \le s_2(t),$

 $V(x,t) = v_1(x,t) - v_2(x,t), \, s_1(t) \le x \le 1,$

for $0 \le t \le t_0$, then U satisfies the following conditions:

$$U_t - \alpha_l U_{x,x} = 0, \quad 0 < x < s_2(t), \quad 0 < t < t_0, \ (12)$$

$$U(0,t) = 0, \quad 0 < t < t_0, \tag{13}$$

$$U(x,0) = 0 \quad 0 \le x \le b_2, \tag{14}$$

$$U(s_2(t), t) = u_1(s_2(t), t) > 0, \quad 0 < t < t_0, \quad (15)$$

$$b_2(0) = b_2.$$
 (16)

By the minimum principle, condition (Eq.15) holds and then $U \ge 0$ in $0 \le x \le s_2(t)$, $0 \le t \le t_0$. At the point $x = s_1(t_0) = s_2(t_0)$, we have $U(s_2(t_0), t_0) = 0$, therefore U attains a minimum at $(s_2(t_0), t_0)$. By the Hopf's lemma, we have

$$U_x(s_2(t_0), t_0) < 0. \tag{17}$$

On the other hand, the function V, satisfies the following conditions:

$$V_t - \alpha_s V_{xx} = 0, \quad s_1(t) < x < 1, \quad 0 < t < t_0,$$
(18)

$$V(s_1(t), t) = -v_2(s_1(t), t) > 0, \quad 0 < t < t_0, \quad (19)$$

$$V(x,0) = 0, \quad b_1 \le x \le 1, \tag{20}$$

 $V(1,t) = 0, \quad 0 < t < t_0, \tag{21}$

$$s_1(0) = b_1.$$
 (22)

In a similar way as above, we obtain that $V(x,t) \ge 0$ in $s_1(t) \le x \le 1$, $0 \le t \le t_0$. Taking into account that $V(s_1(t_0), t_0) = 0$, then V has a minimum at $(s_1(t_0), t_0)$ and from the Hopf's lemma, it results

$$V_x(s_1(t_0), t_0) > 0. (23)$$

But, from the Stefan condition (Eq.6), (Eq.11) and (Eq.17) we obtain:



Figure 2: Free boundaries for $\ell_1 < \ell_2$

 $V_{x}(s_{1}(t_{0}), t_{0}) = v_{1x}(s_{1}(t_{0}), t_{0}) - v_{2x}(s_{2}(t_{0}), t_{0})$ $= \frac{\rho}{k_{s}} \left[\ell_{1}s_{1}'(t_{0}) - \ell_{2}s_{2}'(t_{0}) \right] + \frac{k_{l}}{k_{s}} U_{x}(s_{2}(t_{0}), t_{0})$ $< \frac{\rho\ell_{1}}{k_{s}} \left[s_{1}'(t_{0}) - s_{2}'(t_{0}) \right] + \frac{k_{l}}{k_{s}} U_{x}(s_{2}(t_{0}), t_{0}) < 0,$ high is a constant diction with $(T_{s}, 22)$. Then one of

which is a contradiction with (Eq.23). Then we conclude that:

 $\begin{cases} s_1(t) > s_2(t), \\ u_1(x,t) \ge u_2(x,t), & 0 \le x \le s_2(t), \\ v_1(x,t) \ge v_2(x,t), & s_1(t) \le x \le 1 \\ for \ 0 \le t \le T. \end{cases}$

Case II: We consider now $b_1 = s_1(0) \ge s_2(0) = b_2$. Let $\delta > 0$, then $b_2 \le b_1 < b_1 + \delta$. Let w_{δ}, s_{δ} be the solution of (P1) whit latent heat ℓ_1 and $s_{\delta}(0) = b_1 + \delta$ in (Eq.9). We define the functions φ_{δ} and ψ_{δ} on the intervals $[0, b_1 + \delta]$ and $[b_1 + \delta, 1]$ respectively by:

$$\varphi_{\delta}(x) = \varphi\left(\frac{b_1 x}{b_1 + \delta}\right)$$

and

$$\psi_{\delta}(x) = \psi\left(\frac{(1-b_1)x - \delta}{1 - b_1 - \delta}\right)$$

From case I, we have that:

$$s_2(t) < s_\delta(t) \ \forall \ t \ge 0.$$

Taking into account the Stefan condition (Eq.10) for s_1 and s_{δ} , and from Case I, it results that:

$$\begin{split} & [s_{\delta}(t) - s_{1}(t)] = \delta + \frac{1}{\rho\ell_{1}} \int_{0}^{t} k_{s} [v_{\delta x}(1,\tau) - v_{1x}(1,\tau)] d\tau - \\ & \frac{1}{\rho\ell_{1}} \int_{0}^{t} k_{l} [u_{\delta x}(0,\tau) - u_{1x}(0,\tau)] d\tau \\ & - \frac{c_{l}}{\ell_{1}} \left[\int_{0}^{s_{\delta}(t)} u_{\delta}(x,t) dx - \int_{0}^{s_{1}(t)} u_{1}(x,t) dx \right] + \\ & \frac{c_{s}}{\ell_{1}} \left[\int_{s_{1}(t)}^{b_{1}+\delta} u_{\delta}(x,0) dx - \int_{0}^{b_{1}} v_{\delta}(x,t) dx \right] \\ & + \frac{c_{l}}{\ell_{1}} \left[\int_{0}^{b_{1}+\delta} u_{\delta}(x,0) dx - \int_{0}^{b_{1}} u_{1}(x,0) dx \right] + \\ & \frac{c_{s}}{\ell_{1}} \left[- \int_{b_{1}}^{1} v_{1}(x,0) dx + \int_{b_{1}+\delta}^{1} v_{\delta}(x,0) dx \right] \\ & < \delta + \frac{1}{\rho\ell_{1}} \int_{0}^{t} k_{s} \left[v_{\delta x}(1,\tau) - v_{1x}(1,\tau) \right] d\tau - \\ & \frac{1}{\rho\ell_{1}} \int_{0}^{t} k_{l} \left[u_{\delta x}(0,\tau) - u_{1x}(0,\tau) \right] d\tau \\ & + \frac{c_{l}}{\ell_{1}} \left[\int_{0}^{b_{1}+\delta} \left[\varphi_{\delta}(x) - \varphi(x) \right] dx + \int_{b_{1}}^{b_{1}+\delta} \varphi_{\delta}(x) dx \right] + \\ & \frac{c_{s}}{\ell_{1}} \left[\int_{b_{1}+\delta}^{1} \left[\psi_{\delta}(x) - \psi(x) \right] dx - \int_{b_{1}}^{b_{1}+\delta} \psi(x) dx \right]. \end{split}$$

If we apply again the maximum principle and Hopf's lemma, we can see that: $\begin{aligned} v_{\delta x}(1,\tau) - v_{1x}(1,\tau) &< 0 \text{ and } u_{\delta x}(0,\tau) - u_{1x}(0,\tau) > 0, \\ \text{then} \\ s_{\delta}(t) - s_{1}(t) &< \delta + \frac{c_{l}}{\ell_{1}} \Big[\int_{0}^{b_{1}} \Big[\varphi_{\delta}(x) - \varphi(x) \Big] dx + \\ \int_{b_{1}}^{b_{1}+\delta} \varphi_{\delta}(x) dx \Big] + \frac{c_{s}}{\ell_{1}} \Big[\int_{b_{1}+\delta}^{1} \Big[\psi_{\delta}(x) - \psi(x) \Big] dx + \\ \int_{b_{1}}^{b_{1}+\delta} \psi(x) dx \Big], \quad \forall \delta > 0, \\ \text{which implies} \\ s_{2}(t) &< s_{\delta}(t) < s_{1}(t) + \delta + \frac{c_{l}}{\ell_{1}} \int_{0}^{b_{1}} \Big[\varphi_{\delta}(x) - \varphi(x) \Big] dx + \\ \frac{c_{l}}{\ell_{1}} \int_{b_{1}}^{b_{1}+\delta} \varphi_{\delta}(x) dx - \frac{c_{s}}{\ell_{1}} \int_{b_{1}+\delta}^{1} \Big[\psi_{\delta}(x) - \psi(x) \Big] dx \\ - \frac{c_{s}}{\ell_{1}} \int_{b_{1}}^{b_{1}+\delta} \psi(x) dx, \quad \forall \delta > 0. \\ \text{Taking the limit when } \delta \to 0, \text{ we have that } s_{2} \leq s_{1} \end{aligned}$

Taking the limit when $\delta \to 0$, we have that $s_2 \leq s_1$ in the common domain of existence. To complete the proof, we consider for i = 1, 2

 $\begin{aligned} w_i(x,t) &= \begin{cases} u_i(x,t) & 0 \leq x \leq s_i(t), \quad 0 < t < T; \\ v_i(x,t) & s_i(t) \leq x \leq 1, \quad 0 < t < T. \\ As we have seen before, w_1 - w_2 \geq 0 \text{ for } 0 \leq x \leq s_i(t) \end{cases} \end{aligned}$

As we have seen before, $w_1 - w_2 \ge 0$ for $0 \le x \le s_2(t)$ and $s_1(t) \le x \le 1$. For $s_2(t) \le x \le s_1(t)$, we have $w_1(x,t) - w_2(x,t) = u_1(x,t) - v_2(x,t) \ge 0$, because $u_1 \ge 0$ and $v_2 \le 0$ and the thesis holds. b) Now, we consider that the thermal conductivity of

the liquid phase changes. Let $k_{l1} < k_{l2}$ and we note as $\{w_1, s_1\}$, $\{w_2, s_2\}$ the solutions of problem (P1) with $k_l = k_{li}$ and $s_i(0) = b_i$ for i = 1, 2. As before, we consider two cases: $s_1(0) > s_2(0)$ and $s_1(0) \ge s_2(0)$.

Case I: $b_1 = s_1(0) < s_2(0) = b_2$ and t_0 is the first time such that $s_1(t_0) = s_2(t_0)$, then we have (see Fig.3):

$$s_1'(t_0) \ge s_2'(t_0).$$
 (24)

If we define the function

,

 $\begin{cases} U(x,t) = u_2(x,t) - u_1(x,t), \ 0 \le x \le s_1(t), & t \ge 0, \\ V(x,t) = v_2(x,t) - v_1(x,t), \ s_2(t) \le x \le 1, & t \ge 0, \\ we \ see \ that \ U \ satisfies \ the \ following \ conditions: \end{cases}$

$$U_t - \frac{k_{l2}}{\rho c_l} U_{xx} \ge 0, \ 0 < x < s_1(t), \quad 0 < t < t_0, \ (25)$$

$$U(0,t) = 0, \quad 0 < t < t_0, \tag{26}$$

$$U(x,0) = 0, \quad 0 \le x \le b_1, \tag{27}$$

$$U(s_1(t), t) = u_2(s_1(t), t) - 0 > 0, \quad 0 < t < t_0, \quad (28)$$

$$s_1(0) = b_1.$$
 (29)

The inequality (Eq.25) is a consequence of part (ii) of Lemma 1, while the condition (Eq.28) is obtained from the minimum principle and then we conclude that $U \ge 0$ in $0 \le x \le s_1(t), 0 \le t \le t_0$. At $x = s_1(t_0) = s_2(t_0)$, we have $U(s_1(t_0), t_0) = 0$, then U has a minimum at $(s_1(t_0), t_0)$ and from the Hopf's lemma, it results

$$U_x(s_1(t_0), t_0) < 0. (30)$$

On the other hand, the function V satisfies the following conditions:

$$V_t - \alpha_s V_{xx} = 0, \ s_2(t) < x < 1, \quad 0 < t < t_0, \quad (31)$$



Figure 3: Free boundaries for $k_{l1} < k_{l2}$

$$V(s_2(t), t) = -v_1(s_2(t), t) > 0, \quad 0 < t < t_0, \quad (32)$$

$$V(x,0) = 0, \quad b_2 \le x \le 1, \tag{33}$$

$$V(1,t) = 0, \quad 0 < t < t_0, \tag{34}$$

$$s_2(0) = b_2. (35)$$

In the same way as above, we have that $V(x,t) \ge 0$ in $s_2(t) \le x \le 1$, $0 \le t \le t_0$. Since $V(s_2(t_0), t_0) =$ 0, V has a minimum at $(s_2(t_0), t_0)$ and from Hopf's lemma, it results

$$V_x(s_2(t_0), t_0) > 0. (36)$$

 $\begin{array}{ll} Therefore, & from & the & Stefan & condition\\ (Eq.6), (Eq.24), (Eq.30) & and & Lemma \ 1 \ (i) & we & have: \\ V_x(s_1(t_0), t_0) = v_{2x}(s_1(t_0), t_0) - v_{1x}(s_1(t_0), t_0) \\ &= \frac{\rho \ell}{k_s} [s_2'(t_0) - s_1'(t_0)] + \left[\frac{k_{l2}}{k_s} U_x(s_1(t_0), t_0) \\ &+ \frac{k_{l2} - k_{l1}}{k_s} u_{1x} \left(s_1(t_0), t_0 \right) \right] < 0, \end{array}$

which is a contradiction with (Eq.36). Then we conclude that:

$$\begin{cases} s_1(t) < s_2(t), \\ u_1(x,t) \le u_2(x,t), & 0 \le x \le s_1(t), \\ v_1(x,t) \le v_2(x,t), & s_2(t) \le x \le 1 \\ \text{for } 0 \le t \le T. \end{cases}$$

Case II: Let $b_1 = s_1(0) \leq s_2(0) = b_2$. Let $\delta > 0$, then $b_1 \leq b_2 < b_2 + \delta$. We note $\{w_{\delta}, s_{\delta}\}$ the solution of the problem (P1) with thermal conductivity of liquid phase k_{l_2} and $s_{\delta}(0) = b_2 + \delta$. We define the functions φ_{δ} and ψ_{δ} on the intervals $[0, b_2 + \delta]$ and $[b_2 + \delta, 1]$ respectively by:

 $\begin{aligned} \varphi_{\delta}(x) &= \varphi\Big(\frac{b_2 \, x}{b_2 + \delta}\Big) \ and \ \psi_{\delta}(x) = \psi\Big(\frac{(1 - b_2) \, x - \delta}{1 - b_2 - \delta}\Big). \end{aligned} \\ From \ case \ I, \ we \ have \ that: \end{aligned}$

$$s_1(t) < s_{\delta}(t) \quad \forall t \ge 0.$$

From the Stefan integral condition (Eq.10) for s_2 and s_{δ} and Case I we obtain:

$$\begin{split} & [s_{\delta}(t) - s_{2}(t)] = \delta + \frac{1}{\rho\ell} k_{s} \int_{0}^{t} \left[v_{\delta x}(1,\tau) - v_{2x}(1,\tau) \right] d\tau + \\ & \frac{1}{\rho\ell} k_{l2} \int_{0}^{t} \left[u_{2x}(0,\tau) - u_{\delta x}(0,\tau) \right] d\tau - \frac{c_{l}}{\ell} \int_{0}^{s_{2}(t)} \left[u_{\delta}(x,t) - u_{2}(x,t) \right] dx - \frac{c_{l}}{\ell} \int_{s_{2}(t)}^{s_{\delta}(t)} u_{\delta}(x,t) dx + \frac{c_{s}}{\ell} \int_{s_{\delta}(t)}^{1} \left[v_{2}(x,t) \right] dx \end{split}$$

$$-v_{\delta}(x,t) \bigg] dx + \frac{c_l}{\ell} \int_{s_2(t)}^{s_{\delta}(t)} v_2(x,t) dx + \frac{c_l}{\ell} \bigg[\int_0^{b_2+\delta} \varphi_{\delta}(x) dx \\ - \int_0^{b_2} \varphi(x) dx \bigg] + \frac{c_s}{\ell} \bigg[\int_{b_2+\delta}^1 \psi_{\delta}(x) dx - \int_{b_2}^1 \psi(x) dx \bigg].$$

If we apply the maximum principle and Hopf's lemma, we obtain that

 $\begin{bmatrix} v_{\delta x}(1,\tau) - v_{2x}(1,\tau) \end{bmatrix} < 0 \qquad an$ $\begin{bmatrix} u_{2x}(0,\tau) - u_{\delta x}(0,\tau) \end{bmatrix} < 0$

and consequently:

$$s_{\delta}(t) - s_{2}(t) < \delta + \frac{c_{l}}{\ell} \Big[\int_{0}^{b_{2}+\delta} \varphi_{\delta}(x) dx - \int_{0}^{b_{2}} \varphi(x) dx \Big] + \frac{c_{s}}{\ell} \Big[\int_{b_{2}+\delta}^{1} \psi_{\delta}(x) dx - \int_{b_{2}}^{1} \psi(x) dx \Big], \quad \forall \delta > 0.$$

When δ tends to zero, we obtain that $s_1(t) \leq s_2(t)$ in the common domain of existence.

c) By a similar argument as above, applying the maximum principle and the Hopf's lemma when it is necessary, we obtain the thesis taking into account that in this case the specific heat of each phase changes. We must consider that in this situation the heat equations (Eq.1) or (Eq.2) are affected.

d) As in c), when the mass density changes, we can show the monotony considering that the heat equations (Eq.1), (Eq.2) and also the Stefan condition (Eq.6) are affected.

III. PROBLEM WITH FLUX BOUNDARY SPECIFICATION AT BOTH FACES

A. Mathematical problem and preliminary results

We consider a similar problem to (P1), but on x = 0 and on x = 1 the conditions (Eq.3) and (Eq.4) are replaced by heat flux conditions respectively. Then, we define the problem (P2) for equations (Eq.1), (Eq.2), (Eq.5) - (Eq.9) and

$$k_l u_x(0,t) = f(t) \le 0, \ 0 < t < T,$$
 (37)

$$k_s v_x(1,t) = g(t) \le 0, \ 0 < t < T.$$
 (38)

In Tarzia (1987), Cannon (1971) and Cannon (1984) under certain hypotheses, it was proved that: *) there is unique solution for problem (P2) in $0 < t < T_0$, for some $T_0 > 0$;

**) the condition (Eq.6) is equivalent to the integral equation for all $t \ge 0$:

$$s(t) = b + \int_0^1 \Phi(x) \, dx - \frac{c_l}{\ell} \int_0^{s(t)} u(x, t) \, dx - \frac{c_s}{\ell} \int_{s(t)}^1 v(x, t) \, dx + \frac{1}{\rho \ell} \int_0^t \left[g(\tau) - f(\tau) \right] d\tau, \quad (39)$$

where $\Phi(x) = \begin{cases} \frac{c_l}{\ell} \varphi(x) & 0 \le x \le b, \\ \frac{c_s}{\ell} \psi(x) & b < x \le 1. \end{cases}$

B. Analytical results of monotony

Proposition 3 Let $\{w(x,t), s(t)\}$ be the solution of (P2). It depends monotonically on the latent heat ℓ , the mass density ρ , the specific heat c_i (i = s, l) and the thermal conductivity of the liquid phase k_l .

Proof.

a) When the latent heat ℓ changes, we consider two cases, as in the proof of Proposition 2:

Case I: $b_1 = s_1(0) > s_2(0) = b_2$ and

Case II: $b_1 = s_1(0) \ge s_2(0) = b_2$.

In both cases we repeat the proof but we consider, for the auxiliary functions U and V, the following flux data:

$$U_x(0,t) = 0, \ 0 < t < t_0,$$

$$V_x(1,t) = 0, \ 0 < t < t_0.$$

In addition, in Case II, as before, we consider the functions φ_{δ} and ψ_{δ} on the intervals $[0, b_1 + \delta]$ and $[b_1 + \delta, 1]$ respectively by:

$$\varphi_{\delta}(x) = \varphi\left(\frac{b_1 x}{b_1 + \delta}\right) \text{ and } \psi_{\delta}(x) = \psi\left(\frac{(1-b_1) x - \delta}{1 - b_1 - \delta}\right).$$

From the integral Stefan condition (Eq.39) for s_1 and s_{δ} and considering Case I we have:

$$\begin{split} &[s_{\delta}(t) - s_{1}(t)] = \\ &\delta + \frac{c_{l}}{\ell_{1}} \int_{0}^{b_{1}} [\varphi_{\delta}(x) - \varphi(x)] dx + \frac{c_{l}}{\ell_{1}} \int_{b_{1}}^{b_{1}+\delta} \varphi_{\delta}(x) dx + \\ &- \frac{c_{s}}{\ell_{1}} \int_{b_{1}}^{b_{1}+\delta} [\psi_{\delta}(x) - \psi(x)] dx \\ &- \frac{c_{s}}{\ell_{1}} \int_{b_{1}}^{b_{1}+\delta} \psi(x) dx - \frac{c_{l}}{\ell_{1}} \left[\int_{0}^{s_{1}(t)} [u_{\delta}(x,t) - u_{1}(x,t)] dx + \\ &\int_{s_{1}(t)}^{s_{\delta}(t)} u_{\delta}(x,t) dx \right] - \frac{c_{s}}{\ell_{1}} \left[\int_{s_{\delta}(t)}^{1} [v_{\delta}(x,t) - v_{1}(x,t)] dx - \\ &\int_{s_{1}(t)}^{s_{\delta}(t)} v_{1}(x,t) dx \right] \leq \delta + \frac{c_{l}}{\ell_{1}} \int_{0}^{b_{1}} [\varphi_{\delta}(x) - \varphi(x)] dx + \\ &\frac{c_{l}}{\ell_{1}} \int_{b_{1}}^{b_{1}+\delta} \varphi_{\delta}(x) dx + \frac{c_{s}}{\ell_{1}} \int_{b_{1}+\delta}^{1} [\psi_{\delta}(x) - \psi(x)] dx \\ &- \frac{c_{s}}{\ell_{1}} \int_{b_{1}}^{b_{1}+\delta} \psi(x) dx, \quad \forall \delta > 0, \\ and \ consequently: \\ s_{2}(t) < s_{\delta}(t) \leq \\ s_{1}(t) + \delta + \frac{c_{l}}{\ell_{1}} \int_{0}^{b_{1}} [\varphi_{\delta}(x) - \varphi(x)] dx + \frac{c_{l}}{\ell_{1}} \int_{b_{1}}^{b_{1}+\delta} \varphi_{\delta}(x) dx \\ &+ \frac{c_{s}}{\ell_{1}} \int_{b_{1}+\delta}^{1} [\psi_{\delta}(x) - \psi(x)] dx - \frac{c_{s}}{\ell_{1}} \int_{b_{1}}^{b_{1}+\delta} \psi(x) dx, \quad \forall \delta > 0. \end{split}$$

Taking the limit as δ tends to 0, we obtain that $s_2 \leq s_1$ in the common domain of existence. In order to complete the proof, we continue as in a) of Proposition 2.

The proof of the Proposition 3 for the other parameters is analogous to the cases b), c) and d) of Proposition 2.

IV. PROBLEM WITH TEMPERATURE SPECIFICATION ON x = 0 AND FLUX SPECIFICATION ON x = 1

A. Mathematical problem and preliminary results

We consider a similar problem to (P1), but on x = 0 and x = 1 the conditions Eq.(3) and Eq.(4) are

replaced by temperature specification and heat flux condition respectively. Then, we define the problem (P3) for e Eqs.(1), (2), (5-9) and

$$u(0,t) = f(t) > 0, \ 0 < t < T,$$
(40)

$$k_s v_x(1,t) = g(t) \le 0, \ 0 < t < T.$$
 (41)

In this case, under suitable hypothesis, it can be prove that the Stefan condition (Eq.6) is equivalent to the following integral equation for all $t \ge 0$:

$$s(t) = b + \int_0^1 \Phi(x,0) dx - \frac{c_l}{\ell} \int_0^{s(t)} u(x,t) dx$$
$$-\frac{c_s}{\ell} \int_{s(t)}^1 v(x,t) dx + \frac{1}{\rho\ell} \int_0^t \left[k_s g(\tau) - k_l u_x(0,\tau) \right] d\tau,$$
(42)
where
$$\Phi(x) = \begin{cases} \frac{c_l}{\ell} \varphi(x), & 0 \le x \le b, \\ \frac{c_s}{\ell} \psi(x) & b < x \le 1. \end{cases}$$

B. Analytical results of monotony

Proposition 4 Let $\{w(x,t), s(t)\}$ be the solution of (P3). It depends monotonically on the latent heat ℓ , the mass density ρ , the specific heat $c_i(i = s, l)$ and the thermal conductivity of the liquid phase k_l . **Proof.**

a) We consider that the thermal conductivity of the liquid phase changes and, as in the proof of item b) of Proposition 2(b), we take into account two cases:

Case I: $b_1 = s_1(0) < s_2(0) = b_2$ and

Case II: $b_1 = s_1(0) \le s_2(0) = b_2$.

We consider, for the auxiliary function V, a flux specification:

$$V_x(1,t) = 0, \ 0 < t < t_0,$$

and in Case II, as before, we consider the functions φ_{δ} and ψ_{δ} on the intervals $[0, b_2 + \delta]$ and $[b_2 + \delta, 1]$ respectively by:

$$\varphi_{\delta}(x) = \varphi\left(\frac{b_2 x}{b_2 + \delta}\right) and$$
$$\psi_{\delta}(x) = \psi\left(\frac{(1 - b_2)x - \delta}{1 - b_2 - \delta}\right).$$

From integral expression of the Stefan condition (Eq.42) for s_2 and s_{δ} and Case I we have:

$$\begin{split} \left[s_{\delta}(t) - s_{2}(t)\right] &= \delta + \frac{c_{l}}{\ell} \int_{0}^{b_{2}} \left[\varphi_{\delta}(x) - \varphi(x)\right] dx + \\ \frac{c_{l}}{\ell} \int_{b_{2}}^{b_{2}+\delta} \varphi_{\delta}(x) dx - \frac{c_{s}}{\ell} \int_{b_{2}}^{b_{2}+\delta} \psi(x) dx \\ &+ \frac{c_{s}}{\ell} \int_{b_{2}+\delta}^{1} \left[\psi_{\delta}(x) - \psi(x)\right] dx + \frac{1}{\rho \ell} k_{l2} \int_{0}^{t} \left[u_{2x}(0,\tau) - u_{\delta x}(0,\tau)\right] d\tau \\ &- \frac{c_{l}}{\ell} \int_{0}^{s_{2}(t)} \left[u_{\delta}(x,t) - u_{2}(x,t)\right] dx - \frac{c_{l}}{\ell} \int_{s_{2}(t)}^{s_{\delta}(t)} u_{\delta}(x,t) dx \\ &+ \frac{c_{s}}{\ell} \int_{0}^{1} \cdots \int_{0}^{t} \left[u_{\ell}(x,t) - u_{\ell}(x,t)\right] dx - \frac{c_{s}}{\ell} \int_{s_{2}(t)}^{s_{\delta}(t)} u_{\delta}(x,t) dx \end{split}$$

$$\begin{array}{l} + \frac{1}{\ell} \int_{s_{\delta}(t)} \left[v_{2}(x,t) - v_{\delta}(x,t) \right] dx + \frac{1}{\ell} \int_{s_{2}(t)} v_{2}(x,t) dx. \\ By \ the \ maximum \ principle \ and \ the \ Hopf's \ lemma \ \end{array}$$

$$\left[u_{2x}(0,\tau) - u_{\delta x}(0,\tau)\right] < 0$$

which implies that

we obtain

$$s_{\delta}(t) - s_{2}(t) \leq \delta + \frac{c_{l}}{\ell} \int_{0}^{b_{2}} \left[\varphi_{\delta}(x) - \varphi(x) \right] dx + \frac{c_{l}}{\ell} \int_{b_{2}}^{b_{2}+\delta} \varphi_{\delta}(x) dx - \frac{c_{s}}{\ell} \int_{b_{2}}^{b_{2}+\delta} \psi(x) dx + \frac{c_{s}}{\ell} \int_{b_{2}+\delta}^{1} \left[\psi_{\delta}(x) - \psi(x) \right] dx, \quad \forall \delta > 0,$$

and conservently, if $\delta \to 0$, we obtain $s_{1}(t) \leq s_{2}(t)$ in

and consequently, if $\delta \to 0$, we obtain $s_1(t) \leq s_2(t)$ in the common domain of existence.

The proof for the other parameters is analogous to the Proposition 2.

Remark: We can not obtain any conclusion, through the maximum principle, about variations in the thermal conductivity k_s of the solid phase. This case will be studied by a numerical approximation in Section 5.

V. NUMERICAL SOLUTIONS AND MONOTONY

A. Numerical scheme

In this section we propose a numerical method in order to approximate the solution $\{w(x,t), s(t)\}$ of the problems P(q), for q = 1, 2, 3. Since the free boundary change with time, the domain of the problem is variable. The numerical solution of free boundary problems of this type can be computed through different methods: front-traking methods and front-fixing methods (Zerroukat and Chatwin, 1994; Meyer, 1971; Crank, 1984). In this paper, we develop a scheme with a time variable mesh We propose a variable grid with fixed time step and with a constant number of space steps. These space steps will be update at each time level so that the free boundary is located on a node of the mesh (Fig. 4).

We consider a fixed time step Δt and:

*) $t_{j+1} = j\Delta t$ for j = 0, 1, ..., N. (N number of time intervals so that $t_{N+1} \leq T$, time of coexistence of both phases).

*) For each time t_{j+1} , we define a partition of [0, 1], and we take *m* steps in each phase as follow:

- in the liquid phase:

$$\Delta x_{lj} = \frac{s_j}{m}$$
, then $x_{ij} = (i-1)\Delta x_{lj}$, $i = 1, ..., m+1$
- in the solid phase:

$$\Delta x_{sj} = \frac{1-s_j}{m}, \text{ in consequence}$$

$$x_{ij} = s_j + (i-m-1)\Delta x_{sj}, i = m+2, ..., 2m+1.$$

where $s_j \simeq s(t_j)$.

*) $w_{i,j+1} \simeq w(x_{ij}, t_{j+1}), i = 1, ..., 2m + 1, j = 0, ..., N.$ *) $(x_{m+1,j}, t_{j+1}) \simeq (s_{j+1}, t_{j+1})$ for j = 0, ..., N.

We define the coefficients:

$$\begin{aligned} r_{l,j} &= \alpha_l \frac{\Delta t}{(\Delta x_{l,j})^2}; \quad \beta_{l,i,j} = \frac{\Delta t \, (i-1)s'_j}{m\Delta x_{l,j}}; \\ r_{s,j} &= \alpha_s \frac{\Delta t}{(\Delta x_{s,j})^2}; \quad \beta_{s,i,j} = \frac{\Delta t \, (2m+1-i) \, s'_j}{1-s_j} \end{aligned}$$

where, we note $s'_j \simeq s'(t_j)$, and in each phase we approach the heat equation by an explicit finite difference scheme.

In order to approximate the distribution of temperature in $(0, 1) \ge (0, T)$ we replace equations (Eq.1) and (Eq.2) with j = 0, ..., N by:

$$w_{i,j+1} = (r_{l,j} - \beta_{l,i,j}) w_{i+1,j} + (1 - 2r_{l,j} + \beta_{l,i,j}) w_{i,j} + r_{l,j} w_{i-1,j}; i = 2, ..., m; (43)$$



Figure 4: scheme for the time variable grid

$$\mathbf{w}_{i,j+1} = (r_{s,j} - \beta_{s,i,j})\mathbf{w}_{i+1,j}$$

$$+(1-2i_{s,j}+p_{s,i,j})w_{i,j}+i_{s,j}w_{i-1,j}, i=m+2,..,2m.$$
(44)

In an analogous way, we make the approximations of the boundary conditions:

for (P1): $w_{1,j+1} = f_{j+1};$

$$\mathbf{w}_{2m+1,j+1} = g_{j+1.} \tag{45}$$

for (P2): w
$$_{1,j+1} = w_{2,j+1} - \frac{f_{j+1}\Delta x_{l,j+1}}{k_l};$$

$$\mathbf{w}_{2m+1,j+1} = \mathbf{w}_{2m,j+1} + \frac{g_{j+1}\Delta x_{s,j+1}}{k_s}.$$
 (46)

for (P3): $w_{1,j+1} = f_{j+1};$

$$\mathbf{w}_{2m+1,j+1} = \mathbf{w}_{2m,j+1} + \frac{g_{j+1}\Delta x_{s,j+1}}{k_s}.$$
 (47)

We obtain the numerical solution through the following algorithm:

Step 1:

* Set the initial conditions and the boundary conditions

* Choose $\Delta t > 0$ and m;

Step 2: Set j = 1

- compute $\Delta x_{l,j}; \Delta x_{s,j},$

- evaluate s'_i from the Stefan condition (Eq.6),

- $w_{m+1,i+1} = 0$ (temperature on the free boundary), - compute $w_{i,j+1}$ i = 2, ..., m, m + 2, ..., 2m, from (Eq.43) and (Eq.44),

- compute $w_{1,j+1}$, $w_{2m+1,j+1}$ from (Eq.45), (Eq.46)or (Eq.47) depending on the problem,

Step 3: While
$$j \leq N$$
 return to Step 2 with $j = j + 1$;
Remark: Since we use an explicit scheme, we must
regard that the following four conditions are satisfied
at each step, in order to have a convergent method:
for $j = 0, ..., N$

for
$$i = 2, ..., m$$

 $r_{l,j} - \beta_{l,i,j} > 0,$ $1 - 2r_{l,j} + \beta_{l,i,j} > 0,$

for i = m + 2, ..., 2m



Figure 5: Free boundary position $(t \leq 600h)$

 $r_{s,j} - \beta_{s,i,j} > 0,$ $1 - 2r_{s,j} + \beta_{s,i,j} > 0.$ The numerical solutions obtained from the above algorithm, are used to find counterexamples that show the nonmonotonicity of the solution with respect to the solid thermal conductivity. For this reason, any analysis about stability or convergence are done.

B. Numerical results

In order to analyze the monotony, we consider the numerical solutions of problem (Pq) (q = 2, 3), for the thermal coefficients of water (Alexiades and Solomon, 1996):

- * mass density $\rho = 1000 \frac{kg}{m^3}$; * latent heat $\ell = 334000 \frac{J}{kg}$;
- * thermal conductivity of liquid phase $k_l = 0.58 \frac{W}{mK}$; * thermal conductivity of solid phase $k_s = 2.24 \frac{W}{mK}$;
- * specific heat of the liquid phase $c_l = 4185.5 \frac{J}{kqC}$;
- * specific heat of the solid phase $c_s = 2090 \frac{J}{kaC}$.

i) Problem (P2)

We consider the numerical solutions of problem (P2)for two different values of the thermal conductivity of the solid phase: $\frac{1}{20}k_s$, $\frac{1}{9}k_s$. For the initial conditions and the boundary conditions we take:

$$\varphi(x) = -76x + 38;$$
 $\psi(x) = -8x + 4$
 $f(t) = -10;$ $g(t) = -8;$

and we choose $\triangle t = 240 seq$ and m = 25.

In Figure 5, we obtain a comparative graphic of the free boundary positions and in Figure 6 we show the temperature distribution at a fixed time t = 600hs.

We can observe that the free boundary position have a monotone behavior but it does not occur for the temperature distributions.

For the numerical solutions of the problem (P3) we consider the same values of the thermal conductivity of the solid phase as above. For the initial conditions and the boundary conditions we take:

 $\varphi(x) = -140x + 70;$ $\psi(x) = -8x + 4$



Figure 6: Temperature distribution at t = 600h

ii) Problem (P3)

f(t) = 70; g(t) = -8;and we choose $\triangle t = 240 seg.$ and m = 25.

In Fig. 7, we obtain a comparative graphic of the free boundary positions, in Figure 8 we show the temperature distribution at a fixed time t = 276hs and in Figure 9 we zoom in Figure 8. We can observe that the free boundary position have a monotone behavior but it does not occur for the temperature distributions.

VI. CONCLUSIONS

It was showed, through the maximum principle, that the solutions of two-phase one-dimensional Stefan problems for finite domains, depend monotonically on the latent heat, the mass density, the specific heat of each phase and the thermal conductivity of liquid phase when we consider the following boundary conditions:

- 1) temperature specification on both fixed faces;
- 2) flux specification on both fixed faces;

3) temperature specification on the left face and flux specification on the right face.

We developed a numerical scheme using finite difference methods with variable space step in order to obtain an approximate solution for those problems. We have showed, through those approximate solutions, that there is no monotony of the solution when the thermal conductivity of the solid phase changes for the problems with boundary conditions 2) and 3).

The monotony of the solution for the problem with temperature boundary condition on both fixed faces when the thermal conductivity of the solid phase changes, is at the moment, an open problem.

REFERENCES

Alexiades, V. and A.D. Solomon, Mathematical modeling of melting and freezing processes, Hemisphere-Taylor & Francis, Washington (1996).



Figure 7: Free boundary position $(t \le 276h)$



Figure 8: Temperature distribution at t = 276 h



Figure 9: Temperature distribution at t = 276 h

- Asako, Y., E. Gonçalves, M. Faghri and M. Charmchi, "Numerical solution of melting processes for fixed and unfixed phase change material in the presence of magnetic field-simulation of low-gravity environment," Num. Heat Transf. Part A, 42, 565-583 (2002).
- Bouciguez, A.C., L.T.Villa and M.A. Lara, "Análisis de sustancias de cambio de fase para su utilización en el envasado y transporte de productos alimenticios," VI Congreso Iberoamericano de aire acondicionado y refrigeración (2001).
- Cannon, J.R. and M. Primicerio, "A two-phase Stefan problem with temperature boundary conditions," *Annali Mat. Pura Appl.*, 88, 177-191 (1971).
- Cannon, J.R., "A two-phase Stefan problem with flux boundary conditions," Annali Mat. Pura Appl., 88, 193-205 (1971).
- Cannon, J.R., *The one-dimensional heat equation*, Addison-Wesley, Menlo Park, California (1984).
- Crank, J., Free and moving boundary problems, Clarendon Press, Osford (1984).
- Farid, M.M., A.M. Khudhair, S.A.K. Razack, S. Al-Hallaj, "A review on phase change energy storage: materials and applications," *Energy Conversion* and Management, 45, 1597-1615 (2004).
- Jiji, L. and S. Gaye, "Analysis of solidification and melting of PCM with energy generation," *Appl. Thermal Eng.*, 26, 568-575 (2006).
- Lamberg, P. and K. Sirén, "Approximate analytical model for solidification in a finite PCM storage with internal fins," *Appl. Math. Modelling*, 27, 491-513 (2003).
- Lamberg, P., "Approximate analytical model for twophase solidification problem in a finned phasechange material storage," *Applied Energy*, 77, 131-152 (2004).
- Lamberg, P., R. Lehtiniemi and A. Henell, "Numerical and experimental investigation of melting and freezing processes in phase change material storage," Int. J. of Thermal Sciences, 43, 277-287 (2004).
- Medina, M., A. Bouciguez and M. Lara, "Diseño de un embalaje para productos biológicos con absorción del calor de respiración a través de un material con cambio de fase" *ERMA-Energías Renovables y Medio Ambiente*, **14**, 39-44 (2004).
- Meyer, G., "A numerical method for two-phase Stefan Problems," *SIAM J. Num. Anal.*, **8**, 555-568 (1971).

- Olguín, M., M.C. Sanziel and D.A. Tarzia, "Behavior of the solution of the Stefan problem by changing thermal coefficients of the substance," *Appl. Math. Comput.*, **190**, 765-780 (2007).
- Protter, M.H. and H.F. Weinberger, Maximum principles in differential equations, Prentice Hall, Englewood Cliffs (1967).
- Tarzia, D.A., "Estudios teóricos básicos en el Problema de Stefan unidimensional a dos fases," *Cuadern. Inst. Mat. Beppo Levi*, 14, 45-75 (1987).
- Zalba, B., J.M. Marín, L.F. Cabeza and H. Mehling, "Review on thermal energy storage with phase change materials, heat transfer analysis and applications," *Appl. Thermal Engineering*, 23, 251-283 (2003).
- Zerroukat, M. and C.R. Chatwin, Computational moving boundary problems, John Wiley & Sons, New York,(1994).
- Zivkovic, B. and I. Fujii "An Analysis of isothermal phase change of phase change material within rectangular and cylindrical containers," *Solar Energy*, **70**, 51-61 (2001).

Received: June 5, 2010 Accepted: October 6, 2010 Recommended by Subject Editor: Pedro Alcântara Pessôa