

# AN INEQUALITY FOR THE COEFFICIENT $\sigma$ OF THE FREE BOUNDARY $s(t) = 2\sigma\sqrt{t}$ OF THE NEUMANN SOLUTION FOR THE TWO-PHASE STEFAN PROBLEM\*

By

DOMINGO ALBERTO TARZIA

*Universidad Nacional de Rosario, Argentina*

**Abstract.** We consider a semi-infinite body (e.g. ice), represented by  $(0, +\infty)$ , with an initial temperature  $-c < 0$  having a heat flux  $h(t) = -h_0/\sqrt{t}$  ( $h_0 > 0$ ) in the fixed face  $x = 0$ . If  $h_0 > ck_1/\sqrt{\pi a_1}$  there exists a solution, of Neumann type, for the resulting two-phase Stefan problem. If we connect it with the Neumann problem (on  $x = 0$  the body has a temperature  $b > 0$ ) we obtain the inequality  $\text{erf}(\sigma/a_2) < (k_2 ba_1/k_1 ca_2)$  for the coefficient  $\sigma$  of the free boundary  $s(t) = 2\sigma\sqrt{t}$ , where  $k_i$  and  $a_i^2$  are respectively the thermal conductivity and thermal diffusivity coefficients of the corresponding  $i$  phase ( $i = 1$ : solid phase,  $i = 2$ : liquid phase). If  $h_0 < ck_1/\sqrt{\pi a_1}$  there is no solution of the initial problem and if  $h_0 = ck_1/\sqrt{\pi a_1}$  the problem has no physical meaning and corresponds to the case where the latent heat of fusion  $L$  tends to infinity.

## Notation.

$\Omega = (0, +\infty)$	semi-infinite body
$x$	space coordinate variable in $\Omega$
$t$	time
$s(t)$	position of the solid-liquid interface (free boundary) at time $t > 0$
$\theta(x, t)$	temperature defined for $x > 0, t > 0$
$\theta_2(x, t)$	water temperature defined for $0 < x < s(t), t > 0$
$\theta_1(x, t)$	ice temperature defined for $x > s(t), t > 0$
$c_2$	specific heat of water
$c_1$	specific heat of ice
$l$	latent heat of fusion
$\rho$	mass density
$k_2$	thermal conductivity of water
$k_1$	thermal conductivity of ice
$-c < 0$	initial temperature
$b > 0$	temperature in the fixed face $x = 0$
$h(t)$	heat flux in the fixed face $x = 0$

\* Received October 1, 1980.

$$\begin{aligned}
 C_2 &= \rho c_2, C_1 = \rho c_1, L = \rho l \\
 a_2 &= (k_2/C_2)^{1/2} & a_1 &= (k_1/C_1)^{1/2} \\
 a_2^2 && & \text{thermal diffusivity of water} \\
 a_1^2 && & \text{thermal diffusivity of ice.}
 \end{aligned}$$

**I. The problem.** We shall consider the two-phase Stefan problem for a semi-infinite body, represented by  $\Omega = (0, +\infty)$  with null change phase temperature (case: water-ice). That is, we shall find the functions  $s = s(t) > 0$  (free boundary), defined for  $t > 0$  with  $s(0) = 0$ , and

$$\begin{aligned}
 \theta(x, t) &= \theta_2(x, t) > 0 & \text{if } 0 < x < s(t) \\
 &= 0 & \text{if } x = s(t) \\
 &= \theta_1(x, t) < 0 & \text{if } s(t) < x,
 \end{aligned} \tag{1}$$

defined for  $x > 0$  and  $t > 0$ , such that they satisfy the following conditions:

$$C_1 \frac{\partial \theta_1}{\partial t} - k_1 \frac{\partial^2 \theta_1}{\partial x^2} = 0, \quad \text{in } s(t) < x, t > 0, \tag{2}$$

$$C_2 \frac{\partial \theta_2}{\partial t} - k_2 \frac{\partial^2 \theta_2}{\partial x^2} = 0, \quad \text{in } 0 < x < s(t), t > 0, \tag{3}$$

$$\theta_1(s(t), t) = 0, \quad \forall t > 0, \tag{4}$$

$$\theta_2(s(t), t) = 0, \quad \forall t > 0, \tag{5}$$

$$k_1 \frac{\partial \theta_1}{\partial x}(s(t), t) - k_2 \frac{\partial \theta_2}{\partial x}(s(t), t) = Ls'(t), \quad \forall t > 0, \tag{6}$$

$$\theta_1(x, 0) = -c < 0, \quad \forall x > 0, \tag{7}$$

$$k_2 \frac{\partial \theta_2}{\partial x}(0, t) = h(t), \quad \forall t > 0. \tag{8}$$

The function  $h(t)$  represents the heat flux that the material  $\Omega$  receives in its fixed face  $x = 0$ .

In the case

$$h(t) = -(h_0/\sqrt{t}) \quad (h_0 > 0), \tag{9}$$

we prove that there is not always a solution of Neumann type [1, 2, 3, 4, 7] for the problem (2) to (9). Moreover, the explicit solution exists if the constant  $h_0$  satisfies a certain inequality (19). This idea was suggested in [5], where simple exact solutions are given for the steady-state two-phase Stefan problem in which the heat flux satisfies an inequality on a given portion of the body's boundary and the temperature has a constant sign (for example, positive) on the remaining body's boundary.

**II. Solution of problem (2)–(9).** Following the idea of Neumann for the two-phase Stefan problem [1, 2, 3, 4, 7], we propose:

$$\begin{aligned}
 \theta_1(x, t) &= A_1 + B_1 f(x/2a_1\sqrt{t}), \\
 \theta_2(x, t) &= A_2 + B_2 f(x/2a_2\sqrt{t}), \\
 s(t) &= 2\omega\sqrt{t}, \quad \omega > 0,
 \end{aligned} \tag{10}$$

where

$$f(y) = \frac{2}{\sqrt{\pi}} \int_0^y \exp(-u^2) du, \quad a_1 = \sqrt{k_1/C_1}, \quad a_2 = \sqrt{k_2/C_2}. \quad (11)$$

The four conditions (4), (5), (7), (8) give rise to the two systems of equations

$$A_1 + B_1 = -c, \quad A_1 + f(\omega/a_1)B_1 = 0, \quad (12)$$

$$B_2 = -\sqrt{\pi} \frac{a_2}{k_2} h_0, \quad A_2 + f(\omega/a_2)B_2 = 0. \quad (13)$$

Solving (12) and (13) as functions of  $\omega$ , we obtain:

$$A_1(\omega) = c \frac{f\left(\frac{\omega}{a_1}\right)}{1 - f\left(\frac{\omega}{a_1}\right)}, \quad B_1(\omega) = \frac{-c}{1 - f\left(\frac{\omega}{a_1}\right)}, \quad (14)$$

$$A_2(\omega) = \sqrt{\pi} \frac{a_2 h_0}{k_2} f\left(\frac{\omega}{a_2}\right), \quad B_2(\omega) = -\sqrt{\pi} \frac{a_2 h_0}{k_2}.$$

The condition (6) is satisfied if  $\omega$  is a solution of the equation

$$F_0(\omega) = \omega, \quad \omega > 0 \quad (15)$$

where

$$F_0(\omega) = \frac{h_0}{L} \exp\left(-\frac{\omega^2}{a_2^2}\right) - \frac{ck_1}{L\sqrt{\pi}a_1} F_1\left(\frac{\omega}{a_1}\right), \quad F_1(x) = \frac{\exp(-x^2)}{1 - f(x)}. \quad (16)$$

Taking into account the following properties of the function  $F_1$ :

$$F_1(0^+) = 1, \quad F_1(+\infty) = +\infty, \quad (17)$$

$$\frac{dF_1}{dx}(x) > 0, \quad \forall x > 0 \quad (\text{cf. Appendix}),$$

we deduce for the function  $F_0$

$$F_0(0^+) = \frac{1}{L} \left( h_0 - \frac{ck_1}{\sqrt{\pi}a_1} \right), \quad (18)$$

$$F_0(+\infty) = -\infty, \quad \frac{dF_0}{d\omega}(\omega) < 0, \quad \forall \omega > 0,$$

and therefore we obtain

**LEMMA 1.** There exists a solution (10) of the problem (2)–(9) iff  $F_0(0^+) > 0$  and iff

$$h_0 > ck_1/\sqrt{\pi}a_1 \quad (19)$$

*Proof.* Using the properties (18), Eq. (15) has a unique solution iff  $F_0(0^+) > 0$ .

**Remark 1.** If  $h_0 < ck_1/\sqrt{\pi}a_1$ , there is no solution of the problem (2)–(9) of the type (10). The limit case  $h_0 = ck_1/\sqrt{\pi}a_1$  has no physical meaning and corresponds to the case  $L \rightarrow +\infty$  (cf. Sec. IV).

**III. Relationship with the Neumann solution.** The temperature in the fixed face  $x = 0$  is given by

$$b_0 = \theta_2(0, t) = A_2(\omega) = \sqrt{\pi} \frac{a_2 h_0}{k_2} f\left(\frac{\omega}{a_2}\right). \quad (20)$$

Since  $b_0 > 0$ , we can consider the two-phase Stefan problem consisting in finding the functions  $s(t)$ ,  $\theta_1(x, t)$ ,  $\theta_2(x, t)$  solutions of (2)–(7) and (8 bis), where:

$$\theta_2(0, t) = b \quad \text{with} \quad b > 0. \quad (8 \text{ bis})$$

The solution of problem (2)–(7) and (8 bis), which is known as Neumann solution [1, 2, 3, 4, 7], is given by:

$$\begin{aligned} \theta_1(x, t) &= \alpha_1 + \beta_1 f\left(\frac{x}{2a_1\sqrt{t}}\right), \\ \theta_2(x, t) &= \alpha_2 + \beta_2 f\left(\frac{x}{2a_2\sqrt{t}}\right), \\ s(t) &= 2\sigma\sqrt{t}, \end{aligned} \quad (21)$$

with

$$\begin{aligned} \alpha_1(\sigma) &= c \left\{ f\left(\frac{\sigma}{a_1}\right) / \left[ 1 - f\left(\frac{\sigma}{a_1}\right) \right] \right\}, \\ \beta_1(\sigma) &= -c / \left[ 1 - f\left(\frac{\sigma}{a_1}\right) \right], \\ \alpha_2(\sigma) &= b, \\ \beta_2(\sigma) &= -b / f\left(\frac{\sigma}{a_2}\right). \end{aligned} \quad (22)$$

Here  $\sigma$  is the unique solution of the equation

$$F(\sigma) = \sigma, \quad \sigma > 0 \quad (23)$$

with

$$F(\sigma) = \frac{k_1}{La_1\sqrt{\pi}} \beta_1(\sigma) \exp\left(-\frac{\sigma^2}{a_1^2}\right) - \frac{k_2}{La_2\sqrt{\pi}} \beta_2(\sigma) \exp\left(-\frac{\sigma^2}{a_2^2}\right) \quad (24)$$

which satisfies:

$$F(0^+) = +\infty, \quad F(+\infty) = -\infty, \quad \frac{dF}{d\sigma}(\sigma) < 0, \quad \forall \sigma > 0. \quad (25)$$

**Remark 2.** Notice that

$$A_1(x) = \alpha_1(x), \quad \forall x > 0, \quad B_1(x) = \beta_1(x), \quad \forall x > 0. \quad (26)$$

**Lemma 2.** If the condition (19) is valid and we take  $b = b_0 > 0$ , then

$$\sigma = \omega. \quad (27)$$

*Proof.* Since  $\beta_2(\omega) = -\sqrt{\pi}(a_2 h_0/k_2)$ , we have:

$$\begin{aligned} F(\omega) &= \frac{-ck_1}{La_1\sqrt{\pi}} \frac{\exp(-\omega^2/a_1^2)}{1-f(\omega/a_1)} + \frac{h_0}{L} \exp(-\omega^2/a_2^2) \\ &= \frac{h_0}{L} \exp(-\omega^2/a_2^2) - \frac{ck_1}{L\sqrt{\pi}a_1} F_1(\omega) = F_0(\omega) = \omega, \end{aligned} \quad (28)$$

and from the uniqueness of  $\sigma$  in Eq. (23), we deduce (27).

**Remark 3.** With the hypothesis of Lemma 2, we have:

$$A_2(x) = \alpha_2(x), \quad \forall x > 0, \quad B_2(x) = \beta_2(x), \quad \forall x > 0. \quad (29)$$

**Remark 4.** With the hypothesis of Lemma 2, we deduce the following equivalence: Problem (2)–(9)  $\Leftrightarrow$  Problem (2)–(7) and (8 bis); this implies the inequality:

$$k_2 b / \sqrt{\pi} a_2 f(\sigma/a_2) > ck_1 / \sqrt{\pi} a_1$$

or

$$f(\sigma/a_2) < k_2 a_1 b / k_1 a_2 c = \frac{b}{c} \sqrt{k_2 C_2 / k_1 C_1}. \quad (30)$$

Notice that this inequality takes into account the temperatures  $b, c$  and the coefficients  $k_1, a_1$  and  $k_2, a_2$  corresponding to the solid and liquid phase, but not the latent heat of fusion  $L$ .

**IV. Limit cases.** Using a method analogous to that in [6], we have

**LEMMA 3.**

$$F_0(0^+) = 0 \Leftrightarrow h_0 = \frac{ck_1}{a_1\sqrt{\pi}} \Leftrightarrow \omega = 0 \Leftrightarrow L = +\infty. \quad (31)$$

Let  $\theta_L(x, t)$  be the function defined by (10), (14) and (15) for each  $L > 0$ . Then:

**Remark 5.** If  $h_0 > ck_1/a_1\sqrt{\pi}$ , the limit of  $\theta_L(x, t)$  as  $L \rightarrow +\infty$  is given by:

$$\begin{aligned} \theta_{L=\infty}(x, t) &= \lim_{L \rightarrow +\infty} \theta_L(x, t) = 0 & \text{if } x = 0, \\ &= -cf \left( \frac{x}{2a_1\sqrt{t}} \right) < 0 & \text{if } x \neq 0. \end{aligned} \quad (32)$$

Moreover,  $\theta_{L=\infty}$  is a continuous function, but its heat flux in  $x = 0$  is given by:

$$k_1 \frac{\partial \theta_{L=\infty}}{\partial x}(0, t) = \frac{-ck_1}{a_1\sqrt{\pi t}} \neq \frac{-h_0}{\sqrt{t}} = h(t). \quad (33)$$

This implies that the limit  $L \rightarrow +\infty$  has no physical meaning, as was remarked in [6].

**Remark 6.** If  $h_0 > ck_1/a_1\sqrt{\pi}$  the limit of  $\theta_L(x, t)$  as  $L \rightarrow 0$  is given by:

$$\theta_{L=0}(x, t) = \lim_{L \rightarrow 0} \theta_L(x, t) = \frac{\sqrt{\pi} a_2 h_0}{k_2} \left( 1 - f\left(\frac{x}{2a_2\sqrt{t}}\right) \right). \quad (34)$$

The function  $\theta_{L=0}$  is continuous and its heat flux in  $x = 0$  is:

$$k_2(\partial\theta_{L=0}/\partial x)(0, t) = -h_0/\sqrt{t} = h(t). \quad (35)$$

**Appendix.** Let

$$\begin{aligned} F_1(x) &= \frac{\exp(-x^2)}{1-f(x)}, & f(x) &= \frac{2}{\sqrt{\pi}} \int_0^x \exp(-u^2) du = \operatorname{erf}(x), \\ H(x) &= \frac{f'(x)}{1-f(x)} = \frac{2}{\sqrt{\pi}} F_1(x), & G(x) &= H(x) - 2x. \end{aligned} \quad (1)$$

We prove the following properties:

**LEMMA.**

- (i)  $H(0) = 2/\sqrt{\pi}, \quad H(+\infty) = +\infty, \quad H(x) > 0, \quad \forall x > 0,$
- (ii)  $G(0) = 2/\sqrt{\pi}, \quad G(+\infty) = 0,$
- (iii)  $H'(x) = G(x) \cdot H(x), \quad G'(x) = H'(x) - 2,$
- (iv)  $G(x) > 0, \quad \forall x > 0,$
- (v)  $F_1'(x) > 0, \quad \forall x > 0,$

*Proof.* (i), (ii) and (iii) are evident by definition or by application of L'Hopital's rule. (iv) we suppose that there exists  $x_0 > 0/G(x_0) = 0$ . It follows that

$$H(x_0) = 2x_0, \quad H'(x_0) = 0, \quad (3)$$

$$G(x_0) = 0, \quad G'(x_0) = -2 < 0 \quad (4)$$

The conditions (4) implies that there exists  $x_1 > x_0$ ,

$$G'(x_1) = 0, \quad G(x_1) < 0. \quad (5)$$

Therefore

$$H'(x_1) = G(x_1)H(x_1) < 0.$$

Then

$$0 = G'(x_1) = H'(x_1) - 2 < -2,$$

which is a contradiction. (v) is evident using (iii) and (iv).

#### REFERENCES

- [1] M. Brillouin, *Sur quelques problèmes non résolus de physique mathématique classique : propagation de la fusion*, Annales de l'Inst. H. Poincaré, **1**, 285-308 (1930/31)
- [2] H. S. Carslaw and J. C. Jaeger, *Conduction of heat in solids*, Clarendon Press, Oxford (1959)

- [3] J. Crank, *The mathematics of diffusion*, Clarendon Press, Oxford (1956)
- [4] L. I. Rubinstein, *The Stefan problem*, Trans. Math. Monographs, 27, Amer. Math. Soc., Providence (1971)
- [5] D. A. Tarzia, *Sobre el caso estacionario del problema de Stefan a dos fases*. Mathematicae Notae, Año 28, 73–89 (1980/81)
- [6] D. A. Tarzia, *La chaleur latente de fusion tend vers l'infini n'a pas de sense physique pour le problème de Stefan*, submitted to Int. J. Heat Mass Transfer
- [7] H. Weber, *Die partiellen Differential-gleichungen der mathematischen Physik, nach Riemanns Vorlesungen*, t. II, Braunschweig (1901), 118–122