

MAT

Serie 

Conferencias, seminarios
y trabajos de Matemática

ISSN: 1515-4904

16

*A Brief Survey on
Lubrication Problems
with Nonlinear
Boundary Conditions*

Mahdi Boukrouche

Departamento
de Matemática,
Rosario,
Argentina
Marzo 2009

UNIVERSIDAD AUSTRAL

FACULTAD DE CIENCIAS EMPRESARIALES



MAT

Serie A: CONFERENCIAS, SEMINARIOS Y TRABAJOS DE MATEMATICA

ISSN: 1515-4904

Propiedad de ACES

DIRECTOR

D. A. TARZIA Departamento de Matemática – CONICET, FCE-UA,
Paraguay 1950, S2000FZF ROSARIO, ARGENTINA.
dtarzia@austral.edu.ar

COMITE EDITORIAL Y CIENTIFICO

L. A. CAFFARELLI Department of Mathematics, Univ. of Texas at Austin,
RLM 8100 Austin, TEXAS 78712, USA.
caffarel@math.utexas.edu

R. DURAN Departamento de Matemática, FCEyN, Univ. de Buenos Aires,
Ciudad Universitaria, Pab. 1, 1428 BUENOS AIRES, ARGENTINA.
rduran@dm.uba.ar

A. FASANO Dipartimento di Matematica “U. Dini”, Univ. di Firenze,
Viale Morgagni 67/A, 50134 FIRENZE, ITALIA.
fasano@math.unifi.it

J.L. MENALDI Department of Mathematics, Wayne State University,
Detroit, MI 48202, USA.
ilm@math.wayne.edu

M. PRIMICERIO Dipartimento di Matematica “U. Dini”, Univ. di Firenze,
Viale Morgagni 67/A, 50134 FIRENZE, ITALIA.
primicer@math.unifi.it

C.V. TURNER FAMAF, Univ. Nac. de Córdoba,
Ciudad Universitaria, 5000 CORDOBA, ARGENTINA.
turner@famaf.unc.edu.ar

R. WEDER Instituto de Investigaciones en Matemáticas Aplicadas y en Sistemas,
Univ. Nac. Autónoma de México (UNAM)
Apartado Postal 20-726, MEXICO, DF 010000.
weder@servidor.unam.mx

N. WOLANSKI Departamento de Matemática, FCEyN, Univ. de Buenos Aires,
Ciudad Universitaria, Pab. 1, 1428 BUENOS AIRES, ARGENTINA.
wolanski@dm.uba.ar

SECRETARIA DE REDACCION

G. GARGUICHEVICH Departamento de Matemática, FCE-UA,
Paraguay 1950, S2000FZF ROSARIO, ARGENTINA.
ggarguichevich@austral.edu.ar

MAT es una publicación del Departamento de Matemática de la Facultad de Ciencias Empresariales de la Universidad Austral (FCE-UA) cuyo objetivo es contribuir a la difusión de conocimientos y resultados matemáticos. Se compone de dos series:

- Serie A: CONFERENCIAS, SEMINARIOS Y TRABAJOS DE MATEMATICA.
- Serie B: CURSOS Y SEMINARIOS PARA EDUCACION MATEMATICA.

La Serie A contiene trabajos originales de investigación y/o recapitulación que presenten una exposición interesante y actualizada de algunos aspectos de la Matemática, además de cursos, conferencias, seminarios y congresos realizados en el Departamento de Matemática. El Director, los miembros del Comité Editorial y Científico y/o los árbitros que ellos designen serán los encargados de dictaminar sobre los merecimientos de los artículos que se publiquen. La Serie B se compone de cursos especialmente diseñados para profesores de Matemática de cada uno de los niveles de educación: Primaria, Secundaria, Terciaria y Universitaria.

Los trabajos publicados en MAT-Serie A están indexados en las bases de datos del Mathematical Reviews, MathSciNet y Zentralblatt für Mathematik. Todos los trabajos, en formato pdf, pueden ser bajados de: www.austral.edu.ar/MAT-SerieA

MAT

**SERIE A: CONFERENCIAS, SEMINARIOS Y
TRABAJOS DE MATEMATICA**

No. 16

**A BRIEF SURVEY ON LUBRICATION PROBLEMS
WITH NONLINEAR BOUNDARY CONDITIONS**

MAHDI BOUKROUCHE

Laboratory of Mathematics EA-3889
University of Saint-Etienne, PRES Lyon University
23 Rue Paul Michelon 42023 Saint-Etienne, France
E-mail: Mahdi.Boukrouche@univ-st-etienne.fr

Rosario - March 2009

Abstract. We consider some lubrication problems in a thin domain with thickness of order ε , with mixed boundary conditions and subject to slip phenomenon on a part of the boundary. We study the existence and uniqueness results for the weak solution of each problem, then we establish the asymptotic behavior of its solutions, when the depth of the thin domain tends to zero.

Résumé. Nous considérons quelques problèmes de lubrification dans un domaine mince d'épaisseur ε , avec des conditions aux limites mixtes et soumis au phénomène de glissement de fluide au parois. Nous étudions les résultats d'existence et d'unicité de la solution faible de chaque problème, puis nous établissons le comportement asymptotique des solutions quand l'épaisseur du domaine tend vers zéro.

Resumen. Se consideran algunos problemas de lubricación en un dominio delgado de espesor ε , con condiciones de contorno mixtas y sometido a un fenómeno de deslizamiento sobre una parte de la frontera. Se estudian resultados de existencia y de unicidad de la solución débil de cada problema y luego se establece el comportamiento asintótico de las soluciones cuando el espesor del dominio tiende a cero.

Keywords: Free boundary problems; Lubrication; Non-isothermal fluid; Fluid-solid conditions; slip phenomenon; Roughness phenomenon; Asymptotic approach, Reynolds equation.

Mots clés: Problèmes à frontière libre; Lubrification; Fluide non-isotherme; conditions d'interface fluide-solide; slip Phénomène de glissement de fluide; Phénomène de rugosité; Approche asymptotique; Equation de Reynolds.

Palabras claves: Problemas de frontera libre; Lubricación; Fluidos no-isotérmicos; Condiciones fluido-sólido; Fenómeno de deslizamiento; Fenómeno de rugosidad; Comportamiento asintótico; Ecuación de Reynolds.

AMS Subject Classification: 35R35; 76B03; 76D05; 76D07; 76F10; 78M35; 74K35; 76E30.

These Notes are the enlarged content of the conference given by Prof. M. Boukrouche at the Department of Mathematics of FCE-UA, Rosario, for the Congress TEM2005 on 5-7 December 2005. They contain the basic ideas of the existence and uniqueness results of some lubrication problems subject to nonlinear boundary condition and the asymptotic behavior of its solutions when one of the dimension of the fluid domain tends to zero.

The manuscript has received and accepted on December 2008.

A BRIEF SURVEY ON LUBRICATION PROBLEMS WITH NONLINEAR BOUNDARY CONDITIONS

M. Boukrouche¹

1. INTRODUCTION

This work gives a survey on some results obtained in a series of papers [6, 9, 10, 11, 14, 15, 17] in which we consider a particular cases of the general equations describing the motion of some fluid flows in bounded thin domain, with slip and mixed boundary conditions.

We comment the basic ideas on existence, uniqueness results of the solutions of the associate problems, and also its behavior when the thickness of the thin domain tends to zero. See also [12, 13, 16].

To study lubrication problems or the fluid equations one requires the knowledge of the velocities on the fluid-solid interface. This subject is often a matter of discussion as a lot of physical parameters are involved like micro-roughness of the surface or the rheological properties of the fluid.

No-slip condition, in which the fluid is assumed to have the same velocity as the surrounding solid boundary, is widely used in mathematical studies [54]. Nevertheless, this boundary condition is sometimes overlooked and it is possible to deal with the "slip condition" which allows the fluid to slip on the surface but not to go through it. The normal component of the velocity is equal to zero while the tangential one is proportional to the tangential stresses. Existence and uniqueness theorems for a weak related formulation are easily obtained (see for example) [2]. The intermediate case in which the slip condition only occurs for sufficiently a large ratio between tangential stresses and normal stresses while the no-slip condition is retained for small ratio have also been introduced [23]. This last case is nothing else than a transposition of the well known Coulomb law between two solids [24] to the fluid solid interface and so leads to a free boundary problem model.

An accurate choice of these boundary conditions is of particular interest in the lubrication area which is concerned with thin film flow behavior. In this case, the difference of velocities between the surrounding surfaces is the governing phenomena that allows the pressure in the fluid to build up and prevent the solid surfaces from being in contact which is the main objective of the lubrication. Continuous experimental studies are being conducted [46, 47] but are still difficult due to the thickness of the gap between the solid surfaces which can be as small as 50 nanometers. In such operating conditions, a no slip condition is induced by chemical bonds between the lubricant and the surrounding surfaces. Conversely, tangential stresses are so high that they tend to destroy the chemical bonds and induce a slip phenomena. Such behavior is then close to the Tresca free boundary friction model in solid mechanics [25].

This phenomenon has been related in a lot of mechanical papers for both Newtonian and non Newtonian cases [36, 37, 49, 50, 51, 55]. Although being implicitly used in numerical

¹Laboratory of Mathematics EA-3889, University of Saint-Etienne, 23 Rue Paul Michelon 42023 Saint-Etienne, PRES Lyon University, France.

procedures in lubrication problems, a Reynolds thin film equation taking account of such slip phenomena seems to have been posed for the first time in a somewhat mathematical aspect in [52]. This study is restricted to one dimensional problems and the existence of the discretized problem is proved.

The aim of this paper is not only to give existence and uniqueness for this problem but also to obtain rigorously the equation describing such phenomena in a thin film flow by way of an asymptotic analysis in which the small parameter is the width of the gap, following the same ideas as in [4], [20]. The departure point is the Stokes equation with the Tresca boundary conditions [6] and so fall into the scope of the work of [23]. Then we generalize our results to [9, 10, 11, 12, 13, 14, 15, 16, 17]. See also [29, 30, 31, 32] for similar boundary conditions.

This brief survey is organized as follows. In Section 2 we present the derivation of the fluid equations from the three conservation laws of mass, momentum and energy. In Subsection 2.1 we formulate seven problems considered. In Section 3 we give the variational formulation of each considered problems and existence and uniqueness results. In Section 4 we study the asymptotic analysis of the first case to obtain the limit problem, when the thickness of the thin domain becomes very small. In Subsection 4.1 we study the limit problem of the first case. In Subsection 4.2 we study its uniqueness. In Subsection 4.3 we study the second case. In Section 5 we study the asymptotic analysis of the case 3. In Section 6 we study the asymptotic analysis of the case 6.

I would like to thank my friend Professor Domingo Alberto Tarzia, who kindly proposed me to write these notes. I hope that this notes can provide some idea and be useful to who is interested by this subject. Also I would like to thank Guy Bayada, Professor (Insa-Lyon), Grzegorz Łukaszewicz (Warsaw University), Lionel Ciuperca (Lyon University), for fruitful collaborations on this subject.

2. ON THE FLUID EQUATIONS

We present the derivation of the problems considered from the three *conservation laws of mass, momentum and energy*. Let a bounded domain $\Omega \subset \mathbb{R}^n$ and a times interval $[0, \tau]$. Let $v : [0, \tau] \times \Omega \rightarrow \mathbb{R}^n$ such that $(t, x) \mapsto v(t, x)$ be the velocity vector of the continuous medium, $\rho : [0, \tau] \times \Omega \rightarrow \mathbb{R}$ such that $(t, x) \mapsto \rho(t, x)$ its density, and $e : [0, \tau] \times \Omega \rightarrow \mathbb{R}$ such that $(t, x) \mapsto e(t, x)$ its specific internal energy, which are the unknowns. Let given the external forces $f : [0, \tau] \times \Omega \rightarrow \mathbb{R}^n$ and a scalar function R representing the energy contribution by unity of mass and times. It is well known (see for example [41]) that the motion of continuous medium is modeled by the following three conservation laws of mass, momentum and energy respectively

$$\frac{\partial \rho}{\partial t} + v \cdot \nabla \rho + \rho \operatorname{div}(v) = 0, \quad (2.1)$$

$$\rho \left(\frac{\partial v}{\partial t} + v \cdot \nabla v \right) = \operatorname{div}(\sigma) + \rho f, \quad (2.2)$$

$$\rho \left(\frac{\partial e}{\partial t} + v \cdot \nabla e \right) = \sigma : D(v) - \operatorname{div}(q) + R, \quad (2.3)$$

where $\sigma = (\sigma_{ij})$ (for $1 \leq i, j \leq n$) is the stress tensor, $D(v)$ is the strain rate tensor, with components

$$d_{ij}(v) = \frac{1}{2} \left(\frac{\partial v_i}{\partial x_j} + \frac{\partial v_j}{\partial x_i} \right), \quad 1 \leq i, j \leq n,$$

$$\sigma : D(v) = \sum_{i,j=1}^n \sigma_{ij} d_{ij}(v), \quad \operatorname{div}(v) = \sum_{i=1}^n \frac{\partial v_i}{\partial x_i}.$$

The first term $\sigma : D(v)$ on the right hand side of (2.3) represents the energy generated by the *deformation of the continuous medium* under the action of the shear forces, the so called *dissipation* term. q is a vector function representing the energy transport, from a macroscopic continuum sense, the heat conduction phenomenon is often described by *Fourier's law*, relating the heat flux q to the temperature T

$$q = -K(T)\nabla T, \quad (2.4)$$

where K is a scalar positive function representing the thermal conductivity, see also a damped version of *Fourier's law* introducing a heat relaxation term, [35, 22]

$$\tau \frac{\partial q}{\partial t} = -(q + K\nabla T),$$

where τ is a relaxation time required to establish a steady state of heat conduction in an element suddenly exposed to heat flux.

The case where the density ρ is not constant in time leads to the compressible Euler equations [41] a physical example is a gas dynamics. Let assume that the continuous medium is an *incompressible fluid* so ρ is constant, then the local mass conservation law (2.1) becomes

$$\operatorname{div}(v) = 0. \quad (2.5)$$

The case where the stress tensor σ is non-symmetric the medium is called *micro-polar fluid* [27, 40]. We assume also in all this study that the stress tensor σ is symmetric [24, 26, 39].

$$\sigma_{ij} = \sigma_{ji} \quad \text{for} \quad 1 \leq i, j \leq n. \quad (2.6)$$

Each stress tensor σ characterizes the kind of the fluid, so for example the power law [26]

$$\sigma = -pI + k(T)\gamma^{r-1}D(v), \quad \gamma = 2\sqrt{D(v)D(v)}, \quad (2.7)$$

where k is a given positive scalar function, r is the power law index, p is the pressure, I is the $n \times n$ identity matrix, and the product $\mu = \frac{k}{2}\gamma^{r-1}$ is the viscosity of the considered fluid.

Remark that when $r = 1$ the fluid is called *non-isothermal Newtonian*. When $r \neq 1$ the fluid is called *non-isothermal non-Newtonian* and the constitutive equation

$$S = k(T)\gamma^{r-1}D(v)$$

represents *shear thinning* for $r < 1$ and *shear thickening* for $r > 1$ fluids.

With (2.7) the equation (2.2) becomes

$$\frac{\partial v}{\partial t} + v \cdot \nabla v = f + 2^{r-1} \operatorname{div} \left(k(T) (D(v)D(v))^{\frac{r-1}{2}} D(v) \right) - \nabla p. \quad (2.8)$$

As $ID(v) = \operatorname{div} v$, so from (2.5) $\sigma : D(v) = k(T)\gamma^{r-1}D(v) : D(v)$, then with the Fourier Law (2.4) the energy conservation law (2.3) becomes

$$\frac{\partial e}{\partial t} + e \cdot \nabla e = k(T)\gamma^{r-1}D(v) : D(v) + \operatorname{div} \left(K(T)\nabla T \right) + R(T).$$

Assume that the internal energy of the fluid is given by

$$\frac{\partial e}{\partial t} + e \cdot \nabla e = C_v(T) \left(\frac{\partial T}{\partial t} + v \cdot \nabla T \right),$$

where $C_v(T)$ is the specific heat with constant volume, then the energy conservation law becomes

$$C_v(T) \left(\frac{\partial T}{\partial t} + v \cdot \nabla T \right) = 2\mu(T)D(v) : D(v) + \operatorname{div} \left(K(T)\nabla T \right) + R(T). \quad (2.9)$$

with the *behavior laws* the equations (2.5), (2.8) (2.9) describe the motion of an incompressible non-isothermal non-Newtonian fluid flow.

2.1. Formulation of the problems considered. Let ω be a fixed bounded domain in \mathbb{R}^2 , for a given function $H : \omega \rightarrow \mathbb{R}^+$, we define the surface

$$x_3 = H(x) = H(x_1, x_2).$$

In the lubrication theory it is natural to assume that the fluid film, between the two surfaces ω and $x_3 = H(x)$, is very thin. So we introduce a small positive parameter ε , and a function h such that $H(x) = \varepsilon h(x)$. Then we denote the fluid domain by

$$\Omega^\varepsilon = \{(x, x_3) \in \mathbb{R}^3 : x \in \omega \text{ and } 0 < x_3 < \varepsilon h(x)\}, \quad (2.10)$$

with $\partial\Omega^\varepsilon = \bar{\omega} \cup \bar{\Gamma}_1^\varepsilon \cup \bar{\Gamma}_L^\varepsilon$ where $\Gamma_1^\varepsilon = \{(x, x_3) \in \mathbb{R}^3 : x_3 = \varepsilon h(x)\}$ and Γ_L^ε is the lateral boundary.

In all the following repeated indices means that the summation convention is used.

Case 1. [6] The motion in the fluid is described by the basic stationary Stokes system

$$\frac{\partial \sigma_{ij}^\varepsilon}{\partial x_j} + f_i^\varepsilon = 0, \quad \operatorname{div} (u^\varepsilon) = 0 \quad \text{in } \Omega^\varepsilon, \quad (2.11)$$

with

$$\sigma_{ij}^\varepsilon = p^\varepsilon \delta_{ij} + 2\nu d_{ij}(u^\varepsilon),$$

where ν is a constant viscosity, δ_{ij} is the Kronecker symbol. The velocities on the boundary, except their tangential components, are given in terms of a given function g . The upper surface being assumed to be fixed no slip condition is given so

$$u^\varepsilon = g = 0 \quad \text{on } \Gamma_1^\varepsilon. \quad (2.12)$$

The velocity is known and parallel to the ω -plane

$$u^\varepsilon = g \quad \text{with} \quad g \cdot n = 0 \quad \text{on } \Gamma_L^\varepsilon. \quad (2.13)$$

There is no flux across ω so that

$$u^\varepsilon \cdot n = g \cdot n = 0 \quad \text{on } \omega. \quad (2.14)$$

The tangential velocity on ω is *unknown* and satisfies the Tresca friction law [25] with the friction coefficient k^ε

$$\left. \begin{aligned} |\sigma_T^\varepsilon| = k^\varepsilon &\implies \exists \lambda \geq 0 \quad u_T^\varepsilon = s - \lambda \sigma_T^\varepsilon \\ |\sigma_T^\varepsilon| < k^\varepsilon &\implies u_T^\varepsilon = s \end{aligned} \right\} \quad \text{on } \omega, \quad (2.15)$$

where $n = (n_1, n_2, n_3)$ is the outward unit normal to Γ^ε , $|\cdot|$ denotes the \mathbb{R}^2 Euclidean norm, s is the velocity of the lower surface ω ; σ_n^ε and σ_T^ε are, respectively, the normal and the tangential components of the stress tensor

$$\sigma_n^\varepsilon = \sigma_{ij}^\varepsilon n_i n_j = (\sigma^\varepsilon \cdot n) \cdot n, \quad \sigma_{T_i}^\varepsilon = \sigma_{ij}^\varepsilon n_j - \sigma_n^\varepsilon n_i,$$

and u_T^ε , is the tangential velocity,

$$u_{T_i}^\varepsilon = u_i^\varepsilon - u_j^\varepsilon n_j n_i.$$

The condition (2.15) means that in each point of ω where the Euclidean norm $|\sigma_T^\varepsilon|$ reaches the upper limit k^ε , there exists an unknown scalar $\lambda \geq 0$, such that the tangential velocity of the fluid u_T^ε is braked by $\lambda \sigma_T^\varepsilon$ according to the velocity s of the lower surface ω . So in these unknown points of the lower surface ω occur the slip of the fluid according to ω .

Case 2. [9] We consider (2.11)-(2.14) and we change the Tresca boundary conditions (2.15) by the following Coulomb friction law [25] with the friction coefficient k^ε :

$$\left. \begin{aligned} |\sigma_T^\varepsilon| = k^\varepsilon |\sigma_n^\varepsilon| &\implies \exists \lambda \geq 0 \quad u_T^\varepsilon = s - \lambda \sigma_T^\varepsilon \\ |\sigma_T^\varepsilon| < k^\varepsilon |\sigma_n^\varepsilon| &\implies u_T^\varepsilon = s \end{aligned} \right\} \quad \text{on } \omega. \quad (2.16)$$

Case 3. [10] We consider (2.11), (2.13)-(2.15) and we change the boundary condition (2.12) by the following Fourier's type

$$u^\varepsilon \cdot n = 0, \quad \sigma_T^\varepsilon(u^\varepsilon) + l^\varepsilon u^\varepsilon = 0 \quad \text{on } \Gamma_1^\varepsilon, \quad (2.17)$$

where $l^\varepsilon > 0$ is a given scalar. This means that on Γ_1^ε there are only friction.

Case 4. [11] We consider the Navier-Stokes case with the Reynolds number ε^γ

$$\varepsilon^\gamma u_j \frac{\partial u_i}{\partial x_j} = f_i + \frac{\partial \sigma_{ij}}{\partial x_j} \quad \text{in } \Omega^\varepsilon, \quad \gamma \in \mathbb{R} \quad (2.18)$$

where

$$\sigma_{ij}^\varepsilon = -p^\varepsilon \delta_{ij} + 2\mu d_{ij}(u^\varepsilon), \quad (2.19)$$

with the boundary conditions (2.12), (2.14), and we change (2.13), (2.15) by

$$u^\varepsilon = \varepsilon^\beta g \quad \text{with} \quad g \cdot n = 0 \quad \text{and} \quad \beta \in \mathbb{R} \quad \text{on } \Gamma_L^\varepsilon. \quad (2.20)$$

$$\left. \begin{aligned} |\sigma_T^\varepsilon| = k^\varepsilon &\implies \exists \lambda \geq 0 \quad u_T^\varepsilon = \varepsilon^\beta s - \lambda \sigma_T^\varepsilon \\ |\sigma_T^\varepsilon| < k^\varepsilon &\implies u_T^\varepsilon = \varepsilon^\beta s \end{aligned} \right\} \quad \text{on } \omega. \quad (2.21)$$

Case 5. [14, 16] We consider the Newtonian non-isothermal case (2.11)-(2.15) with

$$\sigma_{ij}^\varepsilon = -p^\varepsilon \delta_{ij} + 2\mu^\varepsilon(T^\varepsilon)d_{ij}(u^\varepsilon),$$

$$\operatorname{div}(K^\varepsilon \nabla T^\varepsilon) + 2\mu(T^\varepsilon)|D(u^\varepsilon)|^2 + R^\varepsilon(T^\varepsilon) = 0, \quad (2.22)$$

$$T^\varepsilon = 0 \quad \text{on } \Gamma_L^\varepsilon \cup \Gamma_1^\varepsilon, \quad (2.23)$$

$$K^\varepsilon \frac{\partial T^\varepsilon}{\partial n} = \theta^\varepsilon(T^\varepsilon) \quad \text{on } \omega, \quad (2.24)$$

where θ^ε is given function on ω .

Case 6. [15] We consider the non-Newtonian non-isothermal case (2.11)-(2.15) with

$$\sigma_{ij}^\varepsilon = -p^\varepsilon \delta_{ij} + 2\mu(T^\varepsilon)|D(u^\varepsilon)|^{r-2}d_{ij}(u^\varepsilon),$$

$$\operatorname{div}(K^\varepsilon \nabla T^\varepsilon) + 2\mu(T^\varepsilon)|D(u^\varepsilon)|^r + R^\varepsilon T^\varepsilon = 0, \quad (2.25)$$

$$T^\varepsilon = 0 \quad \text{on } \Gamma_L^\varepsilon \cup \Gamma_1^\varepsilon, \quad (2.26)$$

$$K^\varepsilon \frac{\partial T^\varepsilon}{\partial n} = b^\varepsilon \quad \text{on } \omega, \quad (2.27)$$

where $r \in \mathbb{R}$ such that $1 < r < \infty$ and b^ε is given function on ω .

Case 7. We consider also in [17] the equations (2.11)-(2.15) taking the roughness phenomenon so the small parameter ε is now related to the roughness wavelength and also to the thickness of the gap between the surfaces $z_3 = 0$ and $z_3 = \lambda\varepsilon h(z, \frac{z}{\varepsilon})$, such that the domain occupied by the fluid is

$$\Omega^\varepsilon = \{(z, z_3) \in \mathbb{R}^3 : \quad z \in \omega \quad 0 < z_3 < \lambda\varepsilon h^\varepsilon(z)\}$$

where

$$h^\varepsilon(z) = h(z, \frac{z}{\varepsilon}) \quad z \in \omega, \quad (2.28)$$

and $\lambda > 0$ is a fixed constant.

3. EXISTENCES UNIQUENESS RESULTS

We assume that the function $g \in (H^{\frac{1}{2}}(\Gamma^\varepsilon))^3$ and such that

$$\int_{\Gamma^\varepsilon} g \cdot n d\sigma = 0, \quad g_3 = 0 \quad \text{on } \Gamma_L^\varepsilon, \quad g = 0 \quad \text{on } \Gamma_1^\varepsilon, \quad g \cdot n = 0 \quad \text{on } \omega. \quad (3.1)$$

So following [33] (lemma 2.2 p.24), there exists a function G^ε such that

$$G^\varepsilon \in (H^1(\Omega^\varepsilon))^3, \quad \operatorname{div}(G^\varepsilon) = 0 \quad \text{in } \Omega^\varepsilon, \quad G^\varepsilon = g \quad \text{on } \Gamma^\varepsilon. \quad (3.2)$$

Let define now the following notations

$$\begin{aligned} V^\varepsilon &= \left\{ v \in (H^1(\Omega^\varepsilon))^3 \quad : \quad v = G^\varepsilon \text{ on } \Gamma_L^\varepsilon \cup \Gamma_1^\varepsilon \quad , \quad v \cdot n = 0 \text{ on } \omega \right\}, \\ V_0^\varepsilon &= \left\{ v \in (H^1(\Omega^\varepsilon))^3 \quad : \quad v = 0 \text{ on } \Gamma_L^\varepsilon \cup \Gamma_1^\varepsilon \quad , \quad v \cdot n = 0 \text{ on } \omega \right\}, \\ V_{div}^\varepsilon &= \left\{ v \in V^\varepsilon : \operatorname{div}(v) = 0 \quad \text{in } \Omega^\varepsilon \right\}, \end{aligned}$$

$$L_0^2(\Omega^\varepsilon) = \{q \in L^2(\Omega^\varepsilon) : \int_{\Omega^\varepsilon} q \, dx dx_3 = 0\},$$

(u, v) denotes the scalar product in $L^2(\Omega^\varepsilon)$

$$a(u, \varphi) = \sum_{i,j=1}^3 2\nu \int_{\Omega^\varepsilon} D_{ij}(u) D_{ij}(\varphi) \, dx dx_3, \quad j(\varphi) = \int_{\omega} k^\varepsilon |\varphi - s| \, dx dx_3$$

So **Case 1**, leads [24] to the following variational formulation: *For G^ε as in (3.2), find $u^\varepsilon \in V_{div}^\varepsilon$ and $p^\varepsilon \in L_0^2(\Omega^\varepsilon)$, such that*

$$a(u^\varepsilon, \varphi - u^\varepsilon) - (p^\varepsilon, \operatorname{div}(\varphi)) + j(\varphi) - j(u^\varepsilon) \geq (f^\varepsilon, \varphi - u^\varepsilon) \quad \forall \varphi \in V^\varepsilon. \quad (3.3)$$

Theorem 1. *Assuming that f^ε in $(L^2(\Omega^\varepsilon))^3$, and the friction coefficient k^ε is a non negative function in $L^\infty(\omega)$, then there exists a unique u^ε and there exists a unique (up to an additive constant) p^ε such that $(u^\varepsilon, p^\varepsilon)$ in $V_{div}^\varepsilon \times L_0^2(\Omega^\varepsilon)$ is a solution to the variational inequality (3.3).*

Proof. [6] The existence and uniqueness of u^ε in V_{div}^ε satisfying the variational inequality of the second kind (3.3) is well known and follows (for example) from [18]. To get p^ε , we apply the duality results of convex optimisation ([28] theorem 4.1 and remark 4.2). recalling [6] we can rewrite (3.3) so that it is defined on the whole of $V(\Omega^\varepsilon)$ by introducing the indicator functions:

$$\psi_{V_{div}^\varepsilon} : (L^2(\Omega^\varepsilon))^3 \rightarrow \overline{\mathbb{R}} \quad \text{such that} \quad \varphi \mapsto \psi_{V_{div}^\varepsilon}(\varphi) = \begin{cases} 0 & \text{if } \varphi \in V_{div}^\varepsilon \\ +\infty & \text{if } \varphi \notin V_{div}^\varepsilon, \end{cases}$$

and

$$H : L^2(\Omega^\varepsilon) \rightarrow \overline{\mathbb{R}} \quad \text{such that} \quad q \mapsto H(q) = \begin{cases} 0 & \text{if } q = 0 \\ +\infty & \text{if } q \neq 0, \end{cases}$$

so (3.3) is equivalent to

$$\begin{aligned} a(u, \varphi - u) + j(\varphi) - j(u) + \psi_{V_{div}^\varepsilon}(\varphi) - \psi_{V_{div}^\varepsilon}(u) &\geq \\ &\geq (f^\varepsilon, \varphi - u) \quad \forall \varphi \in V_0^\varepsilon, \quad \operatorname{div}(\varphi) = 0, \end{aligned} \quad (3.4)$$

and the unique solution of (3.3) minimizes the functional:

$$\inf_{\varphi \in V_0^\varepsilon} \left\{ \frac{1}{2} a(\varphi, \varphi) - (f^\varepsilon, \varphi) + j(\varphi) + H(\operatorname{div}(\varphi)) + \psi_{V_{div}^\varepsilon}(\varphi) \right\} \quad (3.5)$$

which can be write in the following form

$$\inf_{\varphi \in V_0^\varepsilon} \{F(\varphi) + G(\Lambda(\varphi))\}, \quad \text{where}$$

$$F : V_0^\varepsilon \rightarrow \mathbb{R} \quad \text{such that} \quad \varphi \mapsto F(\varphi) = \frac{1}{2} a(\varphi, \varphi) - (f, \varphi),$$

$$\Lambda : V_0^\varepsilon \rightarrow Y = L^2(\omega) \times L^2(\Omega^\varepsilon) \times V_0^\varepsilon,$$

$$\varphi \mapsto \Lambda(\varphi) = (\Lambda_1 \varphi, \Lambda_2 \varphi, \varphi) = (\varphi|_\omega, \operatorname{div}(\varphi), \varphi),$$

$$G : Y \rightarrow \overline{\mathbb{R}} \quad \text{such that} \quad q \mapsto G(q) = j(q_1) + H(q_2) + \psi_{V_{div}^\varepsilon}(q_3).$$

Then, the dual problem (to (3.5)) is given by: *Find p^* in $Y^* = L^2(\omega) \times L^2(\Omega^\varepsilon) \times V_0^{\varepsilon*}$ solution of the problem*

$$\sup_{q^* \in Y^*} \{-F^*(\Lambda^* q^*) - G^*(-q^*)\}, \quad (3.6)$$

where

$$F^*(\Lambda^*q^*) = \sup_{\varphi \in V_0^\varepsilon} \{ \langle \Lambda_1^*q_1^*, \varphi \rangle + \langle \Lambda_2^*q_2^*, \varphi \rangle + \langle \Lambda_3^*q_3^*, \varphi \rangle - F(\varphi) \},$$

$$G^*(-q^*) : = \sup_{q \in Y} \{ \langle -q^*, q \rangle - G(q) \} = \sup_{q_1 \in L^2(\omega)} \{ \langle -q_1^*, q_1 \rangle - j(q_1) \} +$$

$$+ \sup_{q_2 \in L^2(\Omega^\varepsilon)} \{ \langle -q_2^*, q_2 \rangle - H(q_2) \} + \sup_{q_3 \in V_0^\varepsilon} \{ \langle -q_3^*, q_3 \rangle - \psi_{V_{div}^\varepsilon}(q_3) \},$$

and from the definition of H , we have for any $q = (q_1, q_2, q_3)$ in $Y = L^2(\omega) \times L^2(\Omega^\varepsilon) \times V_0^\varepsilon$

$$G^*(-q^*) \geq \{ \langle -q_1^*, q_1 \rangle - j(q_1) \} + \{ \langle -q_3^*, q_3 \rangle - \psi_{V_{div}^\varepsilon}(q_3) \}.$$

As the function G^* from $Y^* \rightarrow \mathbb{R}$, is continuous, then the hypothesis of [28] (see chap.III, Theorem 4.1), are satisfied for the dual problem (3.6), and imply the existence of p^* in Y^* satisfying

$$\{F(u^\varepsilon) + G(\Lambda(u^\varepsilon))\} + \{F^*(\Lambda^*p^*) + G^*(-p^*)\} = 0,$$

which can be written

$$\{F(u^\varepsilon) + j(\Lambda_1 u^\varepsilon) + H(\Lambda_2 u^\varepsilon) + \psi_{V_{div}^\varepsilon}(\Lambda_3 u^\varepsilon)\}$$

$$+ \{F^*(\Lambda^*p^*) + j^*(-p_1^*) + (\psi_{V_{div}^\varepsilon})^*(-p_3^*)\} = 0.$$

Let us remark from the definition of H and by choosing $q = \Lambda\varphi$ for any φ in V^ε that

$$F(u^\varepsilon) - F(\varphi) + j(\Lambda_1 u^\varepsilon) - j(\Lambda_1 \varphi) + \psi_{V_{div}^\varepsilon}(\Lambda_3 u^\varepsilon) - \psi_{V_{div}^\varepsilon}(\Lambda_3 \varphi) + \langle p_2^*, \Lambda_2 \varphi \rangle$$

$$- \langle p_2^*, \Lambda_2 u^\varepsilon \rangle \leq \{H(\Lambda_2 u^\varepsilon) - \langle p_2^*, \text{div}(u^\varepsilon) \rangle\} \leq 0,$$

which is exactly

$$a(u^\varepsilon, \varphi - u^\varepsilon) + j(\varphi) - j(u^\varepsilon) + \psi_{V_{div}^\varepsilon}(\Lambda_3 \varphi) - \psi_{V_{div}^\varepsilon}(\Lambda_3 u^\varepsilon)$$

$$- \langle p_2^*, \text{div}(\varphi - u^\varepsilon) \rangle \geq (f^\varepsilon, \varphi - u^\varepsilon) \quad \forall \varphi \in V_0^\varepsilon. \quad (3.7)$$

So taking in (3.7) φ and u^ε in V_{div}^ε , we get exactly (3.3).

Using Green's formula with $\varphi = u^\varepsilon \pm \phi$ for any ϕ in $(H_0^1(\Omega^\varepsilon))^3$, (3.7) induces

$$\nabla p_2^* = \nu \Delta u^\varepsilon + f^\varepsilon \quad a.e. \quad in \quad \Omega^\varepsilon,$$

then as u^ε is unique in V_{div}^ε , we deduce the uniqueness (up to an additive constant) of p_2^* in $L^2(\Omega^\varepsilon)$. \square

Case 2, leads to the similar variational inequality (3.3) where the functional j is now

$$j(\varphi) = \int_{\omega} k^\varepsilon |\sigma_n^\varepsilon| |\varphi - s| dx.$$

Since $j(u^\varepsilon)$ has no meaning for $u^\varepsilon \in V_{div}^\varepsilon$, we consider (cf. [24]) a regularization operator S from $H^{-\frac{1}{2}}(\omega)$ into $L_+^2(\omega)$ defined, for all $\tau \in H^{-\frac{1}{2}}(\omega)$ and $S(\tau) \in L_+^2(\omega)$, by

$$S(\tau)(x) = | \langle \tau, \psi(x-t) \rangle_{H^{-\frac{1}{2}}(\omega), H_{00}^{\frac{1}{2}}(\omega)} |, \quad \forall x \in \omega, \quad (3.8)$$

where $\psi \in D(\omega)$ is a given positive function. Here $H^{-\frac{1}{2}}(\omega)$ is the topological dual space of $H_{00}^{\frac{1}{2}} = \{\psi|_\omega : \psi \in H^1(\Omega), \psi = 0 \text{ on } \Gamma_1 \cup \Gamma_L\}$, $L_+^2(\omega)$ is the subspace of $L^2(\omega)$ of non-negative functions. So we put $S(\sigma_n^\varepsilon)$ instead of $|\sigma_n^\varepsilon|$ in the value of functional j , which give a correct meaning of $j(\varphi)$.

We have the same Theorem 1. For the proof, we apply first Tichonov's fixed point theorem to deduce the existence of u^ε , and then the existence of p^ε is obtained using the same duality results of convex optimisation. As for the solid-solid Coulomb interface law, cf., for example, [5], [43], the uniqueness is obtained for small k^ε .

In this case we state see [9] the main results concerning the existence of a weak limit (u^*, p^*) of $(u^\varepsilon, p^\varepsilon)$, the strong convergence of u^ε to u^* in a convenient space, a specific Reynolds equation in a weak form, the limit form of the Coulomb boundary conditions, and the uniqueness of (u^*, p^*) .

Case 3 [10], leads to : For G^ε as in (3.2), find $u^\varepsilon \in V_{div}^\varepsilon$ and $p^\varepsilon \in L_0^2(\Omega^\varepsilon)$, such that

$$a_1(u^\varepsilon, \varphi - u) - (p^\varepsilon, \operatorname{div} \varphi) + j(\varphi) - j(u^\varepsilon) \geq (f^\varepsilon, \varphi - u^\varepsilon) \quad \forall \varphi \in V^\varepsilon \quad (3.9)$$

where

$$a_1(u, \varphi) = a(u, \varphi) + \int_{\Gamma_1^\varepsilon} l^\varepsilon u \varphi ds$$

the integral on Γ_1^ε comes from the Fourier condition (2.17). The bilinear form a_1 is continuous symmetric and coercive indeed following [53] suppose that a_1 is not coercive so there exists a subsequence $(w_n) \in V_{div}^\varepsilon$ such that $a_1(w_n, w_n) < \frac{1}{n} \|w_n\|_{H^1(\Omega^\varepsilon)}^2 \quad \forall n \in \mathbb{N}^*$.

Let $u_n = \frac{w_n}{\|w_n\|_{H^1(\Omega^\varepsilon)}}$, then $\|u_n\|_{H^1(\Omega^\varepsilon)} = 1$ and $a_1(u_n, u_n) < \frac{1}{n}$ for all $n \in \mathbb{N}^*$. As

$$\|u_n\|_{H^1(\Omega^\varepsilon)}^2 = a(u_n, u_n) + \|u_n\|_{L^2(\Omega^\varepsilon)}^2 \leq a_1(u_n, u_n) + \|u_n\|_{L^2(\Omega^\varepsilon)}^2 \leq \frac{1}{n} + 1 \leq 2$$

so there exists $u \in H^1(\Omega^\varepsilon)$ such that $u_n \rightharpoonup u$ in $H^1(\Omega^\varepsilon)$ weak and then in $L^2(\Omega^\varepsilon)$ strong thus $\|u\|_{L^2(\Omega^\varepsilon)} = 1$. And by

$$0 \leq \int_{\Gamma_1^\varepsilon} l^\varepsilon u_n^2 ds \leq \liminf_{n \rightarrow +\infty} \int_{\Gamma_1^\varepsilon} l^\varepsilon u_n^2 ds \leq \liminf_{n \rightarrow +\infty} \frac{1}{n} = 0$$

$$0 \leq a(u, u) \leq \liminf_{n \rightarrow +\infty} a(u_n, u_n) \leq \liminf_{n \rightarrow +\infty} \frac{1}{n} = 0$$

we obtain that $u = 0$ in Ω^ε which is impossible with $\|u\|_{L^2(\Omega^\varepsilon)} = 1$. Thus a_1 is coercive so here also the theorem 1 remains valid for this problem 2.

Case 4, leads to : Find $u^\varepsilon \in V_{div}^\varepsilon$, $p^\varepsilon \in L_0^2(\Omega^\varepsilon)$, such that

$$a(u^\varepsilon, \phi - u^\varepsilon) + \varepsilon^\gamma b(u^\varepsilon, u^\varepsilon, \phi - u^\varepsilon) - (p^\varepsilon, \operatorname{div}(\phi)) + j(\phi) - j(u^\varepsilon) \geq (f^\varepsilon, \phi - u^\varepsilon) \quad \forall \phi \in V^\varepsilon,$$

where

$$b : V^\varepsilon \times V^\varepsilon \times V^\varepsilon \rightarrow \mathbb{R} \quad : \quad (u, v, w) \rightarrow b(u, v, w) = \int_{\Omega^\varepsilon} u_i v_{j,i} w_j dx dx_3,$$

$$j(v) = \int_{\omega} k^\varepsilon |v - \varepsilon^\beta s| ds.$$

Theorem 2. *There exists μ_0 such that for $\mu > \mu_0$, this problem has at least one solution $(u^\varepsilon, p^\varepsilon)$, under the condition $\beta \geq \frac{1}{2} - \gamma$. There exists $\varepsilon^1 > 0$ such that for $\varepsilon \leq \varepsilon^1$, then u^ε , such that $(u^\varepsilon, p^\varepsilon)$ is solution of this problem, is unique.*

Proof. The condition $\beta \geq \frac{1}{2} - \gamma$ allow us to obtain the existence of a constant $C > 0$ such that the following application will be well defined

$$\Lambda : B_C \rightarrow B_C : \quad \text{such that} \quad \xi \rightarrow u^\varepsilon$$

where B_C is the $(H^1(\Omega^\varepsilon))^3$ closed ball of radius C , u^ε is the unique solution of the following variational inequality:

$$a(u^\varepsilon, \phi - u^\varepsilon) + \varepsilon^\gamma b(\xi, u^\varepsilon, \phi - u^\varepsilon) + j(\phi) - j(u^\varepsilon) \geq (f^\varepsilon, \phi - u^\varepsilon) \quad \forall \phi \in V_{div}^\varepsilon. \quad (3.10)$$

then by Schauder fixed point theorem, there exists at least one solution u^ε for the following variational inequality:

$$a(u^\varepsilon, \phi - u^\varepsilon) + \varepsilon^\gamma b(u^\varepsilon, u^\varepsilon, \phi - u^\varepsilon) + j(\phi) - j(u^\varepsilon) \geq (f^\varepsilon, \phi - u^\varepsilon) \quad \forall \phi \in V_{div}^\varepsilon. \quad (3.11)$$

the existence of the pressure p^ε comes as in Theorem 1 (see also [6]). The uniqueness follows from some estimates on the gradient of the velocity and it is valid under the condition that the fluid domain must be thin enough. For all the proof see [11]. \square

Case 5, leads to the non-isothermal coupled problem : Find

$$u^\varepsilon \in V_{div}^\varepsilon \cap (H^2(\Omega^\varepsilon))^3, \quad p^\varepsilon \in L_0^2(\Omega^\varepsilon) \cap H^1(\Omega^\varepsilon), \quad T^\varepsilon \in H_{\Gamma_L^\varepsilon \cup \Gamma_1^\varepsilon}^1(\Omega^\varepsilon) \cap C^{0,1}(\overline{\Omega^\varepsilon}),$$

such that

$$a(T^\varepsilon; u^\varepsilon, \varphi - u^\varepsilon) - (p^\varepsilon, \text{div}(\varphi)) + j(\varphi) - j(u^\varepsilon) \geq (f^\varepsilon, \varphi - u^\varepsilon), \quad \forall \varphi \in V^\varepsilon, \quad (3.12)$$

$$\int_{\Omega^\varepsilon} K^\varepsilon \nabla T^\varepsilon \nabla \psi + R^\varepsilon(T^\varepsilon) \psi \, dx, = \int_{\Omega^\varepsilon} 2\mu^\varepsilon(T^\varepsilon) |D(u)|^2 \psi \, dx + \int_{\omega} \theta^\varepsilon(T^\varepsilon) \psi \, ds, \quad \forall \psi \in H_{\Gamma_L^\varepsilon \cup \Gamma_1^\varepsilon}^1(\Omega^\varepsilon), \quad (3.13)$$

where

$$H_{\Gamma_L^\varepsilon \cup \Gamma_1^\varepsilon}^1(\Omega^\varepsilon) = \left\{ \chi \in H^1(\Omega^\varepsilon) \quad : \quad \chi = 0 \text{ on } \Gamma_L^\varepsilon \cup \Gamma_1^\varepsilon \right\}.$$

$$a(T; u, v) = \int_{\Omega^\varepsilon} 2\mu^\varepsilon(T) D(u) : D(v) \, dx' dx_3,$$

Note that the first term of $c(u; T, \psi)$ is well defined for $u \in V^\varepsilon \cap (H^2(\Omega^\varepsilon))^3$.

We study first the two following intermediate problems:

Given $T \in H_{\Gamma_L^\varepsilon \cup \Gamma_1^\varepsilon}^1(\Omega^\varepsilon) \cap C^{0,1}(\overline{\Omega^\varepsilon})$, find $v^\varepsilon \in V_{div}^\varepsilon \cap (H^2(\Omega^\varepsilon))^3$ such that

$$a(T; v^\varepsilon, \varphi - v^\varepsilon) - (p^\varepsilon, \text{div}(\varphi)) + j(\varphi) - j(v^\varepsilon) \geq (f^\varepsilon, \varphi - v^\varepsilon), \quad \forall \varphi \in V^\varepsilon.$$

Given $u \in V_{div}^\varepsilon \cap (H^2(\Omega^\varepsilon))^3$, find T^ε in $H_{\Gamma_L^\varepsilon \cup \Gamma_1^\varepsilon}^1(\Omega^\varepsilon) \cap C^{0,1}(\overline{\Omega^\varepsilon})$ such that (3.13) hold.

The main result in this case is to establish the needed regularity results. Note that the boundary $\partial\Omega^\varepsilon$ is decomposed of three connected compact components ω , Γ_1^ε , and Γ_L^ε , the angles at the corner at the intersections $\Gamma_1^\varepsilon \cap \Gamma_L^\varepsilon$ and $\omega \cap \Gamma_L^\varepsilon$ are less or equal to 90 degree. So we use the local regularity theory in a neighborhood of the boundary [44] and the partition of unity.

We obtain the regularity results in interior and near Γ_1^ε , Γ_L^ε and ω , following [21, 38, 48, 8, 34, 42], the difference here is that the coefficient of our bilinear form $a(\cdot, \cdot)$ depends on T^ε . So we obtain the following estimate

$$\|v^\varepsilon\|_{2,\Omega^\varepsilon} + \|p^\varepsilon\|_{1,\Omega^\varepsilon} \leq C (\|f^\varepsilon\|_{0,\Omega^\varepsilon} + \|k^\varepsilon\|_{1/2,\omega} + \|G^\varepsilon\|_{2,\Omega^\varepsilon}) \quad (3.14)$$

but the constant C depend on some data μ_* , μ^* , C_{μ^ε} , C_K , Ω^ε and also on $\|T^\varepsilon\|_{C^{0,1}(\overline{\Omega^\varepsilon})}$. Thus we establish the needed regularity result of the temperature T^ε , then we deduce with (3.14) the needed regularity of the velocity v^ε . Then with the Banach fixed point theorem we establish the existence and uniqueness results of the weak solution to the above coupled problem [14].

Case 6, we first assume that the function $g \in (W^{1-\frac{1}{r},r}(\Gamma^\varepsilon))^3$. So with (3.1) it is well known [3] (Lemma 3.3) that there exists a function G^ε such that

$$G^\varepsilon \in (W^{1,r}(\Omega^\varepsilon))^3 \text{ with } \operatorname{div}(G^\varepsilon) = 0 \text{ in } \Omega^\varepsilon, \text{ and } G^\varepsilon = g \text{ on } \Gamma^\varepsilon. \quad (3.15)$$

So we consider the following functional framework on Ω^ε :

$$\begin{aligned} W_{\Gamma_1^\varepsilon \cup \Gamma_L^\varepsilon}^{1,r}(\Omega^\varepsilon) &= \{\psi \in W^{1,r}(\Omega^\varepsilon) \quad : \quad \psi = 0 \text{ on } \Gamma_1^\varepsilon \cap \Gamma_L^\varepsilon\} \\ V^\varepsilon &= \{v \in (W^{1,r}(\Omega^\varepsilon))^3 \quad : \quad v = G^\varepsilon \text{ on } \Gamma_1, \quad v \cdot n = 0 \text{ on } \omega\} \\ L_0^{r'}(\Omega^\varepsilon) &= \{\varphi \in L^{r'}(\Omega^\varepsilon) \quad : \quad \int_{\Omega^\varepsilon} \varphi(x) dx = 0\} \end{aligned}$$

then we denote by r' , q' the conjugates of r and q respectively. This non-Newtonian non-isothermal leads to the following variational formulation : Find $u^\varepsilon \in V_{div}^\varepsilon$, $p^\varepsilon \in L_0^{r'}(\Omega^\varepsilon)$, and $T^\varepsilon \in W_{\Gamma_L^\varepsilon \cup \Gamma_1^\varepsilon}^{1,q}(\Omega^\varepsilon)$ such that

$$\begin{aligned} a(T^\varepsilon; u^\varepsilon, \varphi - u^\varepsilon) + (p^\varepsilon, \operatorname{div}(\varphi)) + j(\varphi) - j(u^\varepsilon) &\geq (f^\varepsilon, \varphi - u^\varepsilon), \quad \forall \varphi \in V^\varepsilon, \quad (3.16) \\ \int_{\Omega^\varepsilon} K^\varepsilon \nabla T^\varepsilon \nabla \psi + R^\varepsilon T^\varepsilon dx &= 2 \int_{\Omega^\varepsilon} \mu^\varepsilon(T^\varepsilon) |D(u^\varepsilon)|^r \psi dx + \int_{\omega} b^\varepsilon \psi ds, \quad \forall \psi \in W_{\Gamma_L^\varepsilon \cup \Gamma_1^\varepsilon}^{1,q'}(\Omega^\varepsilon), \end{aligned} \quad (3.17)$$

where

$$a(T^\varepsilon; u^\varepsilon, \varphi) = 2 \int_{\Omega^\varepsilon} \mu^\varepsilon(T^\varepsilon) |D(u^\varepsilon)|^{r-2} D(u^\varepsilon) : D(\varphi) dx$$

This non-Newtonian case is a direct generalized of the Newtonian **Case 5**. Here by the Sobolev inequalities [1], for $q' = q/(q-1) > 3$ the injection of $W_{\Gamma_L^\varepsilon \cup \Gamma_1^\varepsilon}^{1,q'}(\Omega^\varepsilon)$ in $L^\infty(\Omega^\varepsilon)$ make sense to the first term of the right-hand side of (3.13). This idea is not possible for the Newtonian **Case 5**.

We assume that there exist μ_* , μ^* , K_\star^ε , K_ε^\star , R_\star^ε , R_ε^\star , C_b^ε in \mathbb{R} such that

$$\left. \begin{aligned} \mu^\varepsilon &\in C^1(\mathbb{R}), \quad 0 < \mu_* \leq \mu^\varepsilon \leq \mu^* ; \\ f^\varepsilon &\in (W^{1,r'}(\Omega^\varepsilon))^3 ; \quad 0 < K_\star^\varepsilon \leq K^\varepsilon \leq K_\varepsilon^\star ; \\ 0 < R_\star^\varepsilon &\leq R^\varepsilon \leq r_\varepsilon^\star ; \quad |b^\varepsilon| \leq C_b^\varepsilon. \end{aligned} \right\} (H)$$

we have the following result

Lemma 1. [15] *Let $\theta \in W_{\Gamma_1^\varepsilon \cup \Gamma_L^\varepsilon}^{1,q}$. We denote by $u_\theta^\varepsilon \in V_{div}^\varepsilon$ the solution of the following inequality*

$$a(\theta; u_\theta^\varepsilon, \phi - u_\theta^\varepsilon) + j^\varepsilon(\phi) - j^\varepsilon(u_\theta^\varepsilon) \geq (f^\varepsilon, \phi - u_\theta^\varepsilon) \quad \forall \phi \in V_{div}^\varepsilon.$$

Then there exists C^ε constante independent of θ such that

$$\|\nabla u_\theta^\varepsilon\|_{L^r(\Omega^\varepsilon)} \leq C^\varepsilon. \quad (3.18)$$

And the application :

$$\theta \longrightarrow u_\theta^\varepsilon \in V_{div}^\varepsilon$$

is strongly continuous.

Theorem 3. Assume (H) hold. For all $r > 1$ the problem (3.16)-(3.17) has at last one solution $(u^\varepsilon, p^\varepsilon, T^\varepsilon)$ in $V_{div}^\varepsilon \times L_0^{r'}(\Omega^\varepsilon) \times W_{\Gamma_1 \cup \Gamma_L}^{1,q}(\Omega^\varepsilon)$.

Proof. [15] Let $\delta > 0$, we consider the function

$$(\theta, u_\theta^\varepsilon) \mapsto m_\delta(\theta) = \frac{2\mu^\varepsilon(\theta) |D(u_\theta^\varepsilon)|^r}{1 + 2\delta\mu^\varepsilon(\theta) |D(u_\theta^\varepsilon)|^r} \quad (3.19)$$

where u_θ^ε satisfies the variational inequality

$$a(\theta; u_\theta^\varepsilon, \phi - u_\theta^\varepsilon) + j^\varepsilon(\phi) - j^\varepsilon(u_\theta^\varepsilon) \geq (f^\varepsilon, \phi - u_\theta^\varepsilon) \quad \forall \phi \in V_{div}^\varepsilon.$$

Using (3.18), $\exists C_1^\varepsilon$ a constant independent of $\delta, \theta, u_\theta^\varepsilon$ such that

$$\|m_\delta\|_{L^1(\Omega^\varepsilon)} \leq C_1^\varepsilon. \quad (3.20)$$

Now, we consider the following problem : Find T_δ^ε such that

$$\int_{\Omega^\varepsilon} K^\varepsilon \nabla T_\delta^\varepsilon \nabla \psi + R^\varepsilon T_\delta^\varepsilon \psi = \int_{\Omega^\varepsilon} m_\delta \psi, + \int_{\omega} b^\varepsilon \psi \quad \forall \psi \in H_{\Gamma_1 \cup \Gamma_L}(\Omega^\varepsilon). \quad (3.21)$$

This problem has a unique solution by Lax-Milgram Lemma. Let define the application

$$\begin{aligned} \gamma : B(0, \tilde{C}) \cap W_{\Gamma_1 \cup \Gamma_L}^{1,q}(\Omega^\varepsilon) &\rightarrow B(0, \tilde{C}) \cap W_{\Gamma_1 \cup \Gamma_L}^{1,q}(\Omega^\varepsilon) \\ \theta &\mapsto T_\delta^\varepsilon \end{aligned}$$

where $B(0, \tilde{C})$ is a closed ball in $W_{\Gamma_1 \cup \Gamma_L}^{1,q}(\Omega^\varepsilon)$. We must look for $\tilde{C} > 0$ such that γ be well defined. So we choose $\psi = \varphi(T_\delta^\varepsilon)$ in (3.21), where φ is defined by

$$\varphi(t) = \xi \operatorname{sign}(t) \int_0^{|t|} \frac{d\tau}{(1+\tau)^{\xi+1}} = \operatorname{sign}(t) \left[1 - \frac{1}{(1+|t|)^\xi} \right], \quad (3.22)$$

with $\xi > 0$. we deduce from (3.20)-(3.21) after some calculations that

$$\int_{\Omega^\varepsilon} |\nabla T_\delta^\varepsilon|^q \leq 2^{\frac{(q^*-1)(2-q)}{2}} \left(\frac{C_1^\varepsilon}{\xi K_\star^\varepsilon} \right)^{\frac{q}{2}} \times \left(|\Omega^\varepsilon|^{\frac{(2-q)}{2}} + A \right). \quad (3.23)$$

A is independent of δ . So we can choose

$$\tilde{C} = \left[2^{(q^*-1)(2-q)/2} \left(\frac{C_1^\varepsilon}{\xi K_\star^\varepsilon} \right)^{q/2} \left(|\Omega^\varepsilon|^{(2-q)/2} + A \right) \right]^{\frac{1}{q}}$$

Using now lemma 1 and Schauder's fixed point theorem with the application γ , so there exists

$$(u_\delta^\varepsilon, p_\delta^\varepsilon, T_\delta^\varepsilon) \in V_{div}^\varepsilon \times L_0^{r'}(\Omega^\varepsilon) \times H^1(\Omega^\varepsilon) \cap B(0, \tilde{C})$$

where $(u_\delta^\varepsilon, p_\delta^\varepsilon)$ solves (3.12), with

$$u_\delta^\varepsilon = u_{T_\delta^\varepsilon}^\varepsilon, \quad p_\delta^\varepsilon = p_{T_\delta^\varepsilon}^\varepsilon, \quad T_\delta^\varepsilon \text{ solves (3.21)}$$

and $m_\delta = m_\delta(u_\delta^\varepsilon, T_\delta^\varepsilon)$ defined by (3.19). We obtain also the following estimation for the pressure p_δ^ε (as in [13])

$$\left\| \frac{\partial p_\delta^\varepsilon}{\partial x_i} \right\|_{W^{-1,r'}(\Omega^\varepsilon)} \leq C \quad i = 1, 2, 3, \quad (3.24)$$

using (3.18), (3.23), (3.24), and taking a subsequence $\delta \rightarrow 0$, we obtain

$$\begin{aligned} u_\delta^\varepsilon &\rightharpoonup u^\varepsilon \quad \text{weakly in } W_{\Gamma_1^\varepsilon \cup \Gamma_L^\varepsilon}^{1,r}, \\ p_\delta^\varepsilon &\rightharpoonup p^\varepsilon \quad \text{weakly in } L^{r'}(\Omega^\varepsilon), \\ T_\delta^\varepsilon &\rightharpoonup T^\varepsilon \quad \text{weakly in } W^{1,q}(\Omega^\varepsilon) \quad \text{and strongly in } L^q(\Omega^\varepsilon). \end{aligned}$$

From Lemma 1, $u_\delta^\varepsilon \rightarrow u^\varepsilon$ strongly in V_{div}^ε consequently as $\mu^\varepsilon \in C^1(\mathbb{R})$ we get

$$m_\delta = m_\delta(T_\delta^\varepsilon, u_\delta^\varepsilon) \rightarrow 2\mu^\varepsilon(T^\varepsilon) |D(u^\varepsilon)|^r \quad \text{in } L^1(\Omega^\varepsilon),$$

we conclude that the limit, $(u^\varepsilon, p^\varepsilon, T^\varepsilon)$ solves Problem (3.16)-(3.17). \square

Case 7, leads to the same variational inequality (3.3) so the existence and uniqueness of the weak solution comes from Theorem 1. The main difficulties is to study the behavior of the weak solution and especially how to pass to the two-scale limit in the variational inequality, due to the term coming from the Tresca fluid-solid boundary conditions. This difficulty induce us to prove (see [17]) a needed result of lower-semicontinuity for the two-scale convergence, using some results on subdifferential and regularization of convex functions.

4. ASYMPTOTIC ANALYSIS

To be able to compare the solutions for various ε and provide the asymptotic analysis, we use the change of variables $y = x_3/\varepsilon$ to define the fixed domain

$$\Omega = \{(x, y) \text{ such that } x \in \omega, \text{ and } 0 < y < h(x)\},$$

and we denote by $\Gamma = \bar{\omega} \cup \bar{\Gamma}_L \cup \bar{\Gamma}_1$ its boundary. For the three cases 1-3, we define the following functions in Ω

$$\begin{aligned} \hat{u}_i^\varepsilon(x, y) &= u_i^\varepsilon(x, x_3) \quad 1 \leq i \leq 2, \quad \hat{u}_3^\varepsilon(x, y) = \frac{1}{\varepsilon} u_3^\varepsilon(x, x_3), \\ \hat{p}^\varepsilon(x, y) &= \varepsilon^2 p^\varepsilon(x, x_3). \end{aligned}$$

Let us define first the ε -independent vector

$$\hat{f}(x, y) = (\hat{f}_1(x, y), \hat{f}_2(x, y), \hat{f}_3(x, y)),$$

then assume the following dependence (with respect to ε) of the data

$$\hat{f}(x, y) = \varepsilon^2 f^\varepsilon(x, x_3), \quad \hat{g}(x, y) = g(x, x_3). \quad (4.1)$$

$$\hat{k} = \varepsilon k^\varepsilon \text{ for the Tresca cases 1, 3, but } \hat{k} = \varepsilon^{-1} k^\varepsilon \text{ for the Coulomb case 2.} \quad (4.2)$$

The first assumption in (4.1) means that the body forces cannot be too big. In (4.2) the first one means that k^ε , the upper limit for the tangential stress has the same order of magnitude as the actual stress inside the fluid, which is the ratio of the tangential velocity and of the gap: $\frac{S}{\varepsilon h}$, while the second one means that, roughly speaking, the friction coefficient $k^\varepsilon \sim \varepsilon$ is the ratio of the tangential stress inside the fluid σ_T^ε and of the normal stress σ_n^ε , while $\sigma_T^\varepsilon \sim \frac{1}{\varepsilon}$ (ratio between the tangential velocity and the thickness of the gap) and $\sigma_n^\varepsilon \sim \frac{1}{\varepsilon^2}$ (order of magnitude of the actual pressure).

Let us define the ε -independent vector $\hat{G}(x, y) = (\hat{G}_1(x, y), \hat{G}_2(x, y), \hat{G}_3(x, y))$ such that

$$\frac{\partial \hat{G}_1}{\partial x_1} + \frac{\partial \hat{G}_2}{\partial x_2} + \frac{\partial \hat{G}_3}{\partial y} = 0 \quad \text{in } \Omega, \quad \hat{G} = \hat{g} \quad \text{on } \Gamma,$$

and recalling that $g_3 = 0$ on Γ_L , then we can choose as G^ε the lift defined by

$$G_i^\varepsilon(x, x_3) = \hat{G}_i(x, y) \quad \text{for } i = 1, 2 \quad \text{and} \quad G_3^\varepsilon(x, x_3) = \varepsilon \hat{G}_3(x, y).$$

Now we define function spaces and sets on Ω we need in our considerations.

$$V = \{\varphi \in (H^1(\Omega))^3 : \varphi \cdot n = 0 \quad \text{on } \omega, \quad v = \hat{G} \quad \text{on } \Gamma_L \cup \Gamma_1\},$$

$$V_{div} = \{\varphi \in V : \quad \text{div } \varphi = 0 \text{ in } \Omega\}$$

$$V_0(\Omega) = \{v \in (H^1(\Omega))^3 : v \cdot n = 0 \quad \text{on } \omega, \quad v = 0 \text{ on } \Gamma_L \cup \Gamma_1\},$$

$$L_0^2(\Omega) = \{q \in L^2(\Omega) : \int_{\Omega} q dx dy = 0\},$$

Then assuming (4.1) and the first of (4.2), there exists a unique \hat{u}^ε in V_{div} and \hat{p}^ε in $L_0^2(\Omega)$, such that the variational inequality (3.3) leads to the following form:

$$\begin{aligned} & \sum_{i,j=1}^2 \int_{\Omega} \left(\varepsilon^2 \nu \left(\frac{\partial \hat{u}_i^\varepsilon}{\partial x_j} + \frac{\partial \hat{u}_j^\varepsilon}{\partial x_i} \right) - \hat{p}^\varepsilon \delta_{i,j} \right) \frac{\partial}{\partial x_j} (\varphi_i - \hat{u}_i^\varepsilon) dx dy + \\ & \sum_{i=1}^2 \int_{\Omega} \nu \left(\frac{\partial \hat{u}_i^\varepsilon}{\partial y} + \varepsilon^2 \frac{\partial \hat{u}_3^\varepsilon}{\partial x_i} \right) \frac{\partial}{\partial y} (\varphi_i - \hat{u}_i^\varepsilon) dx dy + \int_{\Omega} (2\nu \varepsilon^2 \frac{\partial \hat{u}_3^\varepsilon}{\partial y} - \hat{p}^\varepsilon) \frac{\partial}{\partial y} (\varepsilon^{-1} \varphi_3 - \hat{u}_3^\varepsilon) dx dy \\ & + \sum_{j=1}^2 \int_{\Omega} \varepsilon^2 \nu \left(\varepsilon^2 \frac{\partial \hat{u}_3^\varepsilon}{\partial x_j} + \frac{\partial \hat{u}_j^\varepsilon}{\partial y} \right) \frac{\partial}{\partial x_j} (\varepsilon^{-1} \varphi_3 - \hat{u}_3^\varepsilon) dx dy \geq \sum_{i=1}^2 \int_{\Omega} \hat{f}_i (\varphi_i - \hat{u}_i^\varepsilon) dx dy \\ & + \int_{\Omega} \varepsilon \hat{f}_3 (\varepsilon^{-1} \varphi_3 - \hat{u}_3^\varepsilon) dx dy + \int_{\omega} \hat{k} (|\varphi - s|) - |\hat{u}^\varepsilon - s| dx \quad \forall \varphi \in K. \end{aligned} \quad (4.3)$$

Theorem 4. *Assuming (4.1) and the first of (4.2) we have the following estimate on \hat{u}^ε*

$$\begin{aligned} & \frac{\nu \varepsilon^2}{2} \left\| \frac{\partial \hat{u}_1^\varepsilon}{\partial x_1} \right\|^2 + \frac{\nu \varepsilon^2}{2} \left\| \frac{\partial \hat{u}_1^\varepsilon}{\partial x_2} \right\|^2 + \frac{\nu \varepsilon^2}{2} \left\| \frac{\partial \hat{u}_2^\varepsilon}{\partial x_1} \right\|^2 + \left(\frac{\nu}{2} - \frac{\delta^2}{4} \right) \left\| \frac{\partial \hat{u}_1^\varepsilon}{\partial y} \right\|^2 \\ & + \left(\frac{\nu}{2} - \frac{\delta^2}{4} \right) \left\| \frac{\partial \hat{u}_2^\varepsilon}{\partial y} \right\|^2 + \frac{\nu \varepsilon^2}{2} \left\| \frac{\partial \hat{u}_2^\varepsilon}{\partial x_2} \right\|^2 + \varepsilon^2 \left(\frac{\nu}{2} - \frac{\delta^2}{4} \right) \left\| \frac{\partial \hat{u}_3^\varepsilon}{\partial y} \right\|^2 \\ & + \frac{\nu \varepsilon^4}{2} \left\| \frac{\partial \hat{u}_3^\varepsilon}{\partial x_1} \right\|^2 + \frac{\nu \varepsilon^4}{2} \left\| \frac{\partial \hat{u}_3^\varepsilon}{\partial x_2} \right\|^2 \leq C_0. \end{aligned} \quad (4.4)$$

where $\|\cdot\|$ denotes here the L^2 -norm in Ω , δ is the diameter of Ω , and C_0 is an independent constant of ε .

Proof. [6] Putting $\varphi_i = \hat{G}_i$ for $i = 1, 2$ and $\varphi_3 = \varepsilon \hat{G}_3$, in (5.8), leads to

$$\begin{aligned}
& \sum_{i,j=1}^2 \int_{\Omega} \left(\varepsilon^2 \nu \left(\frac{\partial \hat{u}_i^\varepsilon}{\partial x_j} + \frac{\partial \hat{u}_j^\varepsilon}{\partial x_i} \right) - \hat{p}^\varepsilon \delta_{i,j} \right) \frac{\partial \hat{u}_i^\varepsilon}{\partial x_j} dx dy + \int_{\Omega} \left(2\nu \varepsilon^2 \frac{\partial \hat{u}_3^\varepsilon}{\partial y} - \hat{p}^\varepsilon \right) \frac{\partial \hat{u}_3^\varepsilon}{\partial y} dx dy \\
& + \sum_{i=1}^2 \int_{\Omega} \nu \left(\frac{\partial \hat{u}_i^\varepsilon}{\partial y} + \varepsilon^2 \frac{\partial \hat{u}_3^\varepsilon}{\partial x_i} \right) \frac{\partial \hat{u}_i^\varepsilon}{\partial y} dx dy + \sum_{j=1}^2 \int_{\Omega} \varepsilon^2 \nu \left(\varepsilon^2 \frac{\partial \hat{u}_3^\varepsilon}{\partial x_j} + \frac{\partial \hat{u}_j^\varepsilon}{\partial y} \right) \frac{\partial \hat{u}_3^\varepsilon}{\partial x_j} dx dy \\
& \leq \sum_{i,j=1}^2 \int_{\Omega} \left(\varepsilon^2 \nu \left(\frac{\partial \hat{u}_i^\varepsilon}{\partial x_j} + \frac{\partial \hat{u}_j^\varepsilon}{\partial x_i} \right) - \hat{p}^\varepsilon \delta_{i,j} \right) \frac{\partial \hat{G}_i}{\partial x_j} dx dy + \int_{\Omega} \left(2\nu \varepsilon^2 \frac{\partial \hat{u}_3^\varepsilon}{\partial y} - \hat{p}^\varepsilon \right) \frac{\partial \hat{G}_3}{\partial y} dx dy \\
& + \sum_{i=1}^2 \int_{\Omega} \nu \left(\frac{\partial \hat{u}_i^\varepsilon}{\partial y} + \varepsilon^2 \frac{\partial \hat{u}_3^\varepsilon}{\partial x_i} \right) \frac{\partial \hat{G}_i}{\partial y} dx dy + \sum_{j=1}^2 \int_{\Omega} \varepsilon^2 \nu \left(\varepsilon^2 \frac{\partial \hat{u}_3^\varepsilon}{\partial x_j} + \frac{\partial \hat{u}_j^\varepsilon}{\partial y} \right) \frac{\partial \hat{G}_3}{\partial x_j} dx dy + \\
& + \int_{\omega} \hat{k} |\hat{G} - s| dx - \sum_{i=1}^2 \int_{\Omega} \hat{f}_i (\hat{G}_i - \hat{u}_i^\varepsilon) dx dy - \int_{\Omega} \varepsilon \hat{f}_3 (\hat{G}_3 - \hat{u}_3^\varepsilon) dx dy, \tag{4.5}
\end{aligned}$$

as \hat{k} is positive.

Using (3.3), the Poincaré inequality, $\varepsilon \leq 1$, and $2ab \leq a^2 + b^2$ we deduce

$$\begin{aligned}
& \frac{\nu \varepsilon^2}{2} \left\| \frac{\partial \hat{u}_1^\varepsilon}{\partial x_1} \right\|^2 + \frac{\nu \varepsilon^2}{2} \left\| \frac{\partial \hat{u}_1^\varepsilon}{\partial x_2} \right\|^2 + \frac{\nu \varepsilon^2}{2} \left\| \frac{\partial \hat{u}_2^\varepsilon}{\partial x_1} \right\|^2 + \left(\frac{\nu}{2} - \frac{\delta^2}{4} \right) \left\| \frac{\partial \hat{u}_1^\varepsilon}{\partial y} \right\|^2 + \left(\frac{\nu}{2} - \frac{\delta^2}{4} \right) \left\| \frac{\partial \hat{u}_2^\varepsilon}{\partial y} \right\|^2 \\
& + \frac{\nu \varepsilon^2}{2} \left\| \frac{\partial \hat{u}_2^\varepsilon}{\partial x_2} \right\|^2 + \varepsilon^2 \left(\frac{\nu}{2} - \frac{\delta^2}{4} \right) \left\| \frac{\partial \hat{u}_3^\varepsilon}{\partial y} \right\|^2 + \frac{\nu \varepsilon^4}{2} \left\| \frac{\partial \hat{u}_3^\varepsilon}{\partial x_1} \right\|^2 + \frac{\nu \varepsilon^4}{2} \left\| \frac{\partial \hat{u}_3^\varepsilon}{\partial x_2} \right\|^2 \leq \\
& \leq \nu \left\| \frac{\partial \hat{G}_1}{\partial x_1} \right\|^2 + \nu \left\| \frac{\partial \hat{G}_1}{\partial x_2} \right\|^2 + \nu \left\| \frac{\partial \hat{G}_2}{\partial x_1} \right\|^2 + \nu \left\| \frac{\partial \hat{G}_1}{\partial y} \right\|^2 + \nu \left\| \frac{\partial \hat{G}_2}{\partial y} \right\|^2 \\
& + \nu \left\| \frac{\partial \hat{G}_2}{\partial x_2} \right\|^2 + \nu \left\| \frac{\partial \hat{G}_3}{\partial y} \right\|^2 + \nu \left\| \frac{\partial \hat{G}_3}{\partial x_1} \right\|^2 + \nu \left\| \frac{\partial \hat{G}_3}{\partial x_2} \right\|^2
\end{aligned}$$

$$+ \|\hat{f}_1\| \|\hat{G}_1\| + \|\hat{f}_2\| \|\hat{G}_2\| + \|\hat{f}_3\| \|\hat{G}_3\| + (\|\hat{f}_1\|^2 + \|\hat{f}_2\|^2 + \|\hat{f}_3\|^2) + \text{const.} \|\hat{k}\|_{L^\infty(\omega)} = C_0,$$

thus (4.4) follows. \square

Theorem 5. *Assuming (4.1), and $\nu > \frac{\delta^2}{2}$ or $f = 0$, the following estimates on p^ε are satisfied.*

$$\left\| \frac{\partial \hat{p}^\varepsilon}{\partial x_i} \right\|_{H^{-1}(\Omega)} \leq C_1 \quad (i = 1, 2) \tag{4.6}$$

$$\left\| \frac{\partial \hat{p}^\varepsilon}{\partial y} \right\|_{H^{-1}(\Omega)} \leq \varepsilon \cdot C_2, \tag{4.7}$$

where C_1 and C_2 denote independent constants of ε .

Proof. [6] Let ψ in $H_0^1(\Omega)$, putting in (5.8) $\varphi_i = \hat{u}_i^\varepsilon$ (for $i = 1, 2$), and $\varphi_3 = \varepsilon \hat{u}_3^\varepsilon \pm \psi$, we deduce

$$\begin{aligned} & - \int_{\Omega} \hat{p}^\varepsilon \frac{\partial \psi}{\partial y} dx dy = - \int_{\Omega} 2\nu \varepsilon^2 \frac{\partial \hat{u}_3^\varepsilon}{\partial y} \frac{\partial \psi}{\partial y} dx dy - \\ & - \sum_{j=1}^2 \int_{\Omega^\varepsilon} \varepsilon^2 \nu \left(\varepsilon^2 \frac{\partial \hat{u}_3^\varepsilon}{\partial x_j} + \frac{\partial \hat{u}_j^\varepsilon}{\partial y} \right) \frac{\partial \psi}{\partial x_j} dx dy + \int_{\Omega} \varepsilon \hat{f}_3 \psi dx dy \end{aligned} \quad (4.8)$$

Taking in (5.8) $\varphi_1 = \hat{u}_1^\varepsilon \pm \psi$, ψ in $H_0^1(\Omega)$, $\varphi_2 = \hat{u}_2^\varepsilon$, $\varphi_3 = \varepsilon \hat{u}_3^\varepsilon$, we get

$$\begin{aligned} & - \int_{\Omega} \hat{p}^\varepsilon \frac{\partial \psi}{\partial x_1} dx dy = - \int_{\Omega} 2\varepsilon^2 \nu \frac{\partial \hat{u}_1^\varepsilon}{\partial x_1} \frac{\partial \psi}{\partial x_1} dx dy - \int_{\Omega} \varepsilon^2 \nu \left(\frac{\partial \hat{u}_1^\varepsilon}{\partial x_2} + \frac{\partial \hat{u}_2^\varepsilon}{\partial x_1} \right) \frac{\partial \psi}{\partial x_1} dx dy \\ & - \int_{\Omega} \nu \left(\frac{\partial \hat{u}_1^\varepsilon}{\partial y} + \varepsilon^2 \frac{\partial \hat{u}_3^\varepsilon}{\partial x_1} \right) \frac{\partial \psi}{\partial y} dx dy + \int_{\Omega} \hat{f}_1 \psi dx dy \quad \forall \psi \in H_0^1(\Omega) \end{aligned} \quad (4.9)$$

In the same way, the choice $\varphi_1 = \hat{u}_1^\varepsilon$, $\varphi_2 = \hat{u}_2^\varepsilon \pm \psi$, ψ in $H_0^1(\Omega)$, $\varphi_3 = \varepsilon \hat{u}_3^\varepsilon$, leads to

$$\begin{aligned} & - \int_{\Omega} \hat{p}^\varepsilon \frac{\partial \psi}{\partial x_2} dx dy = - \int_{\Omega} 2\varepsilon^2 \nu \frac{\partial \hat{u}_2^\varepsilon}{\partial x_2} \frac{\partial \psi}{\partial x_2} dx dy - \int_{\Omega} \varepsilon^2 \nu \left(\frac{\partial \hat{u}_1^\varepsilon}{\partial x_2} + \frac{\partial \hat{u}_2^\varepsilon}{\partial x_1} \right) \frac{\partial \psi}{\partial x_1} dx dy \\ & - \int_{\Omega} \nu \left(\frac{\partial \hat{u}_2^\varepsilon}{\partial y} + \varepsilon^2 \frac{\partial \hat{u}_3^\varepsilon}{\partial x_2} \right) \frac{\partial \psi}{\partial y} dx dy + \int_{\Omega} \hat{f}_2 \psi dx dy \quad \forall \psi \in H_0^1(\Omega) \end{aligned} \quad (4.10)$$

then from (4.8) using (4.4) we get (4), and from (4.9)-(4.10) using (4.4) we get (5.2). \square

We define now the Banach space

$$V_y = \{v \in (L^2(\Omega))^2 : \frac{\partial v}{\partial y} \in (L^2(\Omega))^2, v = 0 \text{ on } \Gamma_1\}$$

with its norm

$$\|v\|_{V_y} = \sum_{i=1}^2 \left(\|v_i\|_{L^2(\Omega)}^2 + \left\| \frac{\partial v_i}{\partial y} \right\|_{L^2(\Omega)}^2 \right).$$

Corollary 1. *Let the assumptions of Theorem 1 and Theorem 5 hold, then there exists u_i^* in V_y ($i = 1, 2$), and p^* in $L_0^2(\Omega)$ such that*

$$\hat{u}_i^\varepsilon \rightharpoonup u_i^* \quad (1 \leq i \leq 2) \quad \text{weakly in } V_y \quad (4.11)$$

where $V_y = \{\psi \in L^2(\Omega) \text{ such that } \frac{\partial \psi}{\partial y} \in L^2(\Omega)\}$.

$$\varepsilon \frac{\partial \hat{u}_i^\varepsilon}{\partial x_j} \rightharpoonup 0 \quad (1 \leq i, j \leq 2) \quad \text{weakly in } L^2(\Omega) \quad (4.12)$$

$$\varepsilon \frac{\partial \hat{u}_3^\varepsilon}{\partial y} \rightharpoonup 0 \quad \text{weakly in } L^2(\Omega) \quad (4.13)$$

$$\varepsilon^2 \frac{\partial \hat{u}_3^\varepsilon}{\partial x_i} \rightharpoonup 0 \quad (1 \leq i \leq 2) \quad \text{weakly in } L^2(\Omega) \quad (4.14)$$

$$\varepsilon \hat{u}_3^\varepsilon \rightharpoonup 0 \quad \text{weakly in } L^2(\Omega) \quad (4.15)$$

$$\hat{p}^\varepsilon \rightharpoonup p^* \quad \text{weakly in } L_0^2(\Omega) \quad (4.16)$$

Proof. [6] From (4.4) there exists a fixed constant C which does not depend on ε such that

$$\left\| \frac{\partial \hat{u}_i^\varepsilon}{\partial y} \right\| \leq C \quad (1 \leq i \leq 2)$$

using the above estimate and the Poincaré inequality in the domain Ω we deduce (4.11). Also (4.12)-(4.14) follows from (4.4), and (4.16) follows from (5.2), (4) and [54]. To prove (4.15), as in [7] we choose q such that $q(x, y) = y\theta(x) - \gamma$ where θ in $C_0^\infty(\omega)$ and

$$\gamma = \left(\int_{\Omega} y\theta dx dy \right) / \left(\int_{\Omega} dx dy \right).$$

Using the Green formula, the boundary conditions on Γ imply

$$-\sum_{i=1}^2 \int_{\Omega} y \hat{u}_i^\varepsilon \frac{\partial \theta}{\partial x_i} dx dy - \int_{\Omega} \theta \hat{u}_3^\varepsilon dx dy = 0,$$

As $\hat{u}_i^\varepsilon \rightharpoonup u_i^*$ in V_y ($i = 1, 2$), then (4.15) holds. \square

4.1. Study of the limit problem of the 1rst Case. We give both the equations satisfied by p^* and u^* in Ω and the inequalities for the trace of the velocity $u^*(x, 0)$ and the stress $\frac{\partial u^*}{\partial y}(x, 0)$ on $\partial\omega$.

Theorem 6. *With the same assumptions as in Theorem 5, (u^*, p^*) satisfy*

$$p^* \in H^1(\omega), \quad (4.17)$$

$$-\nu \frac{\partial^2 u_i^*}{\partial y^2} + \frac{\partial p^*}{\partial x_i} = \hat{f}_i \quad (i = 1, 2) \quad \text{in } L^2(\Omega). \quad (4.18)$$

Proof. [6] We choose in (5.8) $\varphi_3 = \hat{u}_3^\varepsilon \pm \psi$ with ψ in $H_0^1(\Omega)$ we deduce

$$\begin{aligned} \sum_{j=1}^2 \int_{\Omega^\varepsilon} \varepsilon^2 \nu \left(\varepsilon^2 \frac{\partial \hat{u}_3^\varepsilon}{\partial x_j} + \frac{\partial \hat{u}_j^\varepsilon}{\partial y} \right) \frac{\partial \psi}{\partial x_j} dx dy + \int_{\Omega} (2\nu \varepsilon^2 \frac{\partial \hat{u}_3^\varepsilon}{\partial y} - \hat{p}^\varepsilon) \frac{\partial \psi}{\partial y} dx dy \\ = \int_{\Omega} \varepsilon f_3 \psi dx dy \end{aligned}$$

Using (4.14) (4.11) (4.13) and the hypothesis of this theorem we obtain

$$\int_{\Omega} p^* \frac{\partial \psi_3}{\partial y} dx dy = 0 \quad \forall \psi \in H_0^1(\Omega),$$

then

$$\frac{\partial p^*}{\partial y} = 0 \quad \text{in } H^{-1}(\Omega). \quad (4.19)$$

Choosing now $\varphi_i = \hat{u}_i^\varepsilon \pm \psi_i$, for ($i = 1, 2$) with ψ_i in $H_0^1(\Omega)$ and $\varphi_3 = \varepsilon \hat{u}_3^\varepsilon$, in (5.8), leads to

$$\sum_{i,j=1}^2 \int_{\Omega} \left(\varepsilon^2 \nu \left(\frac{\partial \hat{u}_i^\varepsilon}{\partial x_j} + \frac{\partial \hat{u}_j^\varepsilon}{\partial x_i} \right) - \hat{p}^\varepsilon \delta_{i,j} \right) \frac{\partial \psi_i}{\partial x_j} dx dy +$$

$$+ \sum_{i=1}^2 \int_{\Omega} \nu \left(\frac{\partial \hat{u}_i^\varepsilon}{\partial y} + \varepsilon^2 \frac{\partial \hat{u}_3^\varepsilon}{\partial x_i} \right) \frac{\partial \psi_i}{\partial y} dx dy = \sum_{i=1}^2 \int_{\Omega} \hat{f}_i \psi_i dx dy \quad (4.20)$$

Using (4.12) (4.16) (4.11) (4.14) and the hypothesis of this theorem, we deduce first with $\psi_1 = 0$ and ψ_2 in $H_0^1(\Omega)$, then with $\psi_2 = 0$ and ψ_1 in $H_0^1(\Omega)$, the following equality

$$- \sum_{i=1}^2 \int_{\Omega} p^* \frac{\partial \psi_i}{\partial x_i} dx dy + \sum_{i=1}^2 \int_{\Omega} \nu \frac{\partial u_i^*}{\partial y} \frac{\partial \psi_i}{\partial y} dx dy = \sum_{i=1}^2 \int_{\Omega} \hat{f}_i \psi_i dx dy \quad (4.21)$$

then using the Green formula, we obtain

$$-\nu \frac{\partial^2 u_i^*}{\partial y^2} + \frac{\partial p^*}{\partial x_i} = \hat{f}_i \quad (i = 1, 2) \quad \text{in } H^{-1}(\Omega). \quad (4.22)$$

To prove that p^* is in $H^1(\omega)$, let us recall first that p^* is a function of (x_1, x_2) only from (5), then following [7] we choose ψ_i in (4.21) such that $\psi_i(x, y) = y(y - h(x))\theta(x)$ with θ in $H_0^1(\omega)$, and using the Green formula we deduce

$$\frac{1}{6} \int_{\omega} p^* \frac{\partial (h^3 \theta)}{\partial x_i} dx - 2\nu \int_{\omega} h \tilde{u}_i^* \theta dx = \int_{\omega} \tilde{f}_i \theta dx$$

where

$$\tilde{u}_i^*(x) = \frac{1}{h(x)} \int_0^{h(x)} u_i^*(x, y) dy, \quad \text{and} \quad \tilde{f}_i(x) = \int_0^{h(x)} y(y - h(x)) \hat{f}_i(x, y) dy.$$

Whence

$$2\nu h \tilde{u}_i^* - \frac{1}{6} h^3 \frac{\partial p^*}{\partial x_i} = \tilde{f}_i \quad (i = 1, 2) \quad \text{in } H^{-1}(\omega). \quad (4.23)$$

As f_i is in $L^2(\Omega)$, u_i^* in V_y then in $L^2(\Omega)$, therefore \tilde{f}_i and \tilde{u}_i^* are in $L^2(\omega)$, then from (4.23) we get p^* in $H^1(\omega)$, then (5) follows. So as f_i belongs to $L^2(\Omega)$, then from (5.10) we have $\frac{\partial^2 u_i^*}{\partial y^2}$ in $L^2(\Omega)$. Whence (5.2) holds, and we also have $\frac{\partial u_i^*}{\partial y}$ in V_y . \square

For convenience, we will denote by $s^*(x) = u^*(x, 0)$ and $\tau^*(x) = \frac{\partial u^*}{\partial y}(x, 0)$, as $\frac{\partial u^*}{\partial y}$ in V_y then τ^* belongs to $L^2(\omega)$, and we have :

Theorem 7. *Under the same hypothesis of Theorem 5, (s^*, τ^*) satisfy the following inequalities*

$$\int_{\omega} \hat{k} (|\psi + s^* - s|) - |s^* - s| dx - \int_{\omega} \nu \tau^* \psi dx \geq 0 \quad \forall \psi \in (L^2(\omega))^2 \quad (4.24)$$

$$\left. \begin{array}{l} \nu |\tau^*| = \hat{k} \implies \exists \lambda \geq 0 \quad s^* = s + \lambda \tau^* \\ \nu |\tau^*| < \hat{k} \implies s^* = s \end{array} \right\} \text{ a.e. in } \omega \quad (4.25)$$

where $|\cdot|$ denotes the \mathbb{R}^2 Euclidean norm.

Proof. [6] Choosing $\varphi = (\varphi_1, \varphi_2, \varepsilon \hat{u}_3^\varepsilon)$ with $\varphi_i = \hat{u}_i^\varepsilon + \psi_i$, for $(i = 1, 2)$ and ψ_i in $H_{\Gamma_1 \cap \Gamma_L}^1(\omega)$ where $H_{\Gamma_1 \cap \Gamma_L}^1(\omega) = \{v \in H^1(\Omega) : v = 0 \text{ on } \Gamma_1 \cap \Gamma_L\}$, in (5.8), leads to

$$\sum_{i,j=1}^2 \int_{\Omega} \left(\varepsilon^2 \nu \left(\frac{\partial \hat{u}_i^\varepsilon}{\partial x_j} + \frac{\partial \hat{u}_j^\varepsilon}{\partial x_i} \right) - \hat{p}^\varepsilon \delta_{i,j} \right) \frac{\partial \psi_i}{\partial x_j} dx dy + \sum_{i=1}^2 \int_{\Omega} \nu \left(\frac{\partial \hat{u}_i^\varepsilon}{\partial y} + \varepsilon^2 \frac{\partial \hat{u}_3^\varepsilon}{\partial x_i} \right) \frac{\partial \psi_i}{\partial y} dx dy$$

$$+ \int_{\omega} \hat{k}(|\psi + \hat{u}^{\varepsilon} - s| - |\hat{u}^{\varepsilon} - s|) dx \geq \sum_{i=1}^2 \int_{\Omega} \hat{f}_i \psi_i dx dy. \quad (4.26)$$

Using Corollary 1, we can pass to the limit in (4.26), to obtain

$$\begin{aligned} & \sum_{i=1}^2 \int_{\Omega} -p^* \frac{\partial \psi_i}{\partial x_i} dx dy + \sum_{i=1}^2 \int_{\Omega} \nu \frac{\partial u_i^*}{\partial y} \frac{\partial \psi_i}{\partial y} dx dy + \\ & + \int_{\omega} \hat{k}(|\psi + s^* - s| - |s^* - s|) dx \geq \sum_{i=1}^2 \int_{\Omega} \hat{f}_i \psi_i dx dy \end{aligned}$$

Using now the Green formula, the equality (5.2) and the fact that $\psi_i = 0$ on $\Gamma_1 \cap \Gamma_L$ and $\cos(n, x_i) = 0$ on ω , we deduce

$$\int_{\omega} \hat{k}(|\psi + s^* - s| - |s^* - s|) dx - \int_{\omega} \nu \tau^* \psi dx \geq 0 \quad \forall \psi \in (H_{\Gamma_1 \cup \Gamma_L}^1(\Omega))^2. \quad (4.27)$$

This inequality remains valid for any ψ in $(D(\omega))^2$ (using the same notations for the trace) and by density of $D(\omega)$ in $L^2(\omega)$ for any ψ in $(L^2(\omega))^2$. Then (5.1) follows.

To prove (4.25), we take $\psi_i = \pm(s_i^* - s_i)$, in (5.1), we obtain

$$\int_{\omega} \left(\hat{k}|s^* - s| - \nu \tau^*(s^* - s) \right) dx = 0, \quad (4.28)$$

taking $\psi = \phi - (s^* - s)$ with ϕ in $(L^2(\omega))^2$, in (5.1), we obtain

$$\int_{\omega} \left(\hat{k}|\phi| - \nu \tau^* \phi \right) dx \geq \int_{\omega} \left(\hat{k}|s^* - s| - \nu \tau^*(s^* - s) \right) dx.$$

And from (4.28) we deduce

$$\int_{\omega} \left(\hat{k}|\phi| - \nu \tau^* \phi \right) dx \geq 0 \quad \forall \phi \in (L^2(\omega))^2, \quad (4.29)$$

taking first $\phi = (\varphi_1, \varphi_2)$ such that $\varphi_i \geq 0$ $i = 1, 2$, in (4.29), we obtain :

$$\int_{\omega} \left(\hat{k}|\phi| - \nu |\tau^*| \cdot |\phi| \cos(\tau^*, \phi) \right) dx = \int_{\omega} \left(\hat{k} - \nu |\tau^*| \cos(\tau^*, \phi) \right) |\phi| dx \geq 0,$$

then:

$$\nu |\tau^*| \cos(\tau^*, \phi) \leq \hat{k} \quad \text{a.e. on } \omega, \quad (4.30)$$

taking now $-\phi$, with $\phi = (\varphi_1, \varphi_2)$ such that $\varphi_i \geq 0$ $i = 1, 2$, in (4.29), we obtain :

$$\int_{\omega} \left(\hat{k}|\phi| + \nu |\tau^*| \cdot |\phi| \cos(\tau^*, \phi) \right) dx = \int_{\omega} \left(\hat{k} + \nu |\tau^*| \cos(\tau^*, \phi) \right) |\phi| dx \geq 0,$$

whence

$$\nu |\tau^*| \cos(\tau^*, \phi) \geq -\hat{k} \quad \text{a.e. on } \omega, \quad (4.31)$$

from (4.30) and (4.31) we get:

$$\nu |\tau^*| \leq \hat{k} \quad \text{a.e. on } \omega, \quad (4.32)$$

then

$$\hat{k}|s^* - s| \geq \nu |\tau^*| \cdot |s^* - s| \geq \nu \tau^* \cdot (s^* - s) \quad \text{a.e. on } \omega$$

so

$$\hat{k}|s^* - s| - \nu\tau^* \cdot (s^* - s) \geq 0 \quad a.e. \quad \text{on } \omega$$

and from (4.28) we deduce that

$$\hat{k}|s^* - s| - \nu\tau^* \cdot (s^* - s) = 0 \quad a.e. \quad \text{on } \omega. \quad (4.33)$$

If $\nu|\tau^*| = \hat{k}$, then from (4.33) we have

$$\nu|\tau^*| \cdot |s^* - s| = \nu\tau^* \cdot (s^* - s) \quad a.e. \quad \text{on } \omega,$$

then $\cos(s^* - s, \nu\tau^*) = 1$, which implies the existence of $\lambda \geq 0$ such that $s^* - s = \lambda\nu\tau^*$. And if $\nu|\tau^*| < \hat{k}$, then from (4.33) we have

$$\hat{k}|s^* - s| - \nu\tau^* \cdot (s^* - s) = 0 \geq (\hat{k} - |\nu\tau^*|)|s^* - s| \quad a.e. \quad \text{on } \omega,$$

whence $s^* - s = 0 \quad a.e. \quad \text{on } \omega$. then (4.25) follows. \square

Theorem 8. *Under the same hypothesis of in Theorem 5, and assuming that \hat{f} is a function of x only, we have*

$$\frac{h^2}{2} \nabla p^*(x) + \nu s^*(x) + \nu h \tau^*(x) - \frac{h^2}{2} \hat{f}(x) = 0 \quad a.e. \quad \text{in } \omega. \quad (4.34)$$

$$\int_{\omega} (h^2 \tau^*(x) + 4h s^*(x)) \nabla \varphi(x) dx = 6 \int_{\partial\omega} \varphi(x) \tilde{g}(x) \cdot n \quad \forall \varphi \in H^1(\omega). \quad (4.35)$$

Proof. [6] Integrate twice (5.2) between 0 and y we obtain

$$\nu u_i^*(x, y) = \frac{y^2}{2} \frac{\partial p^*(x)}{\partial x_i} + \nu u_i^*(x, 0) + \nu y \frac{\partial u_i^*(x, 0)}{\partial y} - \frac{y^2}{2} \hat{f}_i(x),$$

and as $u_i^*(x, h) = 0$, then (4.34) follows. On the other hand, taking the average of the preceding expression we have

$$h\nu \tilde{u}_i^*(x) = \int_0^{h(x)} \nu u_i^*(x, y) dy = \frac{h^3}{6} \frac{\partial p^*(x)}{\partial x_i} + \nu h u_i^*(x, 0) + \nu \frac{h^2}{2} \frac{\partial u_i^*(x, 0)}{\partial y} - \frac{h^3}{6} \hat{f}_i(x) \quad (4.36)$$

Otherwise, for all φ in $H^1(\omega)$, and as $\text{div}(\hat{u}^\varepsilon) = 0$ in Ω we have:

$$\begin{aligned} \int_{\Omega} \varphi \text{div}(\hat{u}^\varepsilon) dx dy &= 0 = \int_{\omega} \varphi(x) \int_0^h \left(\sum_{i=1}^2 \frac{\partial \hat{u}_i^\varepsilon}{\partial x_i} + \frac{\partial \hat{u}_3^\varepsilon}{\partial y} \right) dy = \\ &= \int_{\omega} \varphi(x) \sum_{i=1}^2 \left(\frac{\partial (h \tilde{u}_i)}{\partial x_i} + \hat{u}_3^\varepsilon(x, h) - \hat{u}_3^\varepsilon(x, 0) \right) dx \end{aligned}$$

then as $\hat{u}_3^\varepsilon = 0$ on $\partial\Omega = \bar{\omega} \cup \bar{\Gamma}_1 \cup \bar{\Gamma}_L$, we have

$$\int_{\omega} \varphi(x) \sum_{i=1}^2 \frac{\partial (h \tilde{u}_i)}{\partial x_i} dx = 0, \quad \text{where} \quad \tilde{u}_i(x) = \frac{1}{h(x)} \int_0^{h(x)} \hat{u}_i^\varepsilon(x, y) dy, \quad \forall x \in \omega,$$

$$\text{and} \quad \tilde{g}_i(x) = \int_0^{h(x)} \hat{g}_i(x, y) dy = h(x) \tilde{u}_i(x), \quad \forall x \in \partial\omega.$$

Using Green's formula we have

$$\sum_{i=1}^2 \int_{\omega} h \tilde{q}_i \frac{\partial \varphi}{\partial x_i} dx = \sum_{i=1}^2 \int_{\partial \omega} h \tilde{q}_i \varphi \cdot \cos(n, x_i) = \sum_{i=1}^2 \int_{\partial \omega} \tilde{g}_i(x) \varphi \cdot \cos(n, x_i)$$

as $\hat{u}_i^\varepsilon \rightharpoonup u_i^*$ in V_y then in $L^2(\omega)$, therefore $\tilde{q}_i \rightharpoonup \tilde{u}_i^*$ in $L^2(\omega)$, and as $\partial \omega \subset \partial \Omega$, we deduce :

$$\sum_{i=1}^2 \int_{\omega} h \tilde{u}_i^* \frac{\partial \varphi}{\partial x_i} dx = \sum_{i=1}^2 \int_{\partial \omega} \varphi(x) \tilde{g}_i(x) \cos(n, x_i) \quad \forall \varphi \in H^1(\omega). \quad (4.37)$$

From (4.36) we have

$$\int_{\omega} \left(\frac{h^3}{6\nu} \nabla p^* + h s^* + \frac{h^2}{2} \tau^* - \frac{h^3}{6\nu} \hat{f} \right) \nabla \varphi dx = \int_{\partial \omega} \varphi \tilde{g} \cdot n. \quad (4.38)$$

Then using (4.34) and (4.38), we obtain the weak formulation of the Reynolds equation:

$$\int_{\omega} \left(\frac{h^3}{12\nu} \nabla p^* - \frac{h}{2} s^* - \frac{h^3}{12\nu} \hat{f} \right) \nabla \varphi dx = \int_{\partial \omega} \varphi \tilde{g} \cdot n. \quad (4.39)$$

Using once again (4.34) and (4.39) we get (4.35). \square

4.2. Study of the uniqueness. In this subsection, we will give another formulation of the limit inequalities for s^* and τ^* on ω which enables us to express s^* as a solution of a variational inequality of the second kind with a convenient decomposition. The basic idea is that we have three unknowns s^* , τ^* and ∇p^* and three relations (4.34) (4.35) and (5.1). A test function in (4.35) appears only to be a gradient function. So it is only possible to control the "gradient" part of s^* and τ^* by this equation which is obtained by a slightly modified version of the well known decomposition of $L^2(\omega)^2$, due to the non constant $h(x)$ coefficients.

Lemma 2. *Let h in $L^\infty(\omega) \cap H^1(\omega)$ such that $h \geq \alpha > 0$. Every function ψ in $(L^2(\omega))^2$ has the following orthogonal decomposition:*

$$\psi = h^2 \nabla \varphi + h^{-1} \mathbf{curl}(\theta) \quad (4.40)$$

where φ in $H^1(\omega)/\mathbb{R}$ is the only solution of the problem

$$\int_{\omega} h^3 \nabla \varphi \nabla \mu dx = \int_{\omega} h \psi \nabla \mu dx \quad \forall \mu \in H^1(\omega), \quad (4.41)$$

and θ in $H_0^1(\omega)$ is the only solution of the problem

$$\int_{\omega} \mathbf{curl}(\theta) \mathbf{curl}(\xi) dx = \int_{\omega} (h \psi - h^3 \nabla \varphi) \mathbf{curl}(\xi) dx \quad \forall \xi \in H_0^1(\omega). \quad (4.42)$$

Proof. [6] As h in $L^\infty(\omega)$, for all ψ in $(L^2(\omega))^2$, we have $h \psi$ in $(L^2(\omega))^2$, following [33](theorem 3.2), the Neumann's problem (4.41) has a unique solution φ in $H^1(\omega)/\mathbb{R}$. This solution φ satisfies $\nabla(h \psi - h^3 \nabla \varphi) = 0$ in $H^{-1}(\omega)$. Hence $h \psi - h^3 \nabla \varphi$ is a divergence-free vector of $H(\operatorname{div}, \omega)$. Moreover, Green's formula applied to (4.41) yields:

$$0 = \int_{\omega} (h \psi - h^3 \nabla \varphi) \nabla \mu dx = \int_{\partial \omega} (h \psi - h^3 \nabla \varphi) \cdot n \mu \quad \forall \mu \in H^1(\omega),$$

implying that $(h\psi - h^3\nabla\varphi).n = 0$ in $H^{-1/2}(\partial\omega)$. Whence $h\psi - h^3\nabla\varphi$ lies in the space $H = \{v \in (L^2(\omega))^2 : \operatorname{div}(v) = 0, v.n = 0\}$. Moreover, as ω is connected, we deduce, from [33] (Theorem 3.1 and its corollary), that the space H is characterized by $H = \{\mathbf{curl}(\mu) : \mu \in H_0^1(\omega)\}$, and the mapping \mathbf{curl} is an isomorphism from $H_0^1(\omega)$ onto H . So there exists a unique stream function θ in $H_0^1(\omega)$ of $h\psi - h^3\nabla\varphi$ satisfying (4.40) and (4.42). \square

Theorem 9. *Let h in $L^\infty(\omega) \cap H^1(\omega)$. Under the same hypothesis of Theorem 5, s^* is uniquely given by $s^* = h^2\nabla C + h^{-1}\mathbf{curl}(D)$, where $U = (C, D)$ is the unique solution of the following variational problem: Find U in $H^1(\omega) \times H_0^1(\omega)$ such that*

$$a(U, \phi - U) + J(\phi) - J(U) \geq L(\phi - U) \quad \forall \phi = (\varphi, \theta) \in H^1(\omega) \times H_0^1(\omega), \quad (4.43)$$

$$\text{where } a(U, \phi) = \int_{\omega} 4\nu h^3 \nabla C \nabla \varphi dx + \int_{\omega} \nu h^{-3} \mathbf{curl}(D) \mathbf{curl}(\theta) dx,$$

$$J(\phi) = \int_{\omega} \hat{k} (|h^2 \nabla \varphi + h^{-1} \mathbf{curl}(\theta) - s|) dx,$$

$$L\phi = \frac{1}{2} \int_{\omega} \hat{f} \mathbf{curl}(\theta) dx + \int_{\partial\omega} 6\nu \tilde{g}.n\varphi.$$

Proof. [6] From (5.1) and the orthogonal decomposition of ψ , we have

$$\begin{aligned} & \int_{\omega} \hat{k} (|h^2 \nabla \varphi + h^{-1} \mathbf{curl}(\theta) + s^* - s| - |s^* - s|) dx \geq \\ & \geq \int_{\omega} \nu \tau^* h^2 \nabla \varphi + \int_{\omega} \nu \tau^* h^{-1} \mathbf{curl}(\theta) dx \quad \forall (\varphi, \theta) \in H^1(\omega) \times H_0^1(\omega) \end{aligned} \quad (4.44)$$

and from (4.35), we have

$$\int_{\omega} \nu h^2 \tau^* \nabla \varphi = - \int_{\omega} 4\nu h s^* \nabla \varphi + \int_{\partial\omega} 6\nu \tilde{g}.n\varphi \quad \forall \varphi \in H^1(\omega) \quad (4.45)$$

then from (4.44) and (4.45), we have for all (φ, θ) in $H^1(\omega) \times H_0^1(\omega)$

$$\begin{aligned} & \int_{\omega} \hat{k} (|h^2 \nabla \varphi + h^{-1} \mathbf{curl}(\theta) + s^* - s| - |s^* - s|) dx \geq \\ & \geq - \int_{\omega} 4\nu h s^* \nabla \varphi + \int_{\partial\omega} 6\nu \tilde{g}.n\varphi + \int_{\omega} \nu \tau^* h^{-1} \mathbf{curl}(\theta) dx. \end{aligned} \quad (4.46)$$

Now as s^* in $(L^2(\omega))^2$, we can use its orthogonal decomposition as $s^* = h^2\nabla C + h^{-1}\mathbf{curl}(D)$, then we deduce for all (φ, θ) in $H^1(\omega) \times H_0^1(\omega)$

$$\begin{aligned} & \int_{\omega} \hat{k} |h^2 \nabla \varphi + h^{-1} \mathbf{curl}(\theta) + h^2 \nabla C + h^{-1} \mathbf{curl}(D) - s| dx \\ & - \int_{\omega} \hat{k} |h^2 \nabla C + h^{-1} \mathbf{curl}(D) - s| dx \geq - \int_{\omega} 4\nu h^3 \nabla C \nabla \varphi \\ & - 4\nu \int_{\omega} \mathbf{curl}(D) \nabla \varphi + \int_{\partial\omega} 6\nu \tilde{g}.n\varphi + \int_{\omega} \nu \tau^* h^{-1} \mathbf{curl}(\theta) dx. \end{aligned} \quad (4.47)$$

Using (4.34) we have

$$\int_{\omega} \nu \tau^* h^{-1} \mathbf{curl}(\tau^*) dx = \int_{\omega} \left(-\frac{1}{2} \nabla p^* - \frac{\nu}{h^2} s^* + \frac{1}{2} \hat{f} \right) \mathbf{curl}(\theta) dx,$$

then

$$\begin{aligned} \int_{\omega} \nu \tau^* h^{-1} \mathbf{curl}(\theta) dx &= - \int_{\omega} \frac{1}{2} \mathbf{curl}(\theta) \nabla p^* dx - \int_{\omega} \nu \mathbf{curl}(\theta) \nabla C dx \\ &\quad - \int_{\omega} \nu h^{-3} \mathbf{curl}(D) \mathbf{curl}(\theta) dx + \frac{1}{2} \int_{\omega} \hat{f} \mathbf{curl}(\theta) dx. \end{aligned}$$

Using Green's formula and that θ in $H_0^1(\omega)$, we have

$$\int_{\omega} \mathbf{curl}(\theta) \nabla p^* dx = - \langle p^*, \operatorname{div}(\mathbf{curl}(\theta)) \rangle + \int_{\partial\omega} \mathbf{curl}(\theta) \cdot n p^* dx = 0$$

by the same argument we also have

$$\int_{\omega} \nu \mathbf{curl}(\theta) \nabla C = \int_{\omega} \mathbf{curl}(D) \nabla \varphi = 0.$$

Then from (4.47) $U = (C, D)$ satisfies for all $\phi = (\varphi, \theta)$ in $H^1(\omega) \times H_0^1(\omega)$

$$\begin{aligned} &\int_{\omega} \{ 4\nu h^3 \nabla C \nabla \varphi + \nu h^{-3} \mathbf{curl}(D) \mathbf{curl}(\theta) \} dx \\ &\quad + \int_{\omega} \hat{k} |h^2 \nabla(\varphi + C) + h^{-1} \mathbf{curl}(\theta + D) - s| dx \\ &\quad - \int_{\omega} \hat{k} |h^2 \nabla C + h^{-1} \mathbf{curl}(D) - s| dx \geq \int_{\omega} \frac{1}{2} \hat{f} \mathbf{curl}(\theta) dx + \int_{\partial\omega} 6\nu \tilde{g} \cdot n \varphi, \end{aligned}$$

taking $\tilde{\varphi} = \varphi + C$ and $\tilde{\theta} = \theta + D$ we deduce the variational inequality (5.11).

As the bilinear form $a(\cdot, \cdot)$ is continuous and coercive, the functional J is convex, proper and continuous, and the linear form L is continuous, the existence and uniqueness of (C, D) in $H^1(\omega) \times H_0^1(\omega)$ follows, and implies the existence and uniqueness of s^* in $(L^2(\omega))^2$. \square

Theorem 10. *Under the same hypothesis of Theorem 9, there exists a unique solution p^* in $H^1(\omega)$ satisfying the weak formulation of the Reynolds equation (4.39). Then τ^* is then unique.*

Proof. [6] From Theorem 9 s^* is unique in $(L^2(\omega))^2$, then the uniqueness of p^* follows from (4.39). Finally τ^* is unique from the uniqueness of p^* and s^* using (4.34). \square

4.3. **Case 2.** We introduce in this case see [9] the definition that $v = (v_1, v_2) \in (L^2(\Omega))^2$ satisfies the condition (D') if

$$\int_{\Omega} \left(v_1 \frac{\partial \theta}{\partial x_1} + v_2 \frac{\partial \theta}{\partial x_2} \right) dx dy = 0, \quad \forall \theta \in C_0^\infty(\omega).$$

And consider

$$\Pi(V_{div}) = \{ \bar{\varphi} \in (H^1(\Omega))^2 : \bar{\varphi} = (\varphi_1, \varphi_2), \quad \varphi_i = \hat{G}_i \quad \text{on} \quad \Gamma_1 \cup \Gamma_L, i = 1, 2 \}$$

$$\tilde{\Pi}(V_{div}) = \{ \bar{\varphi} \in \Pi(V_{div}) : \bar{\varphi} \text{ satisfies condition } (D') \},$$

and also the Banach space

$$V_y = \{ v = (v_1, v_2) \in (L^2(\Omega))^2 : \frac{\partial v_i}{\partial y} \in L^2(\Omega), i = 1, 2, \quad v = 0 \quad \text{on} \quad \Gamma_1 \}$$

with its norm $\| \cdot \|_{V_y}$,

$$\| v \|_{V_y}^2 = \sum_{i=1}^2 \left(\| v_i \|_{L^2(\Omega)}^2 + \left\| \frac{\partial v_i}{\partial y} \right\|_{L^2(\Omega)}^2 \right),$$

and define its linear subspace (endowed with the same topology)

$$\tilde{V}_y = \{ v \in V_y : v \text{ satisfies condition } (D') \}.$$

We obtain also the following similar main results

Theorem 11. [9] *There exist $u^* = (u_1^*, u_2^*)$ in \tilde{V}_y , p^* in $L_0^2(\Omega)$, and a subsequence $\varepsilon \rightarrow 0$ such that (4.11)-(4.16) hold. We have also, $\hat{u}_i^\varepsilon \rightarrow u_i^*$ strongly in V_y for $i = 1, 2$. The limit functions u^*, p^* satisfy (5)-(5.2). Moreover u^*, p^* satisfy the inequality*

$$\begin{aligned} & \int_{\Omega} \nu \frac{\partial u^*}{\partial y} \frac{\partial (\varphi - u^*)}{\partial y} dx dy - \int_{\Omega} p^* \left(\frac{\partial \varphi_1}{\partial x_1} + \frac{\partial \varphi_2}{\partial x_2} \right) dx dy \\ & + \int_{\omega} \hat{k} S(-p^*) (|\varphi - s| - |u^* - s|) dx \geq \sum_{i=1}^2 \int_{\Omega} \hat{f}_i (\varphi_i - u_i^*) dx dy \quad \forall \varphi \in \Pi(K). \end{aligned} \quad (4.48)$$

Theorem 12. [9] *The pair (u^*, p^*) satisfies the same weak form of the Reynolds equation (4.39) where $u^* = u^*(\cdot, 0)$. Moreover, the traces $\tau^* = \frac{\partial u^*}{\partial y}(x, 0)$ and $u^*(x, 0)$ satisfy the following limit form of the Coulomb boundary conditions (2.16)*

$$\left. \begin{aligned} \nu |\tau^*| = \hat{k} S(-p^*) &\implies \exists \lambda \geq 0 \quad u^* = s + \lambda \tau^* \\ \nu |\tau^*| < \hat{k} S(-p^*) &\implies u^* = s \end{aligned} \right\} \text{ a.e. in } \omega.$$

Theorem 13. [9] *There exists a positive constant k^* such that for $\|\hat{k}\|_{L^\infty(\omega)} \leq k^*$ the solution (u^*, p^*) in $\tilde{V}_y \times (L_0^2(\omega) \cap H^1(\omega))$ of inequality ((5.1) is unique.*

5. STUDY OF THE case 3

For this case (see [10]) two technical difficulties to study the asymptotic analysis of this problem. The first, we cannot use the usual Korn inequality as we do not assume that the velocity vanishes at one of the boundaries (on the top or the bottom) as is usually assumed in lubrication problems. We thus derive an analogue of the Korn inequality suitable for our boundary conditions and such that the constants can be controlled appropriately as the gap between the surfaces approach zero. This leads us to the main uniform estimate of the velocity fields and to the limit variational inequality, in consequence. The second, to be able to make use of the latter, we have to characterize precisely the limit solution space and the set of admissible test functions. As the limit variational inequality is written in terms of the first two components of the velocity field, we have to characterize - in this very limit case - projections of the convexes appearing in the weak form of the Stokes flow. This allows us, in particular, to obtain a stronger convergence of the velocity fields as in usually expected. We give here only the main results

Lemma 3. [10] "Poincaré inequality"

$$\int_{\Omega^\varepsilon} |u|^2 \leq 2\varepsilon h_M \int_{\Gamma_1^\varepsilon} |u|^2 + 2(\varepsilon h_M)^2 \int_{\Omega^\varepsilon} \left| \frac{\partial u}{\partial x_3} \right|^2. \quad (5.1)$$

Proof.

$$u(x, t) = u(x, h^\varepsilon(x)) - \int_t^{h^\varepsilon(x)} \frac{\partial u}{\partial z}(x, z) dz$$

We integrate over $t \in [0, h^\varepsilon(x)]$ to get

$$\int_0^{h^\varepsilon(x)} |u(x, t)|^2 dt \leq 2h^\varepsilon(x) |u(x, h^\varepsilon(x))|^2 + 2(h^\varepsilon(x))^2 \int_0^{h^\varepsilon(x)} \left| \frac{\partial u}{\partial z}(x, z) \right|^2 dz,$$

and, after integration over ω we get (5.1). □

Lemma 4. [10] "Korn's Inequality" assumming that $h \in C^2(\bar{\omega})$, we have

$$\int_{\Omega^\varepsilon} |\nabla(u - G^\varepsilon)|^2 \leq a(u - G^\varepsilon, u - G^\varepsilon) + C(\Gamma_1^\varepsilon) \int_{\Gamma_1^\varepsilon} |u - G^\varepsilon|^2,$$

where

$$C(\Gamma_1^\varepsilon) = 2\varepsilon \|D_2 h\|_{C(\bar{\omega})} (1 + \varepsilon^2 \|D_1 h\|_{C(\bar{\omega})}^2).$$

Observe that $C(\Gamma_1^\varepsilon)$ est d'ordre ε .

Proof. We have

$$\begin{aligned} a(v, v) &= \frac{1}{2} \int_{\Omega^\varepsilon} \left(\frac{\partial v_i}{\partial x_k} + \frac{\partial v_k}{\partial x_i} \right)^2 = \int_{\Omega^\varepsilon} \left(\frac{\partial v_i}{\partial x_k} \frac{\partial v_i}{\partial x_k} + \frac{\partial v_i}{\partial x_k} \frac{\partial v_k}{\partial x_i} \right) = \int_{\Omega^\varepsilon} |\nabla v|^2 + \int_{\Omega^\varepsilon} \frac{\partial v_i}{\partial x_k} \frac{\partial v_k}{\partial x_i} \\ &= \int_{\Omega^\varepsilon} |\nabla v|^2 - \int_{\Omega^\varepsilon} \frac{\partial^2 v_i}{\partial x_k \partial x_i} v_k + \int_{\partial\Omega^\varepsilon} \frac{\partial v_i}{\partial x_k} v_k n_i \\ &= \int_{\Omega^\varepsilon} |\nabla v|^2 + \int_{\Omega^\varepsilon} \frac{\partial v_i}{\partial x_i} \frac{\partial v_k}{\partial x_k} - \int_{\partial\Omega^\varepsilon} \frac{\partial v_i}{\partial x_i} v_k n_k + \int_{\partial\Omega^\varepsilon} \frac{\partial v_i}{\partial x_k} v_k n_i. \end{aligned}$$

As $\operatorname{div} v = 0$ then we have

$$\int_{\Omega^\varepsilon} |\nabla v|^2 = a(v, v) + \int_{\partial\Omega^\varepsilon} \frac{\partial v_i}{\partial x_i} v_k n_k - \int_{\partial\Omega^\varepsilon} \frac{\partial v_i}{\partial x_k} v_k n_i.$$

For $v = u - G^\varepsilon$, we have $(u - G^\varepsilon)n|_{\partial\Omega^\varepsilon} = 0$, so

$$\int_{\partial\Omega^\varepsilon} \frac{\partial v_i}{\partial x_i} v_k n_k = 0,$$

in other hand, $u - G^\varepsilon = 0$ sur Γ_L^ε . So

$$\int_{\partial\Omega^\varepsilon} \frac{\partial v_i}{\partial x_k} v_k n_i = \int_{\Gamma_1^\varepsilon \cup \omega} \frac{\partial v_i}{\partial x_k} v_k n_i = \int_{\Gamma_1^\varepsilon \cup \omega} (u_k - G_k^\varepsilon) \frac{\partial (u_i - G_i^\varepsilon)}{\partial x_k} n_i.$$

as

$$(u - G^\varepsilon) \cdot n|_{\Gamma_1^\varepsilon \cup \omega} = 0, \quad \text{alors} \quad \frac{\partial}{\partial x_k} \{(u - G^\varepsilon) \cdot n\} = 0,$$

that is

$$\frac{\partial (u_i - G_i^\varepsilon)}{\partial x_k} n_i = -(u_i - G_i^\varepsilon) \frac{\partial n_i}{\partial x_k}.$$

but $\frac{\partial n_i}{\partial x_k}|_\omega = 0$, then from (5.2), (5) we have

$$\left| \int_{\partial\Omega^\varepsilon} \frac{\partial v_i}{\partial x_k} v_k n_i \right| = \left| \int_{\Gamma_1^\varepsilon} (u_k - G_k^\varepsilon) \frac{\partial (u_i - G_i^\varepsilon)}{\partial x_k} n_i \right| \leq 2 \max_{q \in \Gamma_1^\varepsilon} \left| \frac{\partial n_i}{\partial x_k}(q) \right| \int_{\Gamma_1^\varepsilon} |u - G^\varepsilon|^2,$$

Γ_1^ε is given by $x_3 = h^\varepsilon(x_1, x_2)$, then the unit normal vector exterior to Γ_1^ε can be written

$$n(q) = \frac{\left(-\frac{\partial h^\varepsilon}{\partial x_1}(x_1, x_2), -\frac{\partial h^\varepsilon}{\partial x_2}(x_1, x_2), 1 \right)}{\sqrt{1 + |\nabla h^\varepsilon(x)|^2}} = n(x_1, x_2)$$

For $i = 1, 2$, we have

$$\frac{\partial n_3}{\partial x_i}(x_1, x_2) = -(1 + |\nabla h^\varepsilon|^2)^{-\frac{3}{2}} \frac{\frac{\partial h^\varepsilon}{\partial x_1} \frac{\partial^2 h^\varepsilon}{\partial x_i \partial x_1} + \frac{\partial h^\varepsilon}{\partial x_2} \frac{\partial^2 h^\varepsilon}{\partial x_i \partial x_2}}{(1 + |\nabla h^\varepsilon|^2)^{\frac{3}{2}}}$$

hence

$$\left| \frac{\partial n_3}{\partial x_i} \right| \leq 2 |D_1 h^\varepsilon| \cdot |D_2 h^\varepsilon| \leq |D_2 h^\varepsilon| (1 + |D_1 h^\varepsilon|^2)$$

similarly

$$\left| \frac{\partial n_j}{\partial x_i} \right| \leq |D_2 h^\varepsilon| (1 + |D_1 h^\varepsilon|^2).$$

□

Lemma 5.

$$\int_{\Omega^\varepsilon} |\nabla u|^2 \leq \frac{4}{\nu} a(u, u) + 10 \int_{\Omega^\varepsilon} |\nabla G^\varepsilon|^2 + 4C(\Gamma_1^\varepsilon) \left\{ \int_{\Gamma_1^\varepsilon} |u|^2 + \int_{\Gamma_1^\varepsilon} |G^\varepsilon|^2 \right\}. \quad (5.2)$$

Proof. [10] We have

$$\int_{\Omega^\varepsilon} |\nabla u|^2 \leq 2 \int_{\Omega^\varepsilon} |\nabla(u - G^\varepsilon)|^2 + 2 \int_{\Omega^\varepsilon} |\nabla G^\varepsilon|^2.$$

and from 4

$$\int_{\Omega^\varepsilon} |\nabla u|^2 \leq \frac{2}{\nu} a(u - G^\varepsilon, u - G^\varepsilon) + 2C(\Gamma_1^\varepsilon) \int_{\Gamma_1^\varepsilon} |u - G^\varepsilon|^2 + 2 \int_{\Gamma_1^\varepsilon} |\nabla G^\varepsilon|^2$$

or

$$a(u - G^\varepsilon, u - G^\varepsilon) \leq 2a(u, u) + 2a(G^\varepsilon, G^\varepsilon) \leq 2a(u, u) + 4\nu \int_{\Omega^\varepsilon} |\nabla G^\varepsilon|^2$$

so (5.2) follows. \square

5.1. Scaling and uniform estimates. Let $\Omega = \Omega^1$, $y = \frac{x_3}{\varepsilon}$ and assume that

$$\hat{l} = \varepsilon l^\varepsilon, \quad \hat{k} = \varepsilon k^\varepsilon, \quad \hat{f}(x, y) = \varepsilon^2 f^\varepsilon(x, x_3).$$

Let consider also $\hat{G}(x, y) = (\hat{G}_1(x, y), \hat{G}_2(x, y), \hat{G}_3(x, y)) \in (H^1(\Omega))^3$ such that

$$\hat{G}.n = 0 \quad \text{sur} \quad \Gamma_1 \cup \omega \quad (\Gamma_1 = \Gamma_1^1)$$

$$\frac{\partial \hat{G}_1}{\partial x_1} + \frac{\partial \hat{G}_2}{\partial x_2} + \frac{\partial \hat{G}_3}{\partial y} = 0 \quad (\text{div } \hat{G} = 0)$$

we define $G^\varepsilon \in (H^1(\Omega^\varepsilon))^3$, by

$$G_i^\varepsilon(x, x_3) = \hat{G}_i(x, y), \quad i = 1, 2, \quad G_3^\varepsilon(x, x_3) = \varepsilon \hat{G}_3(x, y),$$

hence

$$\int_{\Omega^\varepsilon} |\nabla G^\varepsilon|^2 dx dx_3 \leq \frac{1}{\varepsilon} \int_{\Omega} |\nabla \hat{G}|^2 dx dy. \quad (5.3)$$

$$\int_{\Gamma_1^\varepsilon} |G^\varepsilon|^2 \leq C_0(\Omega) \int_{\Omega} (|\hat{G}|^2 + |\nabla \hat{G}|^2). \quad (5.4)$$

with $C_0(\Omega)$ independant of ε .

$$\|f^\varepsilon\|_{0, \Omega^\varepsilon}^2 = \frac{1}{\varepsilon^3} \|\hat{f}\|_{0, \Omega}^2 \quad (5.5)$$

Lemma 6. Assume that there exists $\hat{l} \in \mathbb{R}^+$ such that

$$\varepsilon l^\varepsilon = \hat{l} \quad \text{and} \quad \frac{C(\Gamma_1^\varepsilon)}{l^\varepsilon} < \frac{3}{\nu}, \quad (5.6)$$

then there exists a constante $C > 0$ independent of ε such that

$$\varepsilon \int_{\Omega^\varepsilon} |\nabla u|^2 \leq C. \quad (5.7)$$

Proof. [10] Choosing $\varphi = G^\varepsilon$ in (3.9), and using Young's inequality we obtain

$$\begin{aligned} \frac{1}{2}a(u, u) + \frac{l^\varepsilon}{2} \int_{\Gamma_1^\varepsilon} u^2 + \int_{\omega} k^\varepsilon |u - s| \leq \frac{5l^\varepsilon}{4} \int_{\Gamma_1^\varepsilon} |G^\varepsilon|^2 + \frac{33\nu}{32} \int_{\Omega^\varepsilon} |\nabla G^\varepsilon|^2 \\ + \left(\frac{32(h_M^\varepsilon)^2}{\nu} + \frac{4h_M^\varepsilon}{l^\varepsilon} \right) \int_{\Omega^\varepsilon} |f^\varepsilon|^2 + \frac{\nu}{32} \int_{\Omega^\varepsilon} |\nabla u|^2. \end{aligned} \quad (5.8)$$

From Lemma 5 we have

$$\int_{\Omega^\varepsilon} |\nabla u|^2 \leq \frac{4}{\nu} a(u, u) + 10 \int_{\Omega^\varepsilon} |\nabla G^\varepsilon|^2 + 4C(\Gamma_1^\varepsilon) \left\{ \int_{\Gamma_1^\varepsilon} |u|^2 + \int_{\Gamma_1^\varepsilon} |G^\varepsilon|^2 \right\}. \quad (5.9)$$

From (5.8) and (5.9) we get

$$\begin{aligned} \int_{\Omega^\varepsilon} |\nabla u|^2 \leq \left(\frac{1}{4} + \frac{\nu C(\Gamma_1^\varepsilon)}{4l^\varepsilon} \right) \int_{\Omega^\varepsilon} |\nabla u|^2 + 10 \left(\frac{1}{\nu} + C(\Gamma_1^\varepsilon) \right) \int_{\Omega^\varepsilon} |G^\varepsilon|^2 \\ + \left(\frac{73}{4} + \frac{33\nu C(\Gamma_1^\varepsilon)}{4l^\varepsilon} \right) \int_{\Omega^\varepsilon} |\nabla G^\varepsilon|^2 + 8 \left(\frac{1}{\nu} + \frac{C(\Gamma_1^\varepsilon)}{l^\varepsilon} \right) \left(\frac{32(h_M^\varepsilon)^2}{\nu} + \frac{4h_M^\varepsilon}{l^\varepsilon} \right) \int_{\Omega^\varepsilon} |f^\varepsilon|^2 \end{aligned}$$

from (5.3), (5.4), (5.5) and (5.6) we deduce the result. \square

Assuming (4.1) and the first of (4.2) then from (5.7) we get the same estimates (5.2) and (5.2). Other estimates follows from the "Poincaré inequality" (Lemma 3) with the new variables,

$$\int_{\Omega} |\hat{u}_i|^2 \leq 2h_M \int_{\Gamma_1} |\hat{u}_i|^2 + 2h_M^2 \int_{\Omega} \left| \frac{\partial \hat{u}_i}{\partial y} \right|^2,$$

for $i = 1, 2, 3$, hence,

$$\|\hat{u}_i\|_{L^2(\Omega)} \leq C_3, \quad \varepsilon^2 \|\hat{u}_3\|_{L^2(\Omega)}^2 \leq C_4. \quad (5.10)$$

So we obtain the same convergence as in Corollary 1. So we obtain the limit variational inequality,

$$\begin{aligned} \sum_{i=1}^2 \int_{\Omega} \frac{1}{2} \frac{\partial u_i^*}{\partial y} \frac{\partial (\hat{\varphi}_i - u_i^*)}{\partial y} - p^*(x) \frac{\partial \hat{\varphi}_i}{\partial x_i} dx dy - \sum_{i=1}^2 \int_{\omega} p^*(x) \hat{\varphi}_i(x, h) \frac{\partial h}{\partial x_i}(x) dx + \\ \sum_{i=1}^2 \hat{l} \int_{\omega} u_i^*(x, h) (\hat{\varphi}_i(x, h) - u_i^*(x, h)) dx + \int_{\omega} \hat{k} (|\hat{\varphi} - s| - |u^* - s|) dx \\ \geq \sum_{i=1}^2 \int_{\omega} \hat{f}_i (\hat{\varphi} - u_i^*) dx dy. \forall \hat{\varphi} = (\hat{\varphi}_1, \hat{\varphi}_2) \in \mathbf{\Pi}(\hat{V}_{div}), \end{aligned} \quad (5.11)$$

$$\begin{aligned} \mathbf{\Pi}(\hat{V}_{div}) &= \{ \hat{\varphi} = (\hat{\varphi}_1, \hat{\varphi}_2) \in (H^1(\Omega))^2 : \exists \hat{\varphi}_3 \in H^1(\Omega), \hat{\varphi} = (\hat{\varphi}_1, \hat{\varphi}_2, \hat{\varphi}_3) \in \hat{V}_{div} \} \\ &= \{ \hat{\varphi} = (\hat{\varphi}_1, \hat{\varphi}_2) \in (H^1(\Omega))^2 : (\hat{\varphi} - \bar{G})|_{\Gamma_L} = 0 \}. \end{aligned}$$

Lemma 7. [10]

$$F_1 = \{u^* = (u_1^*, u_2^*) \in (L^2(\Omega))^2 : \frac{\partial u_i^*}{\partial y} \in L^2(\Omega), \quad i = 1, 2, u^* \text{ satisfait } (D')\}$$

is contained in the closure of $\Pi(\hat{V}_{div})$ in the topology of $V_y \times V_y$.

Lemma 8. The convergence $\hat{u}_i \rightarrow u_i^*$ when $\varepsilon \rightarrow 0$, for $i = 1, 2$ is strong in V_y .

Proof. [10] From the ‘‘Korn inequality’’ (5.2) we have pour $\hat{\varphi} \in \hat{V}_{div}$

$$\sum_{i=1}^2 \left\| \frac{\partial(\hat{u}_i - \hat{\varphi}_i)}{\partial y} \right\|_{L^2(\Omega)}^2 \leq \hat{a}(\hat{u} - \hat{\varphi}, \hat{u} - \hat{\varphi}) + C(\Gamma_1, h) \int_{\Gamma_1} |\hat{u} - \hat{\varphi}|^2$$

and

$$\begin{aligned} \hat{a}(\hat{u} - \hat{\varphi}, \hat{u} - \hat{\varphi}) &= \hat{a}(\hat{\varphi}, \hat{\varphi} - \hat{u}) + \hat{a}(\hat{u}, \hat{u} - \hat{\varphi}) \leq \hat{a}(\hat{\varphi}, \hat{\varphi} - \hat{u}) + \hat{l} \int_{\Gamma_1} \hat{u}(\hat{\varphi} - \hat{u}) \\ &\quad + \hat{k} \int_{\omega} (|\hat{\varphi} - s| - |\hat{u} - s|) dx + \int_{\Omega} \hat{f}(\hat{\varphi} - \hat{u}) \end{aligned}$$

so

$$\begin{aligned} \sum_{i=1}^2 \left\| \frac{\partial(\hat{u}_i - \hat{\varphi}_i)}{\partial y} \right\|_{L^2(\Omega)}^2 + \hat{k} \int_{\omega} (|\bar{u} - s| - |\bar{\varphi} - s|) dx &\leq \int_{\Omega} \frac{\partial \bar{\varphi}}{\partial y} \frac{\partial(\bar{\varphi} - u^*)}{\partial y} dx dy \\ + \hat{l} \int_{\Gamma_1} u^*(\bar{\varphi} - u^*) + C(\Gamma_1, h) \int_{\Gamma_1} |u^* - \bar{\varphi}|^2 + \sum_{i=1}^2 \int_{\Omega} f_i(\bar{\varphi}_i - u_i^*). \end{aligned}$$

using Lemma 7 we can pass to the limit $\bar{\varphi}_i \rightarrow u_i^*$ in the right part to obtain the strong convergence of $\hat{u}_i \rightarrow u_i^*$ in the left part. \square

Theorem 14. The limit functions u^*, p^* satisfy

$$p^*(x_1, x_2, y) = p^*(x_1, x_2) \quad \text{a.e. in } \Omega, \quad p^* \in H^1(\omega),$$

$$-\frac{\partial^2 u_i^*}{\partial y^2} + \frac{\partial p^*}{\partial x_i} = f_i \quad (i = 1, 2) \quad \text{ds } L^2(\Omega).$$

satisfies the following weak form of the Reynolds equation

$$\int_{\omega} \left(\frac{h^3}{12} \nabla p^* - \frac{h}{2} s^* - \frac{h}{2} s_h^* + \tilde{f} \right) \nabla \varphi dx + \int_{\omega} s_h^* \nabla h \varphi dx + \int_{\partial \omega} \varphi \tilde{g}.n. = 0 \quad \forall \varphi \in H^1(\omega).$$

where

$$\begin{aligned} s^*(x) &:= u^*(x, 0), \quad s_h^*(x) := u^*(x, h(x)) \\ \tilde{g}(x) &:= \int_0^{h(x)} \hat{g}(x, y) dy \quad \forall x \in \partial \omega. \end{aligned}$$

Moreover, the traces

$$\begin{aligned} \tau^* &:= \frac{\partial u^*}{\partial y}(\cdot, 0), \quad \tau_h^* := \frac{\partial u^*}{\partial y}(\cdot, h(x)), \\ s_h^* &:= u^*(\cdot, h(x)), \quad s^* := u^*(\cdot, 0) \end{aligned}$$

satisfy the following limit form of the Tresca and Fourier boundary conditions

$$\left. \begin{aligned} |\tau^*| = \hat{k} &\implies \exists \lambda \geq 0 \quad u^* = s + \lambda \tau^* \\ |\tau^*| < \hat{k} &\implies u^* = s \end{aligned} \right\} p.p. \text{ sur } \omega.$$

$$\tau_h^* \nabla h \cdot n + \hat{l} s_h^* = 0 \text{ p.p. sur } \Gamma_1.$$

6. STUDY OF THE CASE 6

To study the asymptotic behavior of the solutions to this Case, we use the same scaling so we introduce the change of the variable $z = \frac{x_3}{\varepsilon}$ and obtain a fixed domain which is independent of ε ,

$$\Omega = \{(x', z) \in \mathbb{R}^3 : (x', 0) \in \omega, 0 < z < h(x')\},$$

$$\partial\Omega = \Gamma = \bar{\omega} \cup \bar{\Gamma}_1 \cup \bar{\Gamma}$$

then we define the following functions in Ω

$$\hat{u}_i^\varepsilon(x', z) = u_i^\varepsilon(x', x_3) \text{ for } i = 1, 2, \quad \hat{u}_3^\varepsilon(x', z) = \varepsilon^{-1} u_3^\varepsilon(x', x_3),$$

$$\hat{p}^\varepsilon(x', z) = \varepsilon^r p^\varepsilon(x', x_3); \quad \hat{T}^\varepsilon(x', z) = T^\varepsilon(x', x_3).$$

And assume the dependence of the data on ε

$$\hat{K}(x', z) = \varepsilon^{-2+r+\alpha} K^\varepsilon(x', x_3), \quad \hat{R}(x', z) = \varepsilon^{r+\alpha} R^\varepsilon(x', z), \quad \hat{\mu} = \mu^\varepsilon$$

with

$$\alpha = \frac{3(2-q)}{3-q}.$$

We suppose that

$$K_* \leq \hat{K} \leq K^*.$$

Let $\hat{G} = (\hat{G}_1, \hat{G}_2, \hat{G}_3)$ be independent on ε :

$$\operatorname{div}_z(\hat{G}) = \frac{\partial \hat{G}_1}{\partial x_1} + \frac{\partial \hat{G}_2}{\partial x_2} + \frac{\partial \hat{G}_3}{\partial z} = 0, \quad \text{and } \hat{G} = \hat{g} \text{ on } \Gamma.$$

Thus the extension G^ε of \hat{g} is defined by

$$G_i^\varepsilon(x', x_3) = \hat{G}_i(x', z) \text{ } i = 1, 2; \quad G_3^\varepsilon(x', x_3) = \varepsilon \hat{G}_3^\varepsilon.$$

Injecting the new data and unknown in (3.12)-(3.13), we deduce that $(\hat{u}^\varepsilon, \hat{p}^\varepsilon, \hat{T}^\varepsilon)$ satisfies the following problem

$$\hat{a}(\hat{T}^\varepsilon; \hat{u}^\varepsilon, \hat{\phi} - \hat{u}^\varepsilon) + (\hat{p}^\varepsilon, \operatorname{div}_z(\hat{\phi} - \hat{u}^\varepsilon)) + \hat{j}(\hat{\phi}) - \hat{j}(\hat{u}^\varepsilon) \geq \sum_{i=1}^2 (f_i, \hat{\phi}_i - \hat{u}_i^\varepsilon) + \varepsilon (f_3, \hat{\phi}_3 - \hat{u}_3^\varepsilon) \quad \forall \hat{\phi} \in V, \quad (6.1)$$

$$\int_{\Omega} \varepsilon^2 \hat{K} \nabla_\varepsilon \hat{T}^\varepsilon \nabla_\varepsilon \hat{\psi} dx' dz + \int_{\Omega} \hat{R} \hat{T}^\varepsilon \hat{\psi} = 2 \int_{\Omega} \varepsilon^\alpha \hat{\mu}(\hat{T}^\varepsilon) | \hat{D}(\hat{u}^\varepsilon) |^r \hat{\psi} dx' dz \quad \forall \hat{\psi} \in W^{1,q'}(\Omega), \quad (6.2)$$

where

$$V = \left\{ v \in (W^{1,r}(\Omega))^3 : v = \hat{G} \text{ on } \Gamma_L \cup \Gamma_1, \quad v \cdot n|_\omega = 0 \right\}$$

$$\begin{aligned} \hat{a}(\hat{T}^\varepsilon; \hat{u}^\varepsilon, \hat{\phi}) &= +\varepsilon^2 \sum_{1 \leq i, j \leq 2} \int_{\Omega} \hat{\mu}(\hat{T}^\varepsilon) | \hat{D}(\hat{u}^\varepsilon) |^{r-2} \left(\frac{\partial \hat{u}_i^\varepsilon}{\partial x_j} + \frac{\partial \hat{u}_j^\varepsilon}{\partial x_i} \right) \frac{\partial \hat{\phi}_i}{\partial x_j} + \\ &+ \sum_{i=1}^2 \int_{\Omega} \hat{\mu}(\hat{T}^\varepsilon) | \hat{D}(\hat{u}^\varepsilon) |^{r-2} \left(\frac{\partial \hat{u}_i^\varepsilon}{\partial z} + \varepsilon^2 \frac{\partial \hat{u}_3^\varepsilon}{\partial x_i} \right) \left(\frac{\partial \hat{\phi}_i}{\partial z} + \varepsilon^2 \frac{\partial \hat{\phi}_3}{\partial x_i} \right) + \\ &+ \varepsilon^2 \int_{\Omega} \hat{\mu}(\hat{T}^\varepsilon) | \hat{D}(\hat{u}^\varepsilon) |^{r-2} \frac{\partial \hat{u}_3^\varepsilon}{\partial z} \frac{\partial}{\partial z} (\hat{\phi}_3 - \hat{u}_3^\varepsilon) dx' dz, \end{aligned}$$

$$| \hat{D}(\hat{u}^\varepsilon) |^2 = \varepsilon^2 \left(\frac{1}{4} \sum_{1 \leq i, j \leq 2} \left(\frac{\partial \hat{u}_i^\varepsilon}{\partial x_j} + \frac{\partial \hat{u}_j^\varepsilon}{\partial x_i} \right)^2 + \left(\frac{\partial \hat{u}_3^\varepsilon}{\partial z} \right)^2 \right) + \frac{1}{2} \sum_{i=1}^2 \left(\frac{\partial \hat{u}_i^\varepsilon}{\partial z} + \varepsilon^2 \frac{\partial \hat{u}_3^\varepsilon}{\partial x_i} \right)^2 = \varepsilon^2 | D(u^\varepsilon) |^2$$

$$\nabla_\varepsilon v = \left(\frac{\partial v_1}{\partial x_1}, \frac{\partial v_2}{\partial x_2}, \frac{1}{\varepsilon} \frac{\partial v_3}{\partial z} \right)^t, \quad \hat{j}(\hat{\phi}) = \int_{\omega} \hat{k} | \hat{\phi} - s | dx'.$$

Let's now introduce the linear subspace

$$V_z^q = \{ v \in W^{1,q}(\Omega) : \frac{\partial v}{\partial z} \in L^q(\Omega), \quad v_{\Gamma_1} = 0 \}.$$

Theorem 15. [15] *Assume (H) hold, and $\alpha = \frac{3(2-q)}{3-q}$, there exists a constant C, independent of ε , such that*

$$\sum_{i,j=1}^2 \left\| \varepsilon \frac{\partial \hat{u}_i^\varepsilon}{\partial x_j} \right\|_{L^r(\Omega)} + \left\| \varepsilon \frac{\partial \hat{u}_3^\varepsilon}{\partial z} \right\|_{L^r(\Omega)} + \sum_{i=1}^2 \left(\left\| \frac{\partial \hat{u}_i^\varepsilon}{\partial z} \right\|_{L^r(\Omega)} + \left\| \varepsilon^2 \frac{\partial \hat{u}_3^\varepsilon}{\partial x_i} \right\|_{L^r(\Omega)} \right) \leq C \quad (6.3)$$

$$\left\| \frac{\partial \hat{p}^\varepsilon}{\partial x_i} \right\|_{W^{-1,r'}(\Omega)} \leq C, \quad i = 1, 2, \quad \left\| \frac{\partial \hat{p}^\varepsilon}{\partial z} \right\|_{W^{-1,r'}(\Omega)} \leq C \varepsilon, \quad (6.4)$$

$$\left\| \frac{\partial \hat{T}^\varepsilon}{\partial z} \right\|_{W^{1,q}(\Omega)} \leq C, \quad \sum_{i=1}^2 \left\| \varepsilon \frac{\partial \hat{T}^\varepsilon}{\partial x_i} \right\|_{W^{1,q}(\Omega)} \leq C. \quad (6.5)$$

Proof. From (6.1) and (6.2), we obtain (6.3)-(6.4), the main difficulty here is to obtain (6.5) which need the technical condition $\alpha = \frac{3(2-q)}{3-q}$ see [15] for all the proof. \square

So the following weak convergences hold

Theorem 16. [15] *Assume (H), there exist*

$$\begin{aligned} u_i^* \in V_z^r \quad i = 1, 2, \quad p^* \in L_0^{r'}(\Omega), \\ \text{and } T^* \in W^{1,q}(\Omega) \quad \text{with } T_{|\Gamma_1}^* = 0, \end{aligned}$$

such that

$$\hat{u}_i^\varepsilon \rightharpoonup u_i^* \quad (1 \leq i \leq 2) \quad \text{weakly in } V_z^r, \quad (6.6)$$

$$\varepsilon \frac{\partial \hat{u}_i^\varepsilon}{\partial x_j} \rightharpoonup 0 \quad (1 \leq i, j \leq 2) \quad \text{weakly in } L^r(\Omega), \quad (6.7)$$

$$\varepsilon \frac{\partial \hat{u}_3^\varepsilon}{\partial z} \rightharpoonup 0 \quad (1 \leq i, j \leq 2) \quad \text{in } L^r(\Omega), \quad (6.8)$$

$$\varepsilon^2 \frac{\partial \hat{u}_3^\varepsilon}{\partial x_i} \rightharpoonup 0 \quad (1 \leq i \leq 2) \quad \text{in } L^r(\Omega), \quad (6.9)$$

$$\varepsilon \hat{u}_3^\varepsilon \rightharpoonup 0 \quad (1 \leq i, j \leq 2) \quad \text{in } L^r(\Omega), \quad (6.10)$$

$$\hat{p}^\varepsilon \rightharpoonup p^* \quad \text{in } L_0^{r'}(\Omega), \quad (6.11)$$

$$\hat{T}^\varepsilon \rightharpoonup T^* \quad \text{in } V_z^q, \quad (6.12)$$

$$\varepsilon \frac{\partial \hat{T}^\varepsilon}{\partial x_i} \rightharpoonup 0 \quad \text{in } L^q(\Omega), \quad i = 1, 2. \quad (6.13)$$

Proof. Readily, from to (6.3)-(6.4) we obtain (6.6)-(6.11), while (6.12)-(6.13) follow from (6.5). \square

Then we can pass to the limit in (6.1) for $\varepsilon \rightarrow 0$ using Minty's Lemma, and in (6.2), to obtain

Theorem 17. [15] *Assume (H), and also that $\hat{K} \in C^1(\mathbb{R}) : (\hat{K})' \in L^\infty(\mathbb{R})$, then u^*, p^*, T^* satisfy*

$$\begin{aligned} \sum_{i=1}^2 \int_{\Omega} \hat{\mu}(T^*) \left(\frac{1}{2} \sum_{i=1}^2 \left(\frac{\partial u_i^*}{\partial z} \right)^2 \right)^{\frac{r-2}{2}} \frac{\partial u_i^*}{\partial z} \frac{\partial}{\partial z} (\hat{\phi}_i - u_i^*) dx' dz - \sum_{i=1}^2 (p^*, \frac{\partial}{\partial x_i} (\hat{\phi}_i - u_i^*)) \\ + \hat{j}(\hat{\phi}) - \hat{j}(u^*) \geq \sum_{i=1}^2 (\hat{f}_i, \hat{\phi}_i - u_i^*) \quad \forall \hat{\phi} \in \Pi(V), \end{aligned} \quad (6.14)$$

$$-\frac{\partial}{\partial z} \left(\hat{\mu}(T^*) \left(\frac{1}{2} \sum_{i=1}^2 \left(\frac{\partial u_i^*}{\partial z} \right)^2 \right)^{\frac{r-2}{2}} \frac{\partial u_i^*}{\partial z} \right) + \frac{\partial p^*}{\partial x_i} = \hat{f}_i, \quad i = 1, 2 \quad \text{in } L^{r'}(\Omega), \quad (6.15)$$

$$p^*(x_1, x_2, z) = p^*(x_1, x_2) \quad \text{a.e in } \Omega, \quad p^* \in W^{1,r'}(\omega), \quad (6.16)$$

$$-\frac{\partial}{\partial z} \left(\hat{K} \frac{\partial T^*}{\partial z} \right) + \hat{R} T^* = 0 \quad \text{in } L^q(\Omega). \quad (6.17)$$

$$T^* = 0 \quad \text{on } \Gamma_1, \quad \text{and} \quad -\hat{K} \frac{\partial T^*}{\partial z} = \hat{b} \quad \text{on } \omega. \quad (6.18)$$

Then we obtain the limit problem

Theorem 18. [15] *Under the same hypothesis as in theorem 4, the traces*

$$s^* = u^*(x', 0), \quad \zeta^* = T^*(x', 0)$$

$$\tau^* = \left(\frac{1}{2} \sum_{i=1}^2 \left(\frac{\partial u_i^*}{\partial z}(x', 0) \right)^2 \right)^{\frac{r-2}{2}} \frac{\partial u_i^*}{\partial z}(x', 0),$$

satisfy the following inequality

$$\int_{\omega} \hat{k} (|\psi + s^* - s| - |s^* - s|) dx' - \int_{\omega} \hat{\mu}(\zeta^*) \tau^* \psi dx' \geq 0 \quad \forall \psi \in (L^r(\omega))^2, \quad (6.19)$$

the limit of Tresca's boundary condition on ω gives:

$$\left. \begin{aligned} \hat{\mu}(\zeta^*) |\tau^*| < \hat{k} &\Rightarrow s^* = s \\ \hat{\mu}(\zeta^*) |\tau^*| = \hat{k} &\Rightarrow \exists \lambda \geq 0 : s^* = s + \lambda \tau^* \end{aligned} \right\} \text{a.e on } \omega.$$

Also u^*, p^* and T^* satisfy the specific weak Reynolds equation

$$\begin{aligned} & \int_{\omega} \left[\int_0^h \int_0^y \hat{\mu}(T^*(x', \xi)) A^*(x', \xi) \frac{\partial u^*}{\partial \xi}(x', \xi) d\xi dy \right] \cdot \nabla \phi(x') dx' \\ & - \int_{\omega} \frac{h}{2} \left[\int_0^h \hat{\mu}(T^*(x', \xi)) A^*(x', \xi) \frac{\partial u^*}{\partial \xi}(x', \xi) d\xi \right] \cdot \nabla \phi(x') dx' \\ & + \int_{\omega} \left[\frac{h^3}{12} \nabla p^*(x') + \tilde{F}(x') \right] \cdot \nabla \phi(x') dx' = 0 \quad \forall \phi \in W^{1,r}(\omega) \end{aligned} \tag{6.20}$$

and the following equation

$$\int_{\Omega} \hat{K} \frac{\partial T^*}{\partial z} \frac{\partial \hat{\psi}}{\partial z} dx' dz + \int_{\Omega} \hat{R} T^* \hat{\psi} dx' dz = 0 \quad \forall \hat{\psi} \in W_{\Gamma_1 \cup \Gamma_L}^{1,q'}(\Omega), \tag{6.21}$$

where

$$A^*(x', \xi) = \left(\frac{1}{2} \sum_{i=1}^2 \left(\frac{\partial u^*}{\partial \xi}(x', \xi) \right)^2 \right)^{\frac{r-2}{2}}$$

$$\tilde{F}(x') = \int_0^h \int_0^y \int_0^{\xi} \hat{f}(x', y) dy d\xi dz - \frac{h}{2} \int_0^h \int_0^{\xi} \hat{f}(x', y) dy d\xi. \tag{6.22}$$

Theorem 19. *The solution (u^*, T^*, p^*) of our limit problem is unique.*

Proof. [15] Let $(U^1, T^1, p^1), (U^2, T^2, p^2)$ be two solutions of the limit problem. Then

$$T = T^1 - T^2$$

satisfies the problem

$$-\frac{\partial}{\partial z} \left(\hat{K} \frac{\partial T}{\partial z} \right) + \hat{R} T = 0, \quad T|_{\Gamma_1} = 0, \quad \hat{K} \frac{\partial T}{\partial z} = 0 \quad \text{on } \omega$$

so $T = 0$, thus $T^1 = T^2 = T^*$.

Taking $\phi = U^2$ and $\phi = U^1$ respectively, as test functions in (6.14) we get

$$\begin{aligned} & \sum_{i=1}^2 \int_{\Omega} \hat{\mu}(T^*) \left[\left(\frac{1}{2} \sum_{i=1}^2 \left(\frac{\partial U_i^1}{\partial z} \right)^2 \right)^{\frac{r-2}{2}} \frac{\partial U_i^1}{\partial z} - \left(\frac{1}{2} \sum_{i=1}^2 \left(\frac{\partial U_i^2}{\partial z} \right)^2 \right)^{\frac{r-2}{2}} \frac{\partial U_i^2}{\partial z} \right] \times \\ & \times \frac{\partial}{\partial z} (U_i^1 - U_i^2) dx' dz \leq 0, \end{aligned} \tag{6.23}$$

using some inequality [45]) we obtain for $r \geq 1$

$$\left\| \frac{\partial}{\partial z} (U^1 - U^2) \right\|_{(L^r(\Omega))^2} = 0,$$

using the Poincare inequality we deduce that

$$\| U^1 - U^2 \|_{V_z} = 0,$$

so u^* is unique. The uniqueness of p^* in $L_0^{r'}(\omega) \cap W^{1,r'}(\omega)$ follows then from the *specific weak Reynolds equation* (6.20), indeed we obtain first

$$\int_{\omega} \frac{h^3}{12} \nabla(p^1 - p^2) \nabla \phi \, dx' = 0,$$

taking

$$\phi = p^1 - p^2$$

and by Poincaré's inequality we get

$$\|p^1 - p^2\|_{L^{r'}(\omega)} = 0.$$

This ends the proof of the uniqueness. □

REFERENCES

- [1] R.A. Adams, Sobolev Spaces, Academic Press, NewYork, 1975.
- [2] G. ALLAIRE, *Homogenization of the Navier Stokes equations with a slip boundary condition*, Comm. on Pure and Applied Maths, XLIV, (1991), 605-641.
- [3] C. AMROUCHE, V. GIRAULT, *Decomposition of vector spaces and application to the Stokes problem in arbitrary dimension*. Czechoslovak Mathematical Journal, 44, (1994).
- [4] A. ASSEMIEN, G. BAYADA AND M. CHAMBAT, *Inertial effects in the asymptotic behaviour of a thin film flow*. Asymptotic Analysis No.9 , (1994), 117-208.
- [5] G. BAYADA, K. LHALOUANI, *Asymptotic and numerical analysis for unilateral contact problem with Coulomb's friction between an elastic body and a thin elastic soft layer*, Asymptotic Analysis 25 (2001) 329-362.
- [6] G.BAYADA, M. BOUKROUCHE, *On a free boundary problem for Reynolds equation derived from the Stokes system with Tresca boundary conditions*. Journal of Mathematical Analysis and Applications, Volume 282, Issue 1, 1 June 2003, Pages 212-231
- [7] G. BAYADA AND M. CHAMBAT, *The transition between the Stokes equation and the Reynolds equation. A mathematical proof*. Applied Maths. and Optim., No.14, (1986), 73-93.
- [8] JA. BELLO, *L^r -regularity for the Stokes and Navier-Stokes problems*. Annali di Matematica Pura ed Applicata, Series IV 170, (1996), 187-206.
- [9] M. BOUKROUCHE, G. LUKASZEWICZ, *Asymptotic behavior of a thin film lubrication with Coulomb fluid solid interface law*. International Journal of Engineering Science 41, (2003) 521-537.
- [10] M. BOUKROUCHE, G. LUKASZEWICZ, *On a lubrication problem with Fourier and Tresca boundary conditions*. Mathematical Models and Methods in Applied Sciences (M3AS) Vol.4, N6, (2004) pages 913-941.
- [11] M. BOUKROUCHE, R. EL MIR, *On the Navier-Stokes system in a thin film flow with Tresca free boundary condition and its asymptotic behavior*. Bull. Math. Soc. Sc. Math. Roumanie Tome 48 (96) No. 2, (2005), 139-163.
- [12] M. BOUKROUCHE, F. BOUGHANIM, H. SMAOUI, *Asymptotic behavior of a non-Newtonian flow with stick-slip condition*. Electron.J.Differ.Equ.Conf., 11 (2004).
- [13] M. BOUKROUCHE, R. EL MIR, *Asymptotic analysis of a nonNewtonian fluid in a thin domain with Tresca law* Nonlinear Analysis, Theory Methods and Applications, Volume 59, Issues 1-2, October 2004, Pages 85-105.
- [14] M. BOUKROUCHE, F. SAIDI, *Non-isothermal lubrication problem with Tresca fluid-solid interface law. Partie I*. Nonlinear Anal. Real World Applications. Volume 7, Issue 5, December 2006, Pages 1145-1166.
- [15] M. BOUKROUCHE, R. EL MIR, *Nonisothermal, non-Newtonian lubrication problem with Tresca fluid-solid law. Existence and asymptotic of weak solutions*. Nonlinear Anal. Real World Applications. Volume 9, Issue 2, April 2008, Pages 674-692.
- [16] M. BOUKROUCHE, F. SAIDI, *Non-isothermal lubrication problem with Tresca fluid-solid interface law. Partie II: Asymptotic behaviour of weak solutions*. Nonlinear Anal. Real World Applications. Volume 9, Issue 4, September 2008, Pages 1680-1701.

- [17] M. BOUKROUCHE, I. CIUPERCA, *Asymptotic behaviour of solutions of lubrication problem in a thin domain with a rough boundary and Tresca fluid-solid interface law*. Quart. Appl. Math. 64 (2006), 561-591.
- [18] H. BREZIS, *Equations et inéquations non linéaires dans les espaces vectoriels en dualité*, Annales de l'institut Fourier, 18, (1968), 115-175.
- [19] D. BRESH, *Sur les fluides incompressibles. Écoulement en eau peu profonde estimation d'énergie et fluide non newtonien*. Thèse Université Clermont-Ferrand Math.Appli., (1997).
- [20] D. BRESH, J. LEMOINE AND J. SIMON, *Écoulement engendré par le vent et la force de Cariolis dans un domaine mince : I Cas stationnaire*. C.R.A.S. Paris, t. 325, Série I, (1997), 807-812.
- [21] L. CATTABRIGA, *Su un problema al contorno relativo al sistema di equazioni di Stokes*. Rendiconti del Seminario Matematico della Università di Padova **31**, (1961), 1-33.
- [22] M.C. CATTANEO, *Sulla Conduzione de Calor*, Atti del Seminario Matematico e Fisico Della Università di Modena, 83-101, 1948.
- [23] L. CONSIGLIERI, *Stationary Solutions for a Bingham Flow with Nonlocal Friction*, Pitman Research Notes in Mathematics Series 274, (1992), 237-243.
- [24] G. DUVAUT, *Équilibre d'un solide élastique avec contact unilatéral et frottement de Coulomb*, C.R.Acad.Sc.Paris, t 290, (1980), 263-265.
- [25] G. DUVAUT AND J.L. LIONS, *Les Inéquations en mécanique et en physique*, Dunod, Paris, (1972).
- [26] U. EISELE, *Introduction to polymer physics*. Springer-Verlag, 1990.
- [27] A.C. ERINGEN, *Theory of micropolar fluids*, J. Math. Mech. 16, num 1 (1966), 1-16.
- [28] I. EKELAND AND R. TEMAM, *Analyse convexe et problèmes variationnels*, Dunod et Gauthier-Villars, Paris, (1974).
- [29] H. FUJITA, *A mathematical analysis of motions of viscous incompressible fluid under leak or slip boundary conditions*. Mathematical fluid mechanics and modeling, (Kyoto, 1994), *Sūrikaiseikikenkyūsho Kōkyūroku*, **888**, (1994), 199-216.
- [30] H. FUJITA, H. KAWARADA, *Variational inequalities for the Stokes equation with boundary conditions of friction type*. Recent developments in domain decomposition methods and flow problems (Kyoto, 1996; Anacapri, 1996), *GAKUTO International Series. Mathematical Sciences and Applications*, **11**, (1998), 15-33.
- [31] H. FUJITA, *Remarks on the Stokes flow under slip and leak boundary conditions of friction type*. Topics in mathematical fluid mechanics, Quaderni di Matematica, Dept. Math., Seconda Univ. Napoli, Caserta, **10**, (2002), 73-94.
- [32] H. FUJITA, *A coherent analysis of Stokes flows under boundary conditions of friction type*. Journal of Computational and Applied Mathematics, **149**(1), (2002), 57-69.
- [33] V. GIRAULT AND P.A. RAVIART, *Finite element Approximation of the Navier-Stokes Equations*. Springer-Verlag, (1979).
- [34] J.M. GHIDAGLIA, *Régularités solutions de certains problèmes aux limites linéaires liés aux équations d'Euler*. Communications in Partial Differential Equations. **9**(13),(1984), 1265-1298.
- [35] M. FANG, R.P. GILBERT, *Non-isothermal, non-Newtonian Hele-Schaw flows within Cattaneo's heat flux law*. Mathematical and Computer Modelling 46 (2007) 765-775.
- [36] B.O.JACOBSON, *At the boundary between lubrication and wear*. First world tribology conference, London, (1997), 291-298.
- [37] B.O. JACOBSON, B.J. HAMROCK, *Non Newtonian fluid model incorporated into elastohydrodynamic lubrication of rectangular contacts*. Journal of Tribology, 106, (1984), 275-284.
- [38] O.A. LADYZHENSKAYA, *The mathematical theory of viscous incompressible flow*. Gordon and Breach, Science Publishers: New York-London-Paris, 1969.
- [39] L. LANDAU, E. LIFCHITZ, *Physique théorique et mécanique des fluides*. Vol.6, 2ième edition, 1989.
- [40] G. ŁUKASZEWICZ, *Micropolar fluids, Theory and applications*, Birkhäuser 1999.
- [41] A. MAJDA, *Compressible Fluid flow and Systems of conservation laws in several space variables*. Applied Mathematical Sciences 53, Springer-Verlag 1984.
- [42] M. MOUSSAOUI M., A.M. ZINE, *Existence and regularity results for the Stokes system with non-smooth boundary data in a polygon*. Mathematical Models and Methods in applied Sciences, **8**,(1998), 1307-1315.
- [43] J. NEČAS, J. JARUSEK AND J. HASLINGER, *On the solution of the variational inequality to the signorini problem with small friction*, Boll. Un. Mat. Ital. B(5) 17 (1980), 796-811.
- [44] J. NEČAS, *Les Méthodes directes en théorie des équations elliptiques*. Masson et Cie: Paris, 1967.

- [45] J.T. ODEN, *Qualitative Methods in Non-linear Mechanics*, Prentice-Hall, Englewood Cliffs, NJ, 1986.
- [46] R. PIT, *Mesure locale de la vitesse à l'interface solide-liquide simple: Glissement et rôle des interactions*. Thèse Physique Université Paris XI, 1999.
- [47] R. PIT, H. HERVET, L. LÉGER, *Direct experimental evidences for flow with slip at hexadecane solid interfaces*. *La revue de Métallurgie-CIT/Science*, February (2001).
- [48] N. SAITO, *On the Stokes equations with the leak and slip boundary conditions of friction type: regularity solutions*. Publications of RIMS, Kyoto University, **40**(2004), 345–383.
- [49] J. SERRIN, *Mathematical principles of classical fluid mechanics*. In *Handbuch der Physik*, **8**(1), (1959), 125–263.
- [50] B.O. JACOBSON AND B.J. HAMROCK, *Non newtonian fluid model incorporated into elastohydrodynamic lubrication of rectangular contacts*, *J. of Tribology*, 106, (1984), 275-284,
- [51] J. SHIEH, B.J.HAMROCK, *Film collapse in ehl and micro-ehl*. *Journal of Tribology*, **113**, (1991), 372–377.
- [52] A. STROZZI, *Formulation of of three lubrication problems in term of complementarity*. *Wear*, 104, (1985), 103-119.
- [53] D. A. TARZIA, *Una familia de problemas que converge hacia el caso estacionario del problema de Stefan a dos fases*. *Mathematicae Notae* Vol. 27, (1979) 157-165.
- [54] R. TEMAM, *Navier-Stokes equations. Theory and Numerical Analysis*. North-Holland, Amsterdam, New York, Oxford (1979).
- [55] J.L. TEVAARWERK, *The shear of hydrodynamic oil films*. *Phd Thesis: Cambridge, England*, 1976.

MAT

INDICE GENERAL

Serie A: CONFERENCIAS, SEMINARIOS Y TRABAJOS DE MATEMÁTICA

ISSN 1515-4904

1 (2000): Elvira Mascolo – Francesco Siepe, “Functionals of the Calculus of Variations with non Standard Growth Conditions”.

2 (2000): Domingo A. Tarzia, “A Bibliography on Moving-Free Boundary Problems for the Heat-Diffusion Equation. The Stefan and Related Problems”.

3 (2001): Domingo A. Tarzia (Ed.), “VI Seminario sobre Problemas de Frontera Libre y sus Aplicaciones”, Primera Parte:

- Ma. Cristina Sanziel, “Conditions to obtain a waiting time for a discrete two-phase Stefan problem”, 1-6.
- Ariel L. Lombardi – Domingo A. Tarzia, “On similarity solutions for thawing processes”, 7-12.
- Ricardo Weder, “Direct and inverse scattering for the nonlinear Schrödinger equation with a potential”, 13-20.
- Domingo A. Tarzia, “Stefan problem for a non-classical heat equation”, 21-26.
- Pedro Morin – Rubén D. Spies, “A quasilinearization approach for parameter identification in nonlinear abstract Cauchy problems”, 27-41.

4 (2001): Domingo A. Tarzia (Ed.), “VI Seminario sobre Problemas de Frontera Libre y sus Aplicaciones”, Segunda Parte:

- Omar Gil, “El problema de Hele-Shaw como un problema límite para la ecuación de los medios porosos”, 1-10.
- Juan C. Reginato – Domingo A. Tarzia, “Estimations of nutrient uptakes by roots of crops through a moving boundary model”, 11-16.
- Oscar D. Quiroga – Luis T. Villa – Fernando Suarez, “Problemas de frontera libre en procesos de transferencia de materia y energía con reacción química”, 17-22.
- Edgardo A. Spiazzi – Rodolfo H. Mascheroni, “Modelo de deshidratación osmótica de alimentos vegetales”, 23-32.
- Eduardo A. Santillan Marcus – Domingo A. Tarzia, “Exact solutions for phase change processes in humid porous half spaces”, 33-38.

5 (2001): Domingo A. Tarzia (Ed.), “VI Seminario sobre Problemas de Frontera Libre y sus Aplicaciones”, Tercera Parte:

- Adriana C. Briozzo – Domingo A. Tarzia, “On a two-phase Stefan problem with nonlinear thermal coefficients”, 1-10.
- Germán Torres – Cristina V. Turner, “Métodos de diferencias finitas para un problema de Bingham unidimensional”, 11-26.
- Analía Gastón – Gustavo Sánchez Sarmiento – Horacio Reggiardo, “Un problemas de frontera libre: Fusión de una vaina de acero dentro de una cuchara de acería”, 27-32.
- Ma. Fernanda Natale – Domingo A. Tarzia, “An exact solution for a one-phase Stefan problem with nonlinear thermal coefficient”, 33-36.
- Claudia Lederman – Juan L. Vazquez – Noemí Wolanski, “Uniqueness of solution to a free boundary problem from combustion with transport”, 37-41.

6 (2002): Federico Talamucci, “Some Problems Concerning with Mass and Heat Transfer in a Multi-Component System”.

7 (2004): Domingo A. Tarzia (Ed.), “Primeras Jornadas sobre Ecuaciones Diferenciales, Optimización y Análisis Numérico”, Primera Parte:

- Adriana B. Verdiell – María C. Maciel – Susana L. Orofino – Tatiana I. Gibelli, “A survey of the spectral gradient method”, 1-14.
- María F. Natale – Domingo A. Tarzia, “An integral equation in order to solve a one-phase Stefan problem with nonlinear thermal conductivity”, 15-24.
- María C. Sanziel – Domingo A. Tarzia, “Optimization on the heat flux in a mixed elliptic problem with temperature constraints”, 25-30.
- Claudia M. Gariboldi – Domingo A. Tarzia, “A new proof of the convergence of distributed optimal controls on the internal energy in mixed elliptic problems”, 31-42.

8 (2004): Domingo A. Tarzia (Ed.), “Primeras Jornadas sobre Ecuaciones Diferenciales, Optimización y Análisis Numérico”, Segunda Parte:

- Rubén D. Spies, “Differentiability of the solutions of a semilinear abstract Cauchy problem with respect to parameters”, 1-10.
- Adriana C. Briozzo – María F. Natale – Domingo A. Tarzia, “An explicit solution for a two-phase Stefan problem with a similarity exponential heat sources”, 11-19.
- Domingo A. Tarzia, “An explicit solution for a two-phase unidimensional Stefan problem with a convective boundary condition at the fixed face”, 21-27.

9 (2005): Micol Amar – Roberto Gianni, “A Brief Survey on Homogenization with a Physical Application”.

10 (2005): Domingo A. Tarzia – Cristina V. Turner (Eds.), “Segundas Jornadas sobre Ecuaciones Diferenciales, Optimización y Análisis Numérico”:

- Marcos Gaudiano – Cristina Turner, “Difusión de un solvente en un polímero vídrioso con una condición de contorno del tipo creciente en el tiempo”, 1-9.
- Adriana C. Briozzo – María F. Natale – Domingo A. Tarzia, “A one-phase Lamé-Clapeyron-Stefan problem with nonlinear thermal coefficients”, 11-16.
- Eduardo A. Santillan Marcus - Domingo A. Tarzia, “Un caso de determinación de coeficientes térmicos desconocidos de un material semiinfinito poroso a través de un problema de desublimación con acoplamiento de temperatura y humedad”, 17-22.

11 (2005): Antonio Fasano, “Mathematical Models of Some Diffusive Processes with Free Boundaries”.

12 (2006): Arturo De Pablo, “An Introduction to the Problem of Blow-up for Semilinear and Quasilinear Parabolic Equations”.

13 (2006): Graciela G. Garguichevich – Claudia M. Gariboldi – Pedro R. Marangunic – Diego Pallara, “Direct methods in the calculus of variations”.

14 (2007): María C. Maciel – Domingo A. Tarzia (Eds.), “Terceras Jornadas sobre Ecuaciones Diferenciales, Optimización y Análisis Numérico”:

- Tatiana I. Gibelli – María C. Maciel, “Large-scale algorithms for minimizing a linear function with a strictly convex quadratic constraint”, 1-12.
- María C. Maciel – Elvio A. Pilotta – Graciela N. Sottosanto, “Thickness optimization of an elastic beam”, 13-23.
- María F. Natale – Eduardo A. Santillan Marcus – Domingo A. Tarzia, “Determinación de dos coeficientes térmicos a través de un problema de desublimación con acoplamiento de temperatura y humedad”, 25-30.
- Rubén D. Spies – Karina G. Temperini, “Sobre la no convergencia del método de mínimos cuadrados en dimension infinita”, 31-34.
- Juan C. Reginato – Domingo A. Tarzia, “An alternative method to compute Michaelis-Menten parameters from nutrient uptake data”, 35-40.

15 (2008): D.A. Tarzia – R.H. Mascheroni (Eds.), “Workshop on Mathematical Modelling of Energy and Mass Transfer Processes, and Applications”:

- María F. Natale – Domingo A. Tarzia, “The classical one-phase Stefan problem with temperature-dependent thermal conductivity and a convective term”, 1-16.
- Carmen Cortazar – Manuel Elgueta, “Non-local diffusion”, 17-24.
- Luis T. Villa, “Consideraciones sobre el buen planteo de un modelo de frontera libre-móvil descriptivo de un proceso de freído por inmersión”, 25-36.
- Ricardo Simpson – Isabel Figueroa – Arthur Teixeira, “Simple, practical, and efficient on-line correction of process deviations in batch retort through simulation”, 37-53.

16 (2009): Mahdi Boukrouche, “A brief survey on lubrication problems with nonlinear boundary conditions”.

Serie B: CURSOS Y SEMINARIOS PARA EDUCACIÓN MATEMÁTICA

ISSN 1515-4912

1 (2000): Domingo A. Tarzia, “Cómo pensar, entender, razonar, demostrar y crear en Matemática”.

2 (2003): Domingo A. Tarzia, “Matemática: Operaciones numéricas y geometría del plano”

INFORMACION PARA LOS AUTORES

Los trabajos han de estar escritos en español o inglés. Excepcionalmente el Director y el Comité Editorial podrán admitir trabajos escritos en otros idiomas ampliamente utilizados. Deberá presentarse el texto mecanografiado o elaborado mediante un procesador de textos, con caracteres de 12 puntos, en un rectángulo de 16cm×24cm y en una sola cara del papel. Trabajos escritos en LATEX o en MS-WORD serán bienvenidos y en ese caso el autor deberá adjuntar un diskette con los archivos correspondientes, o bien enviarlos por correo electrónico.

En cada trabajo deberá constar, en la primera página, a partir de la quinta línea, el título en letras mayúsculas y sin punto final, el nombre del o de los autores, su identificación institucional y su correspondiente dirección postal y electrónica. Se acompañará un resumen que no exceda las 200 palabras en español y otro en inglés, añadiendo en ambos las palabras claves. También se solicita la inclusión de la correspondiente AMS-Mathematics Subject Classification. Al final de la última página deberán dejarse al menos dos líneas en blanco para incluir los datos de recepción. Las tablas y gráficos deberán insertarse en el texto y estar numeradas en forma correlativa.

Las referencias bibliográficas se compondrán sólo de los trabajos mencionados en el texto y se incluirán al final, por orden alfabético de autores y en orden cronológico, si existieran varios trabajos del mismo autor; cada una precedida por el correspondiente número de orden, entre corchetes. Las citas en el texto se efectuarán según los siguientes modelos: [1]; Caffarelli & Vazquez [1]; Caffarelli & Vazquez (1995, [1]). Y en la referencia final:

[1] CAFFARELLI L. A. & VAZQUEZ J.L., *A free-boundary problem for the heat equation arising inflame propagation*, Trans. Amer. Math. Soc., 347 (1995), pp. 411-441.

[2] FASANO A. & PRIMICERIO M., *Blow-up and regularization for the Hele-Shaw problem*, in *Variational and free boundary problems*, Friedman A. & Spruck J. (Eds.), IMA Math. Appl. Vol. 53, Springer Verlag, New York (1993), pp. 73-85.

[3] RODRIGUES J. F., *Obstacle problems in mathematical physics*, North-Holland Mathematics Studies N. 134, North-Holland, Amsterdam (1987).

INTERCAMBIOS

Departamento de Matemática – Biblioteca, Servicio de Canje
Facultad de Ciencias Empresariales -Universidad Austral
Paraguay 1950, S2000FZF ROSARIO, ARGENTINA

NUMEROS APARECIDOS

Serie A:

- #1 (2000): E. Mascolo – F. Siepe, “Functionals of the Calculus of Variations with non Standard Growth Conditions”.
- #2 (2000): D.A. Tarzia, “A Bibliography on Moving-Free Boundary Problems for the Heat-Diffusion Equation. The Stefan and Related Problems”.
- # 3 (2001): D.A. Tarzia (Ed.), “VI Seminario sobre Problemas de Frontera Libre y sus Aplicaciones”, Primera Parte.
- #4 (2001): D.A. Tarzia (Ed.), “VI Seminario sobre Problemas de Frontera Libre y sus Aplicaciones”, Segunda Parte.
- #5 (2001): D.A. Tarzia (Ed.), “VI Seminario sobre Problemas de Frontera Libre y sus Aplicaciones”, Tercera Parte.
- #6 (2002): F. Talamucci, “Some Problems Concerning with Mass and Heat Transfer in a Multi-Component System”.
- #7 (2004): D.A. Tarzia (Ed.), “Primeras Jornadas sobre Ecuaciones Diferenciales, Optimización y Análisis Numérico”, Primera Parte.
- #8 (2004): D.A. Tarzia (Ed.), “Primeras Jornadas sobre Ecuaciones Diferenciales, Optimización y Análisis Numérico”, Segunda Parte.
- #9 (2005): M. Amar – R. Gianni, “A Brief Survey on Homogenization with a Physical Application”.
- #10 (2005): D.A. Tarzia – C.V. Turner (Eds.), “Segundas Jornadas sobre Ecuaciones Diferenciales, Optimización y Análisis Numérico”.
- #11 (2005): A. Fasano, “Mathematical Models of Some Diffusive Processes with Free Boundaries”.
- #12 (2006): A. De Pablo, “An Introduction to the Problem of Blow-up for Semilinear and Quasilinear Parabolic Equations”.
- #13 (2006): G.G. Garguichevich – C.M. Gariboldi – P.R. Marangunic – D. Pallara, “Direct methods in the calculus of variations”.
- #14 (2007): M.C. Maciel – D.A. Tarzia (Eds.), “Terceras Jornadas sobre Ecuaciones Diferenciales, Optimización y Análisis Numérico”.
- #15 (2008): D.A. Tarzia – R.H. Mascheroni (Eds.), “Workshop on Mathematical Modelling of Energy and Mass Transfer Processes, and Applications”.
- #16 (2009): M. Boukrouche, “A brief survey on lubrication problems with nonlinear boundary conditions”.

Serie B:

- #1(2000): D.A. Tarzia, “Cómo pensar, entender, razonar, demostrar y crear en Matemática”.
- #2(2003): D.A. Tarzia, “Matemática: Operaciones numéricas y geometría del plano”.

