

# MAT

Serie  A

Conferencias, seminarios  
y trabajos de Matemática

ISSN: 1515-4904

 8

*Primeras Jornadas  
sobre Ecuaciones  
Diferenciales,  
Optimización y  
Análisis Numérico*

*Segunda Parte*

Departamento  
de Matemática,  
Rosario,  
Argentina  
2004

UNIVERSIDAD AUSTRAL

FACULTAD DE CIENCIAS EMPRESARIALES



# MAT

## SERIE A: CONFERENCIAS, SEMINARIOS Y TRABAJOS DE MATEMÁTICA

No. 8

### PRIMERAS JORNADAS SOBRE ECUACIONES DIFERENCIALES, OPTIMIZACIÓN Y ANÁLISIS NUMÉRICO

Segunda Parte

**Domingo A. Tarzia (Ed.)**

#### INDICE

**Ruben D. Spies**, “Differentiability of the solutions of a semilinear abstract Cauchy problem with respect to parameters”, 1-10.

**Adriana C. Briozzo – María F. Natale – Domingo A. Tarzia**, “An explicit solution for a two-phase Stefan problem with a similarity exponential heat sources”, 11-19.

**Domingo A. Tarzia**, “An explicit solution for a two-phase unidimensional Stefan problem with a convective boundary condition at the fixed face”, 21-27.

**Rosario, Octubre 2004**

**DIFFERENTIABILITY OF THE SOLUTIONS OF A SEMILINEAR  
ABSTRACT CAUCHY PROBLEM WITH RESPECT TO  
PARAMETERS**

RUBEN D. SPIES<sup>†,\*</sup>

INSTITUTO DE MATEMÁTICA APLICADA DEL LITORAL, IMAL-CONICET  
AND  
DEPARTAMENTO DE MATEMÁTICA, FACULTAD DE INGENIERÍA QUÍMICA  
UNIVERSIDAD NACIONAL DEL LITORAL, SANTA FE, ARGENTINA

ABSTRACT. The Fréchet differentiability with respect to a parameter  $q$  of the solutions  $z(t; q)$  of Cauchy problems of the form  $\frac{d}{dt}z(t) = A(q)z(t) + F(q, t, z(t))$  is analyzed. Sufficient conditions on the operator  $A(q)$  and on  $F$  are derived and the corresponding sensitivity equations for the Fréchet derivative  $D_q z(t; q)$  are found.

1. INTRODUCTION

We consider the problem of continuous dependence and differentiability with respect to a parameter  $q$  of the solutions  $z(t; q)$  of the semilinear abstract Cauchy problem

$$(\mathcal{P})_q \begin{cases} \frac{d}{dt}z(t) = A(q)z(t) + F(q, t, z(t)) & z(t) \in Z, \\ z(0) = z_0 & t \in [0, T] \end{cases}$$

where  $Z$  is a Banach space,  $q \in Q_{ad} \subset Q$ , a normed linear space ( $Q_{ad}$  is an open subset of  $Q$ ), and  $A(q)$  is the infinitesimal generator of an analytic semigroup  $T(t; q)$  on  $Z$  for all  $q \in Q_{ad}$ .  $Z$  and  $Q$  are the state space and the parameter space, respectively, while  $Q_{ad}$  is called the admissible parameter set.

Identification problems associated to system  $(\mathcal{P})_q$  and other similar type of equations ([2], [5], [7]) are usually solved by direct methods such as quasilinearization. These methods require that solutions be differentiable with respect to the parameter  $q$ . In addition, their numerical implementation require an approximation to the corresponding Fréchet derivative.

Problems of the type  $\frac{d}{dt}z(t) = A(q)z(t) + u(t)$ , where  $A(q)$  generates a strongly continuous semigroup and  $A(q) = A + B(q)$  where  $B(q)$  is assumed to be bounded were studied by Clark and Gibson ([4]), Brewer ([1]). Burns et al ([3]) studied problems of the type  $\frac{d}{dt}z(t) = Az(t) + F(q, t, z(t))$ . The parameter  $q$  here did not appear in the linear part of the equation.

Here, we prove that, under certain conditions, the solutions of the general abstract Cauchy problem  $(\mathcal{P})_q$  are Fréchet differentiable with respect to  $q$  and we find the corresponding sensitivity equations.

---

*Key words and phrases.* Abstract Cauchy problem, analytic semigroup, infinitesimal generator, Fréchet differentiability, Fréchet derivative.

\* E-mail: rspies@imalpde.ceride.gov.ar

† Supported in part by CONICET, Consejo Nacional de Investigaciones Científicas y Técnicas, through projects PIP 02823 and PEI 6181, by UNL, Universidad Nacional del Litoral through project CAI+D 2002 PE 222, and by Fundación Antorchas of Argentina.

## 2. PRELIMINARY RESULTS

The following standing hypotheses are considered:

**H1:** There exist  $\varepsilon_0 > 0$  such that the type of  $T(t; q)$ , call it  $w_q$ , is less than or equal to  $-\varepsilon_0$  for all  $q \in Q_{ad}$  and there exists  $C_q > 0$  such that  $\|T(t; q)\| \leq C_q e^{-\varepsilon_0 t}$  for all  $t \geq 0$  and  $q \in Q_{ad}$ . The constant  $C_q$  depends on  $q$  but it can be chosen independent of  $q$  on compact subsets of  $Q_{ad}$ .

**H2:**  $\mathcal{D}(A(q)) = D$  is independent of  $q$  and  $D$  is a dense subspace of  $Z$ .

We shall denote by  $Z_\delta$  the space  $D((-A(q))^\delta)$  imbedded with the norm of the graph of  $-A(q)^\delta$ . Since  $0 \in \rho(A(q))$  it follows that this norm is equivalent to  $\|z\|_{q,\delta} \doteq \|(-A(q))^\delta z\|$ . Also, there exists a constant  $M_q$  such that  $\|(-A(q))^\delta T(t; q)\| \leq M_q \frac{e^{-\varepsilon_0 t}}{t^\delta}$ , for all  $t > 0$  (see [13], Theorem 2.6.13).

**H3:** There exists  $\delta \in (0, 1)$  such that

$$\|F(q, t_1, z_1) - F(q, t_2, z_2)\|_Z \leq L(|t_1 - t_2| + \|z_1 - z_2\|_{q,\delta})$$

for  $(t_i, z_i) \in U$ , where  $L$  can be chosen independent of  $q$  on any compact subset of  $Q_{ad}$ .

This last regularity condition guarantees existence and uniqueness of solutions of problem  $(\mathcal{P})_q$ , provided that the initial condition  $z_0$  is in  $Z_\delta$ . See [12] and [11] for details.

The next results can be easily proved by using the Closed Graph Theorem.

LEMMA 1: Under hypotheses H1 and H2, for any  $q_1, q_2 \in Q_{ad}$  and  $\delta \in (0, 1)$  we have:

i)  $A(q_1)(-A(q_2))^{-\delta}$  is bounded on  $Z_{1-\delta}$ .

ii)  $A(q_1)T(\cdot; q_2) \in L^1(0, \infty; \mathcal{L}(Z))$  and  $A(q_1)T(\cdot; q_2) \in L^\infty(\eta, \infty; \mathcal{L}(Z))$ , for each  $\eta > 0$ .

iii)  $T(\cdot; q_2) \in L^1(0, \infty; \mathcal{L}(Z, Z_{q_1, \delta}))$  and  $T(\cdot; q_2) \in L^\infty(\eta, \infty; \mathcal{L}(Z; Z_{q_1, \delta}))$ , for each  $\eta > 0$ .

**Note:** This result implies that the operator  $A(q_1)T(t; q_2)$  is bounded for each  $t > 0$ . However, no uniform bound can be found for  $t$  near zero. For  $q_1 = q_2 = q$ , it implies, in particular, that the derivative  $\frac{d}{dt}T(t; q)$  of the solution operator of the homogeneous equation associated with  $(\mathcal{P})_q$  is integrable near  $t = 0$ .

We will also assume that  $A(q)$  satisfies the following hypothesis:

**H4:** For  $\delta$  as in H3 and for any  $q_1, q_2 \in Q_{ad}$  there are constants  $M(q_1, q_2)$  and  $C(q_1, q_2)$  both depending on  $q_1$  and  $q_2$ , such that  $\|(-A(q_1))^\delta (-A(q_2))^{-\delta}\|_{\mathcal{L}(Z)} \leq M(q_1, q_2)$ ,  $\|A(q_1)[A(q_2)]^{-1} - I\| \leq C(q_1, q_2)$  and  $C(q_1, q_2) \rightarrow 0$  as  $q_1 \rightarrow q_2$ .

**Note:** It is sufficient to request that H4 be true for  $\delta = 1$ .

We also consider the hypothesis:

**H4':** For each  $q_0 \in Q_{ad}$  there exists  $C = C(q_0)$  such that

$$\|(A(q) - A(q_0))z\| \leq C\|q - q_0\| \|A(q_0)z\| \quad z \in D, \quad q \in Q_{ad}.$$

THEOREM 2: Assume H1-H4 hold. Then for any  $q_0 \in Q_{ad}$  and  $\varepsilon > 0$ , there exists  $\tilde{\delta} > 0$  such that

$$\|A(q)T(\cdot, q_0)z - A(q_0)T(\cdot, q_0)z\|_{L^1(0, \infty; Z)} \leq \varepsilon \|z\|$$

for all  $z \in Z$ , and for all  $q \in Q_{ad}$  satisfying  $\|q - q_0\| < \tilde{\delta}$ , that is

$$\|A(q)T(\cdot, q_0) - A(q_0)T(\cdot, q_0)\|_{L^1(0, \infty; \mathcal{L}(Z))} \leq \varepsilon,$$

or equivalently, for every fixed  $q_0 \in Q_{ad}$  the mapping from  $Q$  into  $L^1(0, \infty; \mathcal{L}(Z))$  defined by

$$q \rightarrow A(q)T(\cdot, q_0)$$

is continuous on  $Q_{ad}$ .

The proof follows immediately using Lemma 1.

### 3. MAIN RESULTS

Recall that for  $z_0 \in Z_\delta$ ,  $z(t; q)$  satisfies

$$z(t; q) = T(t; q)z_0 + \int_0^t T(t-s; q)F(q, s, z(s; q))ds \doteq T(t; q)z_0 + S(t; q), \quad t \in [0, T].$$

Consider now the following standing hypothesis concerning the  $q$ -regularity of  $\frac{d}{dt}T(t; q)$ .

**H5:** The mapping  $q \rightarrow A(q)T(\cdot; q_0)$  from  $Q$  into  $L^1(0, \infty; \mathcal{L}(Z))$  is Fréchet differentiable at  $q_0$  for all  $q_0 \in Q_{ad}$  (under H1-H4, we already know that this mapping is continuous, by virtue of Theorem 2).

**THEOREM 3:** *Suppose H1-H5 hold. It follows that*

i) *The mapping  $q \rightarrow T(\cdot; q)$  from  $Q \rightarrow L^\infty(0, \infty; \mathcal{L}(Z))$  is Fréchet differentiable at  $q_0$ , for each  $q_0 \in Q_{ad}$ . Moreover, for any  $t > 0$  and  $h \in Q_{ad}$  the  $q$ -Fréchet derivative of  $T(t; q)$  evaluated at  $q_0 \in Q_{ad}$  and applied to  $h \in Q$ , i.e.  $[D_q T(t; q_0)]h$ , is the solution  $v_h(t)$  of the following linear IVP, the so called “sensitivity equation” for  $T(t; q)$ , in  $\mathcal{L}(Z)$*

$$(S_1) : \begin{cases} \frac{d}{dt}v_h(t) = A(q_0)v_h(t) + [D_q A(q)T(t; q_0)|_{q=q_0}]h \\ v_h(0) = 0, \end{cases}$$

and ii) for every  $q_0 \in Q_{ad}$ ,  $D_q T(\cdot; q_0) = D_q T(\cdot; q)|_{q=q_0} \in L^\infty(0, \infty; \mathcal{L}(Q; \mathcal{L}(Z)))$ .

**PROOF:** For  $q_0 \in Q_{ad}$  we have

$$(2) \quad [D_q T(t; q_0)z_0](\cdot) = \int_0^t T(t-s; q_0) [D_q A(q)T(s; q_0)z_0|_{q=q_0}](\cdot) ds.$$

It remains to show the Fréchet differentiability of the mapping  $q \rightarrow T(\cdot; q)$  when viewed as a mapping from  $Q$  into  $L^\infty(0, \infty; \mathcal{L}(Z))$ , i.e. in the stronger  $L^\infty(0, \infty; \mathcal{L}(Z))$  norm. Let  $\varepsilon > 0$ ,  $t > 0$  and  $q_0 \in Q_{ad}$ . First note that for any  $h \in Q$  with  $\|h\| < \tilde{\delta}$ , ( $\tilde{\delta}$  as in Theorem 2) we have

$$\begin{aligned} \frac{d}{dt}[T(t; q_0 + h)z_0 - T(t; q_0)z_0] &= A(q_0 + h)T(t; q_0 + h)z_0 - A(q_0)T(t; q_0)z_0 \\ &= A(q_0 + h)[T(t; q_0 + h)z_0 - T(t; q_0)z_0] + (A(q_0 + h) - A(q_0))T(t; q_0)z_0. \end{aligned}$$

From Theorem 2 and [13] (Corollary 2.2) it follows that

$$(3) \quad T(t; q_0 + h)z_0 - T(t; q_0)z_0 = \int_0^t T(t-s; q_0 + h) (A(q_0 + h) - A(q_0))T(s; q_0)z_0 ds.$$

and therefore for all  $h \in Q$  with  $\|h\| < \tilde{\delta}$ , we have

$$\begin{aligned}
\|T(t; q_0 + h)z_0 - T(t; q_0)z_0\|_Z &\leq \int_0^t M_{q_0+h} e^{-\varepsilon_0(t-s)} \|(A(q_0 + h)T(s; q_0) - A(q_0)T(s; q_0))z_0\|_Z ds \\
&\leq C \|(A(q_0 + h)T(\cdot; q_0) - A(q_0)T(\cdot; q_0))z_0\|_{L^1(0, \infty; Z)} \\
&\leq C\varepsilon \|z_0\|_Z,
\end{aligned}$$

Thus for  $t > 0$

$$(4) \quad \|T(t; q_0 + h) - T(t; q_0)\|_{\mathcal{L}(Z)} \leq C\varepsilon, \quad \text{for } \|h\| < \tilde{\delta},$$

and, since the constant  $C$  above does not depend on  $t$ ,

$$\|T(\cdot; q_0 + h) - T(\cdot; q_0)\|_{L^\infty(0, \infty; \mathcal{L}(Z))} \leq C\varepsilon, \quad \text{for } \|h\| < \tilde{\delta}.$$

The following estimate then follows

$$\begin{aligned}
&\left\| T(t; q_0 + h) - T(t; q_0) - \int_0^t T(t-s; q_0) [D_q A(q)T(s; q_0)|_{q=q_0}] h ds \right\|_{\mathcal{L}(Z)} \\
&\leq (\varepsilon + 1)C \|A(q_0 + h)T(\cdot; q_0) - A(q_0)T(\cdot; q_0) - [D_q A(q)T(\cdot; q_0)|_{q=q_0}] h\|_{L^1(0, \infty; \mathcal{L}(Z))} \\
(5) \quad &+ \varepsilon C \|[D_q A(q)T(\cdot; q_0)|_{q=q_0}] h\|_{L^1(0, \infty; \mathcal{L}(Z))}.
\end{aligned}$$

Now by (H5) for the given  $\varepsilon > 0$  there exists  $\xi > 0$  such that

$$(6) \quad \|A(q_0 + h)T(\cdot; q_0) - A(q_0)T(\cdot; q_0) - [D_q A(q)T(\cdot; q_0)|_{q=q_0}] h\|_{L^1(0, \infty; \mathcal{L}(Z))} \leq \varepsilon \|h\|$$

for  $\|h\| < \xi$ , and since  $D_q A(q)T(\cdot; q_0)|_{q=q_0} \in \mathcal{L}(Q, L^1(0, \infty; \mathcal{L}(Z)))$  there exists  $M, 0 < M < \infty$  such that

$$(7) \quad \|D_q A(q)T(\cdot, q_0)|_{q=q_0}\|_{\mathcal{L}(Q, L^1(0, \infty, \mathcal{L}(Z)))} \leq M.$$

Now, employing (6) and (7) in (5) we get that for  $\|h\| < \min(\tilde{\delta}, \xi)$

$$\begin{aligned}
&\left\| T(t; q_0 + h) - T(t; q_0) - \int_0^t T(t-s; q_0) [D_q A(q)T(s; q_0)|_{q=q_0}] h ds \right\|_{\mathcal{L}(Z)} \\
&\leq (\varepsilon + 1)C\varepsilon \|h\| + \varepsilon CM \|h\| \leq K\varepsilon \|h\|.
\end{aligned}$$

Hence the mapping from  $Q$  into  $L^\infty(0, \infty; \mathcal{L}(Z))$  defined by

$$q \rightarrow T(\cdot; q)$$

is Fréchet  $q$ -differentiable at  $q_0$  and

$$(8) \quad [D_q T(t; q_0)](\cdot) = \int_0^t T(t-s; q_0) [D_q A(q)T(s; q_0)|_{q=q_0}](\cdot) ds.$$

Since  $q_0 \in Q_{ad}$  is arbitrary, part (i) of the Theorem follows. To prove (ii) we first note that by H5, for  $q_0 \in Q_{ad}$ , there exists  $C = C(q_0)$  such that for  $h \in Q$

$$(9) \quad \|D_q A(q)T(\cdot; q_0)|_{q=q_0} h\|_{L^1(0, \infty; \mathcal{L}(Z))} \leq C(q_0) \|h\|.$$

Now, it follows from (8) that for  $t > 0$ ,  $q_0 \in Q_{ad}$  and  $h \in Q$ , one has  $\| [D_q T(t; q_0)] h \|_{\mathcal{L}(Z)} \leq \tilde{C}(q_0) \|h\|$ . Thus  $\|D_q T(t; q_0)\|_{\mathcal{L}(Q; \mathcal{L}(Z))} \leq \tilde{C}(q_0)$ , and since  $\tilde{C}(q_0)$  does not depend on  $t > 0$ , it follows that  $D_q T(\cdot; q_0) \in L^\infty(0, \infty; \mathcal{L}(Q; \mathcal{L}(Z)))$ . ■

Under slightly stronger assumptions on the mapping  $q \rightarrow A(q)T(\cdot; q_0)$ , it is possible to obtain the Lipschitz continuity of the mapping  $q \rightarrow D_q T(\cdot; q_0)$  as a mapping from  $Q$  into  $L^\infty(0, \infty; \mathcal{L}(Q; \mathcal{L}(Z)))$  and from  $Q$  into  $L^\infty(0, \infty; \mathcal{L}(Q; \mathcal{L}(Z, Z_\delta)))$ . In fact, consider the following hypothesis.

**H6:** The mapping  $q \rightarrow D_q A(q)T(\cdot; q_0)$  from  $Q$  into  $L^1(0, \infty; \mathcal{L}(Q; \mathcal{L}(Z)))$  is locally Lipschitz continuous at  $q_0$ , for all  $q_0 \in Q_{ad}$ .

**THEOREM 4:** *Let  $q_0 \in Q_{ad}$  and assume hypotheses H1-H6 hold. Then the mapping  $q \rightarrow D_q T(\cdot; q_0)$  from  $Q$  into  $L^\infty(0, \infty; \mathcal{L}(Q; \mathcal{L}(Z)))$  is locally Lipschitz continuous at  $q_0$ .*

**PROOF:** Choose  $h \in Q$  such that  $\|h\| < \tilde{\delta}$  ( $\tilde{\delta}$  as in Theorem 2) and denote  $G_q(t; q_0)(\cdot) = D_q A(q)T(t; q_0)|_{q=q_0}(\cdot) \in \mathcal{L}(Q, \mathcal{L}(Z))$ . Theorem 3 together with the appropriate choice of  $\alpha(h)$ ,  $0 \leq |\alpha(h)| \leq 1$ , yield

$$\begin{aligned} & \|D_q T(t; q_0 + h)(\cdot) - D_q T(t; q_0)(\cdot)\|_{\mathcal{L}(Q; \mathcal{L}(Z))} \\ & \leq M_{q_0+h} \|G_q(\cdot; q_0 + h) - G_q(\cdot; q_0)\|_{L^1(0, \infty; \mathcal{L}(Q; \mathcal{L}(Z)))} \\ & \quad + \|D_q T(\cdot; q_0 + \alpha(h)h)\|_{L^\infty(0, \infty; \mathcal{L}(Q; \mathcal{L}(Z)))} \|G_q(\cdot; q_0)\|_{L^1(0, \infty; \mathcal{L}(Q; \mathcal{L}(Z)))} \|h\| \\ & \leq C \|h\|. \end{aligned}$$

The last inequality follows from H6 and Theorem 3(ii), and by the fact that  $G_q(\cdot, q_0) \in L^1(0, \infty; \mathcal{L}(Q; \mathcal{L}(Z)))$ , which is a result of H6. ■

In order to obtain the  $q$ -Fréchet differentiability of  $S(\cdot; q)$ , we will need the local Lipschitz continuity of the mapping  $q \rightarrow D_q T(\cdot; q_0)$  when viewed as a mapping from  $Q$  into  $L^\infty(0, \infty; \mathcal{L}(Q; \mathcal{L}(Z; Z_\delta)))$ . This can be achieved by requiring the following hypothesis.

**H7:** For every  $q_0 \in Q_{ad}$ ,  $D_q A(q)T(\cdot; q_0)|_{q=q_0} \in L^1(0, \infty; \mathcal{L}(Q; \mathcal{L}(Z; Z_\delta)))$  and the mapping  $q \rightarrow D_q A(q)T(\cdot; q_0)$  from  $Q$  into  $L^1(0, \infty; \mathcal{L}(Q; \mathcal{L}(Z; Z_\delta)))$  is locally Lipschitz continuous at  $q_0$ , for all  $q_0 \in Q_{ad}$ .

Clearly H7 implies H6 (since the  $Z_\delta$ -norm is stronger than the  $Z$ -norm).

**THEOREM 5:** *Assume H1-H5 and H7 hold. Then, for all  $q_0 \in Q_{ad}$ ,  $D_q T(\cdot; q_0) \in L^\infty(0, \infty; \mathcal{L}(Q; \mathcal{L}(Z; Z_\delta)))$  and, the mapping  $q \rightarrow D_q T(\cdot; q)$  from  $Q$  into the space  $L^\infty(0, \infty; \mathcal{L}(Q; \mathcal{L}(Z; Z_\delta)))$  is locally Lipschitz continuous at  $q_0$ .*

**PROOF:** For  $t > 0$ ,  $z \in Z$ ,  $h \in Q$ , it follows that

$$\begin{aligned} & \| [D_q T(t; q_0)h] z \|_{Z_\delta} = \left\| (-A(q_0))^\delta ([D_q T(t; q_0)] h) z \right\|_Z \\ & = M_{q_0} \|h\| \|z\|_Z \|D_q A(q)T(\cdot; q_0)|_{q=q_0}\|_{L^1(0, t; \mathcal{L}(Q; \mathcal{L}(Z; Z_\delta)))} \\ & \leq C(q_0) \|h\| \|z\|_Z \quad (\text{by virtue of H7}) \end{aligned}$$

Hence,  $\|D_q T(t; q_0)h\|_{\mathcal{L}(Z; Z_\delta)} \leq C(q_0) \|h\|$ , and  $\|D_q T(t; q_0)\|_{\mathcal{L}(Q; \mathcal{L}(Z; Z_\delta))} \leq C(q_0)$ .

Since  $C(q_0)$  does not depend on  $t > 0$ , it follows that  $D_q T(\cdot; q_0) \in L^\infty(0, \infty; \mathcal{L}(Q; \mathcal{L}(Z; Z_\delta)))$ . The Lipschitz continuity of this mapping is obtained following the same steps as in Theorem 4. ■

This result implies that  $q \rightarrow T(\cdot; q)$  is Fréchet differentiable as a mapping from  $Q$  into  $L^\infty(0, \infty; \mathcal{L}(Q; \mathcal{L}(Z; Z_\delta)))$ . In fact, THEOREM 6: *Under the same hypotheses of Theorem 5,  $T(\cdot; q)$  is Fréchet differentiable at  $q_0$ , for each  $q_0 \in Q_{ad}$ , when viewed as a mapping from  $Q$  into  $L^\infty(0, \infty; \mathcal{L}(Z; Z_\delta))$ .*

PROOF: For  $h \in Q$  with  $\|h\| < \tilde{\delta}$  so that  $q_0 + \alpha h \in Q_{ad}$ ,  $\alpha$  satisfying  $|\alpha| \leq 1$ ,  $\beta(h)$  appropriately chosen,  $0 \leq |\beta(h)| \leq 1$ , and any  $t > 0$  it follows that

$$\begin{aligned} & \|T(t; q_0 + h) - T(t; q_0) - [D_q T(t; q_0)] h\|_{\mathcal{L}(Z; Z_\delta)} \\ & \leq C(q_0) \|\beta(h)h\| \|h\| \\ & \leq C(q_0) \epsilon \|h\|, \quad \text{for } \|h\| < \epsilon, \text{ for all } \epsilon \text{ such that } 0 < \epsilon \leq \tilde{\delta}. \end{aligned}$$

■

Note that Theorems 3 and 6 imply that the solution  $z_h(t; q)$  of the linear homogeneous problem associated to  $(\mathcal{P})_q$  is Fréchet differentiable with respect to  $q$ , both as a mapping into  $Z$  and into  $Z_\delta$ , respectively. Theorems 4 and 5 imply, moreover, that the corresponding Fréchet derivatives are locally Lipschitz continuous.

The following generalization of Growall's Lemma for singular kernels will be needed later. Its proof can be found in [6], Lemma 7.1.1.

LEMMA 7: *Let  $L, T, \delta$  be positive constants,  $\delta < 1$ ,  $a(t)$  a real valued, nonnegative, locally integrable function on  $[0, T]$  and  $\mu(t)$  a real-valued function on  $[0, T]$  satisfying*

$$\mu(t) \leq a(t) + L \int_0^t \frac{\mu(s)}{(t-s)^\delta} ds, \quad t \in [0, T].$$

*Then, there exists a constant  $K$  depending only on  $\delta$  such that*

$$\mu(t) \leq a(t) + KL \int_0^t \frac{a(s)}{(t-s)^\delta} ds, \quad t \in [0, T].$$

Before proving the Fréchet differentiability of the mapping  $q \rightarrow S(\cdot; q)$  from  $Q \rightarrow L^\infty(0, T; Z_\delta)$ , we will show that if  $F(q, t, z)$  satisfies appropriate regularity properties, such a mapping is locally Lipschitz continuous at  $q_0$ , for all  $q_0 \in Q_{ad}$ . This result will be needed later.

Consider the hypothesis:

**H8:** The mapping  $q \rightarrow F(q, \cdot; z)$  from  $Q$  into  $L^\infty(0, T; Z)$  is locally Lipschitz continuous for all  $z \in Z_\delta$  with Lipschitz constant independent of  $z$  on  $Z_\delta$ -bounded sets.

THEOREM 8: *Let  $q_0 \in Q_{ad}$ ,  $z_0 \in D_\delta$  and assume H1-H5, H7 and H8 hold. Then the mapping  $q \rightarrow S(\cdot; q)$  from  $Q \rightarrow L^\infty(0, T; Z_\delta)$  is locally Lipschitz continuous at  $q_0$ .*

PROOF: Using Theorem 3 we write

$$\begin{aligned}
S(t; q_0 + h) - S(t; q_0) &= \\
&= \int_0^t T(t-s; q_0 + h) [F(q_0 + h, s, z(s; q_0 + h)) - F(q_0, s, z(s; q_0 + h))] ds \\
&\quad + \int_0^t T(t-s; q_0 + h) [F(q_0, s, z(s; q_0 + h)) - F(q_0, s, z(s; q_0))] ds \\
&\quad + \int_0^t D_q T(t-s; q_0 + \beta(h)h) h F(q_0, s, z(s; q_0)) ds,
\end{aligned}$$

where  $q_0 + h \in Q_{ad}$  for all  $\|h\| \leq \gamma_1$  and  $\beta(h)$  is an appropriately selected constant satisfying  $0 \leq |\beta(h)| \leq 1$ .

Using H8, H3 and Theorem 5 it then follows that

$$\begin{aligned}
&\|S(t; q_0 + h) - S(t; q_0)\|_\delta \\
&\leq \int_0^t \frac{M_{q_0+h} e^{-\varepsilon_0(t-s)}}{(t-s)^\delta} C_1 \|h\| ds + \int_0^t \frac{M_{q_0+h} e^{-\varepsilon_0(t-s)}}{(t-s)^\delta} L \|z(s; q_0 + h) - z(s; q_0)\|_\delta + C_2 \|h\| \\
&\leq C_3 \|h\| + C_4 \int_0^t \frac{\| [D_q T(s; q_0 + \beta(h)h) h] z_0 + S(s; q_0 + h) - S(s; q_0) \|_\delta}{(t-s)^\delta} ds \\
&\leq C_5 \|h\| + C_4 \int_0^t \frac{\|S(s; q_0 + h) - S(s; q_0)\|_\delta}{(t-s)^\delta} ds.
\end{aligned}$$

Hence, by Lemma 7, there exist a constant  $K$  such that

$$\|S(t; q_0 + h) - S(t; q_0)\|_\delta \leq C_5 \|h\| + KC_4 C_5 \|h\| \int_0^T \frac{1}{(t-s)^\delta} ds \doteq C_6 \|h\|, \quad t \in [0, T],$$

provided  $\|h\| \leq \gamma_1$ . The Theorem follows.  $\blacksquare$

It is appropriate to note at this point that this result together with Theorem 6 imply that the mapping  $q \rightarrow z(\cdot; q)$  from  $Q$  into  $L^\infty(0, T; Z_\delta)$  is locally Lipschitz continuous at  $q_0$ . We proceed now to prove the Fréchet differentiability of the mapping  $q \rightarrow S(t; q)$ , corresponding to the nonlinear part of problem  $(\mathcal{P})_q$ . For that purpose, we consider the following hypothesis.

**H9:** The mapping  $(q, z(\cdot)) \rightarrow F(q, \cdot, z(\cdot))$  from  $Q_{ad} \times L^1(0, T; Z_\delta)$  into  $L^\infty(0, T; Z)$  is Fréchet differentiable in both variables, the mapping  $(q, z(\cdot)) \rightarrow F_q(q, \cdot, z(\cdot))$  from  $Q \times L^\infty(0, T; Z_\delta)$  into  $L^\infty(0, T; \mathcal{L}(Q; Z_\delta))$  is locally Lipschitz continuous with respect to  $q$  and  $z$ , with Lipschitz constant independent of  $z$  on  $Z_\delta$ -bounded sets and  $F_z(q, \cdot, z(\cdot; q)) \in L^\infty(0, T; \mathcal{L}(Z; Z_\delta))$ .

Clearly H9 is stronger than H8.

**THEOREM 9:** *Let  $q_0 \in Q_{ad}$ ,  $z_0 \in D_\delta$  and suppose H1-H5, H7 and H9 hold. Then the mapping  $q \rightarrow S(t; q) = \int_0^t T(t-s; q) F(q, s, z(s; q)) ds$  from  $Q \rightarrow L^\infty(0, T; Z_\delta)$  is Fréchet differentiable at  $q_0$ . Moreover, for any  $t \in [0, T]$ , and any  $h \in Q_{ad}$ ,  $[D_q S(t; q_0)]h \doteq w_h(t)$  satisfies the integral equation*

$$\begin{aligned}
w_h(t) = \int_0^t \left\{ T(t-s; q_0) \left[ F_q(q_0, s, z(s; q_0))h + F_z(q_0, s, z(s; q_0)) [D_q T(s; q_0)z_0]h \right. \right. \\
(10) \qquad \qquad \qquad \left. \left. + F_z(q_0, s, z(s; q_0))w_h(s) \right] + [D_q T(t-s; q_0)F(q_0, s, z(s; q_0))] h \right\} ds,
\end{aligned}$$

and  $w_h(t)$  is the solution of the following non-homogeneous linear IVP, the so-called “sensitivity equation” for  $S(t; q)$ , in  $Z$ :

$$(S_2) \begin{cases} \frac{d}{dt} w_h(t) = (A(q_0) + F_z(q_0, t, z(t; q_0)))w_h(t) + F_q(q_0, t, z(t; q_0))h + \\ \qquad + F_z(q_0, t, z(t; q_0))[D_q T(t; q_0)z_0]h + \int_0^t D_q A(q)T(t-s; q_0)|_{q=q_0} h F(q_0, s, z(s; q_0)) ds \\ w_h(0) = 0. \end{cases}$$

PROOF: That the solution  $w_h(t)$  of  $(S_2)$  satisfies (10) follows immediately  $(S_1)$  in Theorem 2 and the fact that  $[D_q T(0; q_0)z]h = 0$  for  $z \in Z$  and  $h \in Q$ .

We write

$$\begin{aligned}
S(t; q_0 + h) - S(t; q_0) - w_h(t) &= \\
&= \int_0^t T(t-s; q_0) [F(q_0 + h, s, z(s; q_0)) - F(q_0, s, z(s; q_0)) - F_q(q_0, s, z(s; q_0))h] ds \\
&\quad + \int_0^t T(t-s; q_0) \left[ F(q_0, s, z(s; q_0 + h)) - F(q_0, s, z(s; q_0)) \right. \\
&\qquad \qquad \qquad \left. - F_z(q_0, s, z(s; q_0))(z(s; q_0 + h) - z(s; q_0)) \right] ds \\
&\quad + \int_0^t T(t-s; q_0) F_z(q_0, s, z(s; q_0)) [S(s; q_0 + h) - S(s; q_0) - w_h(s)] ds \\
&\quad + \int_0^t T(t-s; q_0) F_z(q_0, z(s; q_0)) \left[ [D_q T(s; q_0 + \alpha(h)h)z_0]h - [D_q T(s; q_0)z_0]h \right] ds \\
&\quad + \int_0^t \left\{ T(t-s; q_0 + h)F(q_0, s, z(s; q_0)) - T(t-s; q_0)F(q_0, s, z(s; q_0)) \right. \\
&\qquad \qquad \qquad \left. - [D_q T(t-s; q_0)F(q_0, s, z(s; q_0))]h \right\} ds \\
&\quad + \int_0^t T(t-s; q_0 + h) [F(q_0 + h, s, z(s; q_0 + h)) - F(q_0, s, z(s; q_0))] ds \\
&\quad - \int_0^t T(t-s; q_0) [F(q_0 + h, s, z(s; q_0)) - 2F(q_0, s, z(s; q_0)) + F(q_0, s, z(s; q_0 + h))] ds \\
&\doteq \sum_{i=1}^7 I_i,
\end{aligned}$$

where  $I_i$  is the  $i^{th}$  term in the expression written above. Here,  $\alpha(h)$  is an appropriately chosen constant satisfying  $0 \leq |\alpha(h)| \leq 1$ .

In what follows,  $C_i$  will denote a generic finite positive constant depending on  $q_0$ . Let  $\gamma_1 > 0$  be such that  $q_0 + \eta \in Q_{ad}$  for all  $\eta \in Q$  satisfying  $\|\eta\| < \gamma_1$ . Then for any  $h \in Q_{ad}$

with  $\|h\| < \gamma_1$  it follows, by virtue of Theorem 8 and hypothesis H9 that there exist positive constants  $C_1$ ,  $C_2$  and  $L$ , such that:

$$\begin{aligned} \|I_6 + I_7\|_\delta &\leq C_1 \|h\|^2 + \int_0^t \frac{L}{(t-s)^\delta} \left( |\alpha_1(h) - \alpha_3(h)| \|h\| + \|z(s; q_0 + h) - z(s; q_0)\|_\delta \right) \|h\| ds \\ &\quad + \int_0^t \frac{C_2}{(t-s)^\delta} \|z(s; q_0 + h) - z(s; q_0)\|_\delta \|h\| ds \\ (11) \quad &\leq C_3 \|h\|^2, \quad \text{provided } \|h\| \leq \gamma_1, \end{aligned}$$

where the last inequality follows from the Lipschitz continuity of the mapping  $q \rightarrow z(\cdot; q)$  from  $Q$  into  $L^\infty(0, T; Z_\delta)$  at  $q_0$ . Now let  $\varepsilon$  be a fixed positive constant. It follows from H9 that there exist  $\gamma_2 > 0$  and  $\gamma_3 > 0$  such that

$$(12) \quad \|I_1\|_\delta \leq \int_0^t \frac{C_4}{(t-s)^\delta} \varepsilon \|h\| ds \leq C_5 \varepsilon \|h\|,$$

provided  $\|h\| \leq \gamma_2$ , and also

$$\begin{aligned} \|I_2\|_\delta &\leq \int_0^t \frac{C_6 \varepsilon}{(t-s)^\delta} \|z(s; q_0 + h) - z(s; q_0)\|_Z ds \\ (13) \quad &\leq C_7 \varepsilon \|h\|, \quad \text{provided } \|h\| \leq \gamma_3. \end{aligned}$$

Also, by using H9 we have that  $F_z(q_0, \cdot, z(\cdot; q_0)) \in L^\infty(0, T; \mathcal{L}(Z; Z_\delta))$ , and therefore there exists a constant  $C_8$  such that

$$(14) \quad \|I_3\|_\delta \leq C_8 \int_0^t \frac{\|S(s; q_0 + h) - S(s; q_0) - w_h(s)\|_\delta}{(t-s)^\delta} ds.$$

On the other hand, the local Lipschitz continuity of  $D_q T(\cdot; q_0)$  (Theorem 4), implies the existence of two finite positive constants  $C_9$  and  $\gamma_4$  such that

$$(15) \quad \|I_4\|_\delta \leq \int_0^t \frac{C_9}{(t-s)^\delta} |\alpha(h)| \|h\|^2 ds \leq C_{10} \|h\|^2, \quad \text{provided } \|h\| \leq \gamma_4.$$

Finally from Theorem 6 and H9, there are two finite positive constants  $C_{10}$  and  $\gamma_5$  such that

$$(16) \quad \|I_5\|_\delta \leq C(q_0) \varepsilon \|h\| \int_0^t \|F(q_0, s, z(s; q_0))\|_Z ds \leq C_{10} \varepsilon \|h\|,$$

provided  $\|h\| \leq \gamma_5$ .

From (11)-(16) we conclude that there exist finite positive constants  $C_{11}$ ,  $C_{12}$ , and  $\gamma$  such that for  $t \in [0, T]$  and  $h \in Q_{ad}$  with  $\|h\| \leq \gamma$

$$\begin{aligned} \|S(t; q_0 + h) - S(t; q_0) - w_h(t)\|_\delta &\leq C_{11} \varepsilon \|h\| + \\ &\quad + C_{12} \int_0^t \frac{\|S(s; q_0 + h) - S(s; q_0) - w_h(s)\|_\delta}{(t-s)^\delta} ds. \end{aligned}$$

Hence, applying Lemma 7 we conclude that

$$\begin{aligned} \|S(t; q_0 + h) - S(t; q_0) - w_h(t)\|_\delta &\leq C_{11} \varepsilon \|h\| + KC_{12}C_{11}\varepsilon \|h\| \int_0^t \frac{1}{(t-s)^\delta} ds \\ &\leq C_{13}\varepsilon \|h\|, \quad t \in [0, T], \quad \|h\| \leq \gamma. \end{aligned}$$

hence the mapping  $q \rightarrow S(\cdot; q)$  from  $Q \rightarrow L^\infty(0, T; Z_\delta)$  is Fréchet differentiable at  $q_0$  and  $w_h(t)$  is the Fréchet derivative of  $S(t; q)$  at  $q_0$ , i.e.  $D_q S(t; q_0) = w_h(t)$ . ■

**THEOREM 10:** *Under the same hypotheses of Theorem 9, the mapping  $q \rightarrow z(\cdot; q)$  from the admissible parameter set  $Q_{ad}$  into the solution space  $L^\infty(0, T; Z_\delta)$ , is Fréchet differentiable at  $q_0$ . Moreover, for any  $h \in Q$ ,  $t \in [0, T]$ , the  $q$ -Fréchet derivative of  $z(t; q)$  evaluated at  $q_0$  and applied to  $h$ , i.e.  $[D_q z(t; q_0)]h$  is the solution  $v_h(t)$  of the following linear non-homogeneous initial value problem in  $Z$ , the sensitivity equation for  $z(t; q)$*

$$(S) \begin{cases} \frac{d}{dt} v_h(t) = (A(q_0) + F_z(q_0, t, z(t; q_0))) v_h(t) + F_q(q_0, t, z(t; q_0))h + \\ \quad + D_q A(q)T(t; q_0)z_0|_{q=q_0} h + \int_0^t D_q A(q)T(t-s; q_0)|_{q=q_0} h F(q_0, s, z(s; q_0)) ds \\ v_h(0) = 0. \end{cases}$$

**PROOF:** The Fréchet differentiability of  $z(t; q) = T(t; q)z_0 + S(t; q)$  follows immediately from Theorems 6 and 9 and the sensitivity equation is readily obtained by combining the sensitivity equations  $(S_1)$  and  $(S_2)$ . ■

#### REFERENCES

- [1] BREWER, D., *The Differentiability with Respect to a Parameter of the Solution of a Linear Abstract Cauchy Problem*, SIAM Journal on Mathematical Analysis, Vol. 13, N 4, 1982, pp. 607-620.
- [2] BREWER, D., BURNS, J. AND CLIFF E., *Parameter Identification for an Abstract Cauchy Problem by Quasilinearization*, Quarterly of Applied Mathematics, Vol 51, 1993, pp. 1-22.
- [3] BURNS, J., MORIN P. AND SPIES R., *Parameter Differentiability of the Solution of a Nonlinear Abstract Cauchy Problem*, Journal of Math. Analysis and Applications, Vol. 252, 2000, pp. 18-31.
- [4] CLARK, L. G. AND GIBSON, J. S., *Sensitivity Analysis for a Class of Evolution Equations*, Journal of Mathematical Analysis and Applications, Vol. 58, 1977, pp. 22-31.
- [5] HAMMER, P. W., *Parameter Identification in Parabolic Partial Differential Equations Using Quasilinearization*, PhD thesis, ICAM Report 90-07-01, Virginia Tech, 1990.
- [6] HENRY, D., *Geometric Theory of Semilinear Parabolic Equations*, Lecture Notes in Mathematics, 840, Springer-Verlag, 1989.
- [7] HERDMAN T., MORIN P. AND SPIES R., *Parameter Identification for Nonlinear Abstract Cauchy Problems Using Quasilinearization*, J. Optimization Theory and Appl., 113,2, 2002, pp. 227-250.
- [8] LUNARDI, A., *Analytic Semigroups and Optimal Regularity in Parabolic Problems*, Birkhäuser, 1995.
- [9] MORIN, P. AND SPIES, R., *Identifiability of the Landau-Ginzburg Potential in a Mathematical Model of Shape Memory Alloys*, J. Math. Analysis and Appl., 212, 1997, pp. 292-315.
- [10] MORIN, P. AND SPIES, R., *A Quasilinearization Approach for Parameter Identification in a Nonlinear Model of Shape Memory Alloys*, Inverse Problems, Vol. 14, 1998, pp. 1551-1563.
- [11] SPIES, R., *A State-Space Approach to a One-Dimensional Mathematical Model for the Dynamics of Phase Transitions in Pseudoelastic Materials*, Journal of Mathematical Analysis and Applications, Vol. 190, 1995, pp. 58-100.
- [12] SPIES, R., *Results on a Mathematical Model of Thermomechanical Phase Transitions in Shape Memory Materials*, Smart Materials and Structures, Vol. 3, 1994, pp. 459-469.
- [13] PAZY, A., *Semigroups of Linear Operators and Applications to Partial Differential Equations*, Springer-Verlag, 1983.